

Patent
238/168

To: Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

NEW APPLICATION TRANSMITTAL
UTILITY

Sir:

Transmitted herewith for filing is a **utility** patent application:

Inventor(s): Peter B. Dervan, et al.

Title: DESIGN, SYNTHESIS AND USE OF SPECIFIC POLYAMIDE DNA-BINDING LIGANDS

I. PAPERS ENCLOSED HEREWITH FOR FILING UNDER 37 CFR § 1.53(b):

228 Page(s) of Written Description

12 Page(s) Claims

1 Page(s) Abstract

14 Sheets of Drawings ☒ Informal ☐ Formal

II. ADDITIONAL PAPERS ENCLOSED IN CONNECTION WITH THIS FILING:

☒ Declaration

☒ Power of Attorney: ☐ Separate or ☒ Combined with Declaration

☒ Assignment to California Institute of Technology and assignment cover sheet

☒ Verified Statement establishing "**Small Entity**" under 37 CFR §§ 1.9 and 1.27

CERTIFICATE OF MAILING
(37 C.F.R. §1.10)

I hereby certify that this paper (along with any referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as 'Express Mail Post Office To Addressee' in an envelope addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

EL088594213 US
Express Mail Label No.

August 11, 1999
Date of Deposit

AKIL PARRISH
Name of Person Mailing Paper

Akil Parrish
Signature of Person Mailing Paper

- ☐ Priority Document No(s): _____
- ☐ Information Disclosure Statement w/PTO 1449 ☐ Copy of Citations
- ☒ Preliminary Amendment
- ☒ Return Postcard

III. THE FILING FEE HAS BEEN CALCULATED AS SHOWN BELOW:

BASIC FILING FEE:							\$760.00
Total Claims	48	-	20	=	28	x \$18.00	\$504.00
Independent Claims	8	-	3	=	5	x \$78.00	\$390.00
Multiple Dependent Claims	\$260	(if applicable)				<input type="checkbox"/>	\$0.00
TOTAL OF ABOVE CALCULATIONS							\$1,654.00
Reduction by ½ for Filing by Small Entity. Note 37 CFR §§ 1.9, 1.27, 1.28. If applicable, Verified Statement must be attached.							\$827.00
Misc. Filing Fees (Recordation of Assignment -- \$40)							\$40.00
TOTAL FEES SUBMITTED HERewith							\$867.00

IV. METHOD OF PAYMENT OF FEES

- ☒ A check in the amount of \$867.00.
- ☐ Charge Lyon & Lyon's Deposit Account No. **12-2475** in the amount of _____.
- ☐ This application is being filed without fee or Declaration under 37 CFR § 1.53.

V. AUTHORIZATION to CHARGE FEES

The Commissioner is authorized to charge Lyon & Lyon's Deposit Account No. **12-2475** for the following:


- ☒ 37 CFR § 1.16(a), (f) or (g) – **(Filing fees)**
- ☒ 37 CFR § 1.16(b), (c) and (d) – **(Presentation of Extra Claims)**
- ☒ 37 CFR § 1.16(e) – **(Surcharge for filing the basic filing fee and/or Declaration on a date later than the filing date of the application)**
- ☒ 37 CFR § 1.17 – **(Any Application processing fees)**

☒ Credit Lyon & Lyon's Deposit Account No. **12-2475** for overpayment of fees.

Respectfully submitted,

LYON & LYON LLP

Dated: 08/11/99

By: 
Daniel M. Chambers
Reg. No. 34,561

633 West Fifth Street, Suite 4700
Los Angeles, California 90071-2066
(213) 489-1600

Applicant or Patentee: Peter B. Dervan and Eldon E. Baird
Serial or Patent No.: To Be Assigned
Filed or Issued: Herewith
For: DESIGN, SYNTHESIS AND USE OF SPECIFIC POLYAMIDE DNA-BINDING LIGANDS

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(d) and 1.27(c)) - SMALL BUSINESS CONCERN**

I hereby declare that I am:

_____ the owner of the small business concern identified below:
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF CONCERN: GENESOFT, INC.
ADDRESS OF CONCERN: Two Corporate Drive
South San Francisco, California 94080

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR 121.3-18, and reproduced in 37 CFR 1.9(d) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third-party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention entitled DESIGN, SYNTHESIS AND USE OF SPECIFIC POLYAMIDE DNA-BINDING LIGANDS, by inventors Peter B. Dervan and Eldon E. Baird described in

☒ the specification filed herewith.
_____ application serial number _____, filed _____
_____ Patent No. _____, issued _____

If the rights held by the above-identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

**NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)*

NAME: California Institute of Technology
ADDRESS: 1200 East California Blvd., Pasadena, California 91125
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☒ NONPROFIT ORGANIZATION


NAME: Prolinx, Inc.
ADDRESS: 22322 20th Avenue S.E., Bothell, Washington 98021
☐ INDIVIDUAL ☒ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING: David B. Singer
TITLE OF PERSON IN ORGANIZATION: President + CEO
ADDRESS OF PERSON SIGNING: Two Corporate Drive
South San Francisco, CA 94080

Signature



Date

8-6-99

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

Peter B. Dervan et al.

Serial No.:

Filed: August 11, 1999

For: DESIGN, SYNTHESIS AND USE OF
SPECIFIC POLYAMIDE DNA-BINDING
LIGANDS

)
) **Group Art Unit:** ArtUnit

)
) **Examiner:** Examiner

PRELIMINARY AMENDMENT

Box New Application
Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend the application as follows:

In the Specification:

On page 1, line 10, please insert after "This application": --claims priority to
PCT/US98/01714 filed January 29, 1998, and--

Respectfully submitted,

LYON & LYON LLP

By: 

Daniel M. Chambers

Reg. No. 34,561

633 West Fifth Street, Suite 4700
Los Angeles, California 90071-2066
(213) 489-1600

APPLICATION

FOR

UNITED STATES LETTERS PATENT

TO THE ASSISTANT COMMISSIONER OF PATENTS:

BE IT KNOWN THAT WE PETER B. DERVAN AND ELDON J. BAIRD

have invented certain new and useful improvements in

**“DESIGN, SYNTHESIS AND USE OF SPECIFIC POLYAMIDE DNA-BINDING
LIGANDS”**

of which the following is a specification:

DESIGN, SYNTHESIS AND USE OF SPECIFIC
POLYAMIDE DNA-BINDING LIGANDS
(Case.No. 98,016)

5 The U.S. Government has certain rights in this invention pursuant to Grant Nos. GM 26453, 27681 and 47530 awarded by the National Institute of Health.

CROSS REFERENCE TO RELATED APPLICATIONS

10 This application is a continuation-in-part of PCT/US97/03332 filed February 20, 1997, Serial No. 08/853,522 filed May 8, 1997 and PCT/US 97/12722 filed July 21, 1997 which are continuation-in-part applications of Serial No. 08/837,524, filed April 21, 1997, Serial No. 08/607,078, filed February 26, 1996, provisional application Serial No. 60/042,022, filed April 16, 1997 and provisional application Serial No. 60/043,444, filed April 8, 1997.

BACKGROUND OF THE INVENTION

Field of the Invention

This invention relates to polyamides which bind to predetermined sequences in the minor groove of double stranded DNA.

Description of the Related Art

20 The design of synthetic ligands that read the information stored in the DNA double helix has been a long standing goal of chemistry. Cell-permeable small molecules which target predetermined DNA sequences are useful for the regulation of gene-expression. Oligodeoxynucleotides that recognize the major groove of double-helical DNA via triple-helix formation bind to a broad range of sequences with high affinity and specificity. Although
30 oligonucleotides and their analogs have been shown to interfere with gene expression, the triple helix approach is limited to purine tracks and suffers from poor cellular uptake. The development of pairing rules for minor groove binding polyamides derived from N-methylpyrrole (Py) and N-methylimidazole (Im) amino acids provides another code to control sequence specificity. An Im/Py pair distinguishes G•C from C•G and both of these from A•T or
35 T•A base pairs. Wade, W.S., Mrksich, M. & Dervan, P.B. describes the design of peptides that bind in the minor groove of DNA at 5'-(A,T)G(A,T)C(A,T)-3' sequences by a dimeric side-by-side motif. *J. Am. Chem. Soc.* **114**, 8783-8794 (1992); Mrksich, M. *et al.* describes antiparallel

side-by-side motif for sequence specific-recognition in the minor groove of DNA by the designed peptide 1-methylimidazole-2-carboxamidenetropsin. *Proc. Natl. Acad. Sci. USA* **89**, 7586-7590 (1992); Trauger, J.W., Baird, E. E. Dervan, P.B. describes the recognition of DNA by designed ligands at subnanomolar concentrations. *Nature* **382**, 559-561 (1996). A Py/Py pair specifies A•T from G•C but does not distinguish A•T from T•A. Pelton, J.G. & Wemmer, D.E. describes the structural characterization of a 2-1 distamycin A-d(CGCAAATTTGGC) complex by two-dimensional NMR. *Proc. Natl. Acad. Sci. USA* **86**, 5723-5727 (1989); White, S., Baird, E. E. & Dervan, P.B. Describes the effects of the A•T/T•A degeneracy of pyrrole-imidazole polyamide recognition in the minor groove of DNA. *Biochemistry* **35**, 12532-12537 (1996); White, S., Baird, E. E. & Dervan, P. B. describes the pairing rules for recognition in the minor groove of DNA by pyrrole-imidazole polyamides. *Chem. & Biol.* **4**, 569-578 (1997); White, S., Baird, E. E. & Dervan, P.B. describes the 5'-3' N-C orientation preference for polyamide binding in the minor groove. New methods of designing selective compounds and the resulting specific polyamide binding ligands that are designed to target an identified sequence of double stranded DNA are needed to overcome the A•T/T•A degeneracy of pyrrole-imidazole polyamide recognition.

SUMMARY OF THE INVENTION

It has been found that a new aromatic amino acid, 3-hydroxy-N-methylpyrrole (Hp) when incorporated into a polyamide and paired opposite Py, provides the means to discriminate A•T from T•A. Unexpectedly, the replacement of a single hydrogen atom on the pyrrole with a hydroxy group in a Hp/Py pair regulates the affinity and the specificity of a polyamide by an order of magnitude. Utilizing Hp together with Py and Im in polyamides to form four aromatic amino acid pairs (Im/Py, Py/Im, Hp/Py, and Py/Hp) provides a code to distinguish all four Watson-Crick base pairs in the minor groove of DNA.


The present invention provides a method for designing specific polyamides suitable for use as DNA-binding ligands, as well as compositions comprising such polyamides, that are selective for an identified target sequence of double stranded DNA. Preferably, the designed specific polyamides are characterized by a dissociation constant of less than 1 nM, as measured by DNase I footprint titration, and greater than ten-fold selectivity for the identified target

sequence over related mismatch sequences, based on the ratio of the corresponding dissociation constants measured by DNase I footprint titrations.

The invention encompasses improved polyamides for binding to the minor groove of double stranded ("duplex") DNA. The polyamides are in the form of a hairpin comprising two groups of at least three consecutive carboxamide residues, the two groups covalently linked by an aliphatic amino acid residue, preferably γ -aminobutyric acid or 2,4 diaminobutyric acid, the consecutive carboxamide residues of the first group pairing in an antiparallel manner with the consecutive carboxamide residues of the second group in the minor groove of double stranded DNA. The improvement relates to the inclusion of a binding pair of Hp/Py carboxamides in the polyamide to bind to a T•A base pair in the minor groove of double stranded DNA or Py/Hp carboxamide binding pair in the polyamide to bind to an A•T base pair in the minor groove of double stranded DNA. The improved polyamides have at least three consecutive carboxamide pairs for binding to at least three DNA base pairs in the minor groove of a duplex DNA sequence that has at least one A•T or T•A DNA base pair, the improvement comprising selecting a Hp/Py carboxamide pair to correspond to a T•A base pair in the minor groove or a Py/Hp carboxamide pair to bind to an A•T DNA base pair in the minor groove. Preferably the binding of the carboxamide pairs to the DNA base pairs modulates the expression of a gene.

In general, the method provides specific polyamides suitable for use as DNA-binding ligands that are selective for identified target sequences of double stranded DNA having a coding strand sequence of the form 5'-WN₁N₂ . . . N_mW-3' where N is a nucleotide chosen from the group A, T, C and G, W is a nucleotide chosen from the group A and T, and with the corresponding paired antiparallel strand 3'-W'N'₁N'₂ . . . N'_mW'-5' where N' is a nucleotide chosen from the group T, A, G and C respectively to form Watson-Crick base pairs, W is a nucleotide chosen from the group T and A respectively to form Watson-Crick base pairs, and m is an integer having a value from 3 to 6 inclusive.

The preferred corresponding designed specific polyamides resulting from this invention are of the form



wherein X₁, X₂, X_m, X_(m+1), X_(2m-1), and X_{2m} are carboxamide residues forming carboxamide binding pairs X₁/X_{2m}, X₂/X_(2m-1), X_m/X_(m+1), and γ is γ -aminobutyric acid or 2,4 diaminobutyric acid and Dp is dimethylaminopropylamide,

and where

carboxamide binding pair X_1/X_{2m} corresponds to base pair $N_1 \bullet N'_1$,
 carboxamide binding pair $X_2/X_{(2m-1)}$ corresponds to base pair $N_2 \bullet N'_2$,
 carboxamide binding pair $X_m/X_{(m+1)}$ corresponds to base pair $N_m \bullet N'_m$.

5

In general, the specific polyamide DNA-binding ligands were designed by using a method that comprises the steps of identifying the target DNA sequence 5'-WN₁N₂ ... N_mW-3'; representing the identified sequence as 5'-W**a**b ... **x**W-3', wherein **a** is a first nucleotide to be bound by the X₁ carboxamide residue, **b** is a second nucleotide to be bound by the X₂ carboxamide residue, and **x** is the corresponding nucleotide to be bound by the X_m carboxamide residue; defining **a** as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified six base pair sequence.

10

Carboxamide residues were selected sequentially as follows: Im was selected as the X₁ carboxamide residue and Py as the X_{2m} carboxamide residue if **a** was G. Py was selected as the X₁ carboxamide residue and Im as the X_{2m} carboxamide residue if **a** was C. Hp was selected as the X₁ carboxamide residue and Py as the X_{2m} carboxamide residue if **a** was T. Py was selected as the X₁ carboxamide residue and Hp as the X_{2m} carboxamide residue if **a** was A.

20

The remaining carboxamide residues were selected in the same fashion. Im was selected as the X₂ carboxamide residue and Py as the X_{2m-1} carboxamide residue if **b** was G. Py was selected as the X₂ carboxamide residue and Im as the X_{2m-1} carboxamide residue if **b** was C. Hp was selected as the X₂ carboxamide residue and Py as the X_{2m-1} carboxamide residue if **b** was T. Py was selected as the X₂ carboxamide residue and Hp as the X_{2m-1} carboxamide residue if **b** was A.


25

The selection of carboxamide residues was continued through m iterations. In the last iteration, Im was selected as the X_m carboxamide residue and Py as the X_{m+1} carboxamide residue if **x** was G. Py was selected as the X_m carboxamide residue and Im as the X_{m+1} carboxamide residue if **x** was C. Hp was selected as the X_m carboxamide residue and Py as the X_{m+1} carboxamide residue if **x** was T. Py was selected as the X_m carboxamide residue and Hp as the X_{m+1} carboxamide residue if **x** was A.

30

In one preferred embodiment, the polyamide includes at least four consecutive carboxamide pairs for binding to at least four base pairs in a duplex DNA sequence. In another preferred embodiment, the polyamide includes at least five consecutive carboxamide pairs for binding to at least five base pairs in a duplex DNA sequence. In yet another preferred embodiment, the polyamide includes at least six consecutive carboxamide pairs for binding to at least six base pairs in a duplex DNA sequence. In one preferred embodiment, the improved polyamides have four carboxamide binding pairs that will distinguish A•T, T•A, C•G and G•C base pairs in the minor groove of a duplex DNA sequence. The duplex DNA sequence can be a regulatory sequence, such as a promoter sequence or an enhancer sequence, or a gene sequence, such as a coding sequence or a non-coding sequence. Preferably, the duplex DNA sequence is a promoter sequence.

More specifically, "polyamide" refers to a polymer of polyamide subunits of the formula.



where R^1 is chosen from H, NH_2 , SH, Cl, Br, F, N-acetyl, or N-formyl.

where R^2 is C_{1-100} alkyl (preferably C_{1-10} alkyl such as methyl, ethyl, isopropyl), C_{1-100} alkylamine (preferably C_{1-10} alkylamine such as ethylamine), C_{1-100} alkyldiamine (preferably C_{1-10} alkyldiamine such as N,N-dimethylpropylamine), a C_{1-100} alkylcarboxylate (preferably a C_{1-10} alkylcarboxylate such as $-CH_2COOH$), C_{1-100} alkenyl (preferably C_{1-10} alkenyl such as $CH_2CH=CH_2$), or a C_{1-100} alkynyl (preferably C_{1-10} alkynyl such as $-CH_2C\equiv CH_3$), or a $C_{1-100}L$, where L groups can be independently chosen from but is not limited to arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, oligodeoxynucleotide, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine,

captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- α -tocopheral. Most preferably R^2 is H, $(CH_2)_mCH_3$, $(CH_2)_mNH_2$, $(CH_2)_mSH$, $(CH_2)_mOH$, $(CH_2)_mNR^5_2$, $(CH_2)_mOR^5$, $(CH_2)_mSR^5$, where $R^5 = (CH_2)_mCH_3$, $(CH_2)_mNH_2$, $(CH_2)_mSH$, $(CH_2)_mOH$ and m is an integer from 0 to 6.

where R^3 is chosen from H, NH_2 , OH, SH, Br, Cl, F, OMe, CH_2OH , CH_2SH , CH_2NH_2 .

where R^4 is $-NH(CH_2)_{0-100}NR^6R^7$ or $NH(CH_2)_pCO NH(CH_2)_{0-100}NR^6R^7$ or NHR^6 or $NH(CH_2)_pCONHR^6$. Where R^6 and R^7 are independently chosen from H, Cl, NO, N-acetyl, benzyl, C_{1-100} alkyl, C_{1-100} alkylamine, C_{1-100} alkyldiamine, C_{1-100} alkylcarboxylate, C_{1-100} alkenyl, a C_{1-100} alkynyl, or a $C_{1-100}L$, where L groups can be independently chosen from but is not limited to arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, oligodeoxynucleotide, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, an oligodeoxynucleotide, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- α -tocopheral. Where p is an integer value ranging from 0 to 12. In the preferred form of the present invention R^6 and R^7 are H, and the resulting amine modified polyamide is coupled to an amine reactive molecule in order to generate a bifunction polyamide conjugate. Where the amine reactive molecule is chosen from but not limited to the following: arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, an oligodeoxynucleotide, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- α -tocopheral.

where X and Y are chosen from the following, N, CH, COH, CCH_3 , CNH_2 , CCl, CF.

a is an integer chosen from values of 0 or 1

b is an integer chosen integer values ranging from 1 to 5.

c is an integer value ranging from 2 to 10.

Hereinafter, N-methylpyrrolocarboxamide may be referred to as "Py". N-methylimidazolecarboxamide may be referred to as "Im", γ -aminobutyric acid may referred to as " γ ", β -alanine may be referred to as " β ", glycine may be referred to as "G",

dimethylaminopropylamide may be referred to as "Dp", and ethylenediaminetetraacetic acid may be referred to as "EDTA".

5 The preparation and the use of polyamides for binding in the minor groove of double stranded DNA are extensively described in the art. This invention is an improvement of the existing technology that uses 3-hydroxy-N-methylpyrrole to provide carboxamide binding pairs for DNA binding polyamides.

10 The invention encompasses polyamides having γ -aminobutyric acid or a substituted γ -aminobutyric acid to form a hairpin with a member of each carboxamide pairing on each side of it. Preferably the substituted γ -aminobutyric acid is a chiral substituted γ -aminobutyric acid such as (R)-2,4-diaminobutyric acid. In addition, the polyamides may contain an aliphatic amino acid residue, preferably a β -alanine residue, in place of a Hp or Py carboxamide. The β -alanine residue is represented in formulas as β . The β -alanine residue becomes a member of a carboxamide binding pair. The invention further includes the substitution as a β/β binding pair for non-Im containing binding pair. Thus, binding pairs in addition to the Im/Py, Py/Im, Hp/Py and Py/Hp are Im/ β , β /Im, Py/ β , β /Py, Hp/ β , β /Hp, and β/β .

20 The polyamides of the invention can have additional moieties attached covalently to the polyamide. Preferably the additional moieties are attached as substituents at the amino terminus of the polyamide, the carboxy terminus of the polyamide, or at a chiral (R)-2,4-diaminobutyric acid residue. Suitable additional moieties include a detectable labeling group such as a dye, biotin or a hapten. Other suitable additional moieties are DNA reactive moieties that provide for sequence specific cleavage of the duplex DNA.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 illustrates the structure of polyamide 1, 2, and 3.

30 Figure 2 illustrates the pairing of polyamides to DNA base pairs.

Figure 3 illustrates the DNase footprint titration of compounds 2 and 3.

Figure 4 illustrates a list of the structures of representative Hp containing polyamides.

Figure 5 schematically illustrates a method for the design of eight carboxamide residue hairpin polyamide compounds suitable for recognition of 6-bp 5'-WNNNNW-3' sequences in the minor groove of double stranded DNA.

5 Figure 6 schematically illustrates a method for determining the position of an aromatic amino acid residue that should be replaced with a β -alanine residue in order to enhance the DNA binding properties of certain eight carboxamide residue hairpin polyamide compounds.

Figure 7 schematically illustrates a method for the design of ten carboxamide residue hairpin polyamide compounds suitable for recognition of 7-bp 5'-WNNNNW-3' sequences in the
10 minor groove of double stranded DNA.

Figure 8 schematically illustrates a method for determining the position of an aromatic amino acid residue that should be replaced with a β -alanine residue in order to enhance the DNA binding properties of certain ten carboxamide residue hairpin polyamide compounds.

Figure 9 schematically illustrates a method for determining the position of an additional aromatic amino acid residue that should be replaced with a β -alanine residue in order to enhance the DNA binding properties of certain ten carboxamide residue hairpin polyamide compounds.

Figure 10 schematically illustrates a method for the design of twelve carboxamide residue hairpin polyamide compounds suitable for recognition of 8-bp 5'-WNNNNW-3' sequences in the minor groove of double stranded DNA.

Figure 11 schematically illustrates a method for determining the position of an aromatic amino acid residue that should be replaced with a β -alanine residue in order to enhance the DNA binding properties of certain twelve carboxamide residue hairpin polyamide compounds.

DETAILED DESCRIPTION OF THE INVENTION

Within this application, unless otherwise stated, definitions of the terms and illustration of the techniques of this application may be found in any of several well-known references such as: Sambrook, J., *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989); Goeddel, D., *ed.*, *Gene Expression Technology, Methods in Enzymology*, **185**, Academic Press, San Diego, CA (1991); "Guide to Protein Purification" in Deutscher, M.P., *ed.*, *Methods in Enzymology*, Academic Press, San Diego, CA (1989); Innis *et al.*, *PCR Protocols: A Guide to Methods and Applications*, Academic Press, San Diego, CA (1990); Freshney, R.I., *Culture of Animal Cells: A Manual of Basic Technique*, 2nd Ed., Alan Liss, Inc. New York, NY (1987); Murray, E.J., *ed.*, *Gene Transfer and Expression Protocols*, pp.
30 109-128, The Humana Press Inc., Clifton, NJ and Lewin, B., *Genes VI*, Oxford University Press, New York (1997).
35

For the purposes of this application, a *promoter* is a regulatory sequence of DNA that is involved in the binding of RNA polymerase to initiate transcription of a gene. A *gene* is a segment of DNA involved in producing a peptide, polypeptide or protein, including the coding region, non-coding regions preceding (“leader”) and following (“trailer”) the coding region, as well as intervening non-coding sequences (“introns”) between individual coding segments (“exons”). Coding refers to the representation of amino acids, start and stop signals in a three base “triplet” code. Promoters are often upstream (“5’ to”) the transcription initiation site of the corresponding gene. Other regulatory sequences of DNA in addition to promoters are known, including sequences involved with the binding of transcription factors, including response elements that are the DNA sequences bound by inducible factors. Enhancers comprise yet another group of regulatory sequences of DNA that can increase the utilization of promoters, and can function in either orientation (5’-3’ or 3’-5’) and in any location (upstream or downstream) relative to the promoter. Preferably, the regulatory sequence has a positive activity, i.e., binding of an endogeneous ligand (e.g. a transcription factor) to the regulatory sequence increases transcription, thereby resulting in increased expression of the corresponding target gene. In such a case, interference with transcription by binding a polyamide to a regulatory sequence would reduce or abolish expression of a gene.

The promoter may also include or be adjacent to a regulatory sequence known in the art as a *silencer*. A silencer sequence generally has a negative regulatory effect on expression of the gene. In such a case, expression of a gene may be increased directly by using a polyamide to prevent binding of a factor to a silencer regulatory sequence or indirectly, by using a polyamide to block transcription of a factor to a silencer regulatory sequence.

It is to be understood that the polyamides of this invention bind to double stranded DNA in a sequence specific manner. The function of a segment of DNA of a given sequence, such as 5’-TATAAA-3’, depends on its position relative to other functional regions in the DNA sequence. In this case, if the sequence 5’-TATAAA-3’ on the coding strand of DNA is positioned about 30 base pairs upstream of the transcription start site, the sequence forms part of the promoter region (Lewin, *Genes VI*, pp. 831-835). On the other hand, if the sequence 5’-TATAAA-3’ is downstream of the transcription start site in a coding region and in proper

register with the reading frame, the sequence encodes the tyrosyl and lysyl amino acid residues (Lewin, *Genes VI*, pp. 213-215).

5 While not being held to one hypothesis, it is believed that the binding of the polyamides of this invention modulate gene expression by altering the binding of DNA binding proteins, such as RNA polymerase, transcription factors, TBF, TFIIB and other proteins. The effect on gene expression of polyamide binding to a segment of double stranded DNA is believed to be related to the function, e.g., promoter, of that segment of DNA.

10 It is to be understood by one skilled in the art that the improved polyamides of the present invention may bind to any of the above-described DNA sequences or any other sequence having a desired effect upon expression of a gene. In addition, U.S. Patent No. 5,578,444 describes numerous promoter targeting sequences from which base pair sequences for targeting an improved polyamide of the present invention may be identified.

It is generally understood by those skilled in the art that the basic structure of DNA in a living cell includes both *major* and a *minor groove*. For the purposes of describing the present invention, the *minor groove* is the narrow groove of DNA as illustrated in common molecular biology references such as Lewin, B., *Genes VI*, Oxford University Press, New York (1997).

25 To affect gene expression in a cell, which may include causing an increase or a decrease in gene expression, a effective quantity of one or more polyamide is contacted with the cell and internalized by the cell. The cell may be contacted *in vivo* or *in vitro*. Effective extracellular concentrations of polyamides that can modulate gene expression range from about 10 nanomolar to about 1 micromolar. Gottesfeld, J.M., *et al.*, *Nature* **387** 202-205 (1997). To determine effective amounts and concentrations of polyamides *in vitro*, a suitable number of cells is plated on tissue culture plates and various quantities of one or more polyamide are added to separate
30 wells. Gene expression following exposure to a polyamide can be monitored in the cells or medium by detecting the amount of the protein gene product present as determined by various techniques utilizing specific antibodies, including ELISA and western blot. Alternatively, gene expression following exposure to a polyamide can be monitored by detecting the amount of messenger RNA present as determined by various techniques, including northern blot and RT-PCR.

Similarly, to determine effective amounts and concentrations of polyamides for *in vivo* administration, a sample of body tissue or fluid, such as plasma, blood, urine, cerebrospinal fluid, saliva, or biopsy of skin, muscle, liver, brain or other appropriate tissue source is analyzed.

5 Gene expression following exposure to a polyamide can be monitored by detecting the amount of the protein gene product present as determined by various techniques utilizing specific antibodies, including ELISA and western blot. Alternatively, gene expression following exposure to a polyamide can be monitored by the detecting the amount of messenger RNA present as determined by various techniques, including northern blot and RT-PCR.

10

The polyamides of this invention may be formulated into diagnostic and therapeutic compositions for *in vivo* or *in vitro* use. Representative methods of formulation may be found in *Remington: The Science and Practice of Pharmacy*, 19th ed., Mack Publishing Co., Easton, PA (1995).

For *in vivo* use, the polyamides may be incorporated into a physiologically acceptable pharmaceutical composition that is administered to a patient in need of treatment or an animal for medical or research purposes. The polyamide composition comprises pharmaceutically acceptable carriers, excipients, adjuvants, stabilizers, and vehicles. The composition may be in solid, liquid, gel, or aerosol form. The polyamide composition of the present invention may be administered in various dosage forms orally, parentally, by inhalation spray, rectally, or topically. The term parenteral as used herein includes, subcutaneous, intravenous, intramuscular, intrasternal, infusion techniques or intraperitoneally.

25

The selection of the precise concentration, composition, and delivery regimen is influenced by, *inter alia*, the specific pharmacological properties of the particular selected compound, the intended use, the nature and severity of the condition being treated or diagnosed, the age, weight, gender, physical condition and mental acuity of the intended recipient as well as the route of administration. Such considerations are within the purview of the skilled artisan.

30 Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods.

Polyamides of the present invention are also useful for detecting the presence of double stranded DNA of a specific sequence for diagnostic or preparative purposes. The sample containing the double stranded DNA can be contacted by polyamide linked to a solid substrate, thereby isolating DNA comprising a desired sequence. Alternatively, polyamides linked to a suitable detectable marker, such as biotin, a hapten, a radioisotope or a dye molecule, can be contacted by a sample containing double stranded DNA.

The design of bifunctional sequence specific DNA binding molecules requires the integration of two separate entities: recognition and functional activity. Polyamides that specifically bind with subnanomolar affinity to the minor groove of a predetermined sequence of double stranded DNA are linked to a functional molecule, providing the corresponding bifunctional conjugates useful in molecular biology, genomic sequencing, and human medicine. Polyamides of this invention can be conjugated to a variety of functional molecules, which can be independently chosen from but is not limited to arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, oligodeoxynucleotides, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- α -tocopheral, psoralen, EDTA, methidium, acridine, Ni(II)•Gly-Gly-His, TO, Dansyl, pyrene, N-bromoacetamide, and gold particles. Such bifunctional polyamides are useful for DNA affinity capture, covalent DNA modification, oxidative DNA cleavage, and DNA photocleavage. Such bifunctional polyamides are useful for DNA detection by providing a polyamide linked to a detectable label. Detailed instructions for synthesis of such bifunctional polyamides can be found in copending U.S. provisional application 60/043,444, the teachings of which are incorporated by reference.

DNA complexed to a labeled polyamide can then be determined using the appropriate detection system as is well known to one skilled in the art. For example, DNA associated with a polyamide linked to biotin can be detected by a streptavidin / alkaline phosphatase system.

The present invention also describes a diagnostic system, preferably in kit form, for assaying for the presence of the double stranded DNA sequence bound by the polyamide of this invention in a body sample, such brain tissue, cell suspensions or tissue sections, or body fluid

samples such as CSF, blood, plasma or serum, where it is desirable to detect the presence, and preferably the amount, of the double stranded DNA sequence bound by the polyamide in the sample according to the diagnostic methods described herein.

5 The diagnostic system includes, in an amount sufficient to perform at least one assay, a specific polyamide as a separately packaged reagent. Instructions for use of the packaged reagent(s) are also typically included. As used herein, the term "package" refers to a solid matrix or material such as glass, plastic (e.g., polyethylene, polypropylene or polycarbonate), paper, foil and the like capable of holding within fixed limits a polyamide of the present invention. Thus, for example, a package can be a glass vial used to contain milligram quantities of a contemplated polyamide or it can be a microliter plate well to which microgram quantities of a contemplated polyamide have been operatively affixed, i.e., linked so as to be capable of being bound by the target DNA sequence. "Instructions for use" typically include a tangible expression describing the reagent concentration or at least one assay method parameter such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent or sample admixtures, temperature, buffer conditions and the like. A diagnostic system of the present invention preferably also includes a detectable label and a detecting or indicating means capable of signaling the binding of the contemplated polyamide of the present invention to the target DNA sequence. As noted above, numerous detectable labels, such as biotin, and detecting or indicating means, such as enzyme-linked (direct or indirect) streptavidin, are well known in the art.

As used herein, "subnanomolar affinity" means binding that is characterized by a dissociation constant, K_d , of less than 1 nM, as measured by DNase I footprint titration.

25 Preferably, polyamides of the present invention are characterized by subnanomolar binding affinity for the identified target DNA sequence. As used herein, the "selectivity" of the binding of a polyamide to a DNA sequence is the ratio of the dissociation constant, K_d , as measured by DNase I footprint titration of binding the polyamide to a mismatch DNA sequence divided by the corresponding dissociation constant of the binding of the polyamide to the identified target DNA sequence. Preferably, polyamides of the present invention are characterized by a selectivity of 5 or greater, more preferably a selectivity of greater than 10.

The exemplary polyamide that illustrates the compositions and methods of the present invention is polyamide **3** of Figure 1, ImImHpPy- γ -ImPyPyPy- β -Dp. This polyamide was designed according to the method of the present invention to target the identified sequence 5'-WGGTCW-3'. See Table 5, below, Sequence No. 36 and the corresponding sequence of
 5 carboxamide binding pairs. Polyamide **3** binds an identified target sequence 5'-TGGTCA-3' with a dissociation constant, as measured by DNase I footprint titration, of 0.48 nM, i.e., with subnanomolar affinity as defined herein (see Table 1, below). The polyamide binds to the mismatch sequence 5'-TGGACA-3' with a dissociation constant of 37 nM, yielding a selectivity, as defined herein, of 77 (Table 1).

Figure 1 shows representative structures of polyamides. ImImPyPy- γ -ImPyPyPy- β -Dp (1), ImImPyPy- γ -ImHpPyPy- β -Dp (2), and ImImHpPy- γ -ImPyPyPy- β -Dp (3). (Hp = 3-hydroxy-N-methylpyrrole, Im = N-methylimidazole, Py = N-methylpyrrole, β = β -alanine, γ = γ -aminobutyric acid, Dp = Dimethylaminopropylamide). Polyamides were synthesized by solid phase methods using Boc-protected 3-methoxypyrrole, imidazole, and pyrrole aromatic amino acids, cleaved from the support by aminolysis, deprotected with sodium thiophenoxide, and purified by reversed phase HPLC. Baird, E. E. & Dervan, P. B. describes the solid phase synthesis of polyamides containing imidazole and pyrrole amino acids. *J. Am. Chem. Soc.* **118**, 6141-6146 (1996); also see PCT US 97/003332. The identity and purity of the polyamides were verified by ¹H NMR, analytical HPLC, and matrix-assisted laser-desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS-monoisotopic): **1** 1223.6 (1223.6 calculated), **2** 1239.6 (1239.6 calculated); **3** 1239.6 (1239.6 calculated).

Figure 2 illustrates binding models for polyamides **1-3** in complex with 5'-TGGTCA-3' and 5'-TGGACA-3' (A•T and T•A in fourth position highlighted). Filled and unfilled circles represent imidazole and pyrrole rings respectively; circles containing an H represent 3-hydroxypyrrole, the curved line connecting the polyamide subunits represents γ -aminobutyric acid, the diamond represents β -alanine, and the + represents the positively charged dimethylaminopropylamide tail group.

Figure 3 shows quantitative DNase I footprint titration experiments with polyamides **2** and **3** on the 3' ³²P labeled 250-bp pJK6 *Eco*RI/*Pvu*II restriction fragment. Lane 1, intact DNA; lanes 2-11 DNase I digestion products in the presence of 100, 50, 20, 10, 5, 2, 1, 0.5, 0.2, 0.1 nM







polyamide, respectively; lane 12, DNase I digestion products in the absence of polyamide; lane 13, adenine-specific chemical sequencing. Iverson, B. L. & Dervan, P. B. describes an adenine-specific DNA chemical sequencing reaction. *Methods Enzymol.* **15**, 7823-7830 (1987). All reactions were done in a total volume of 400 μ L. A polyamide stock solution or H₂O was added to an assay buffer containing radiolabeled restriction fragment, with the final solution conditions of 10 mM Tris-HCl, 10 mM KCl, 10 mM MgCl₂, 5 mM CaCl₂, pH 7.0. Solutions were allowed to equilibrate for 4-12 h at 22 °C before initiation of footprinting reactions. Footprinting reactions, separation of cleavage products, and data analysis were carried out as described. White, S., Baird, E. E. & Dervan, P. B. Effects of the A•T/T•A degeneracy of pyrrole-imidazole polyamide recognition in the minor groove of DNA. *Biochemistry* **35**, 12532-12537 (1996).

Figure 4 shows the structure and equilibrium dissociation constant for numerous compounds of the present invention. Polyamides are shown in complex with their respective match site. Filled and unfilled circles represent imidazole (Im) and pyrrole (Py) rings, respectively; circles containing an H represent 3-hydroxypyrrole (Hp), the curved line connecting the polyamide subunits represents γ -aminobutyric acid (γ), the diamond represents β -alanine (β), and the + represents the positively charged dimethylaminopropylamide tail group (Dp). The equilibrium dissociation constants are the average values obtained from three DNase I footprint titration experiments. The standard deviation for each set is less than 15% of the reported number. Assays were carried out in the presence of 10 mM Tris•HCl, 10 mM KCl, 10 mM MgCl₂, and 5 mM CaCl₂ at pH 7.0 and 22°C.

Four-ring polyamide subunits, covalently coupled to form eight-ring hairpin structures, bind specifically to 6-bp target sequences at subnanomolar concentrations. Trauger, J.W., Baird, E. E. & Dervan, P.B. describe the recognition of DNA by designed ligands at subnanomolar concentrations. *Nature* **382**, 559-561 (1996); Swalley, S. E., Baird, E. E. & Dervan, P. B. describe the discrimination of 5'-GGGG-3', 5'-GCGC-3', and 5'-GGCC-3' sequences in the minor groove of DNA by eight-ring hairpin polyamides. *J. Am. Chem. Soc.* **119**, 6953-6961 (1997). The DNA-binding affinities of three eight-ring hairpin polyamides shown in Figure 1 as compound **1**, **2**, and **3** containing pairings of Im/Py, Py/Im opposite G•C, C•G and either Py/Py, Hp/Py, or Py/Hp at a common single point opposite T•A and A•T has been determined. Equilibrium dissociation constants (K_d) for ImImPyPy- γ -ImPyPyPy- β -Dp **1**, ImImPyPy- γ -ImHpPyPy- β -Dp **2**, ImImHpPy- γ -ImPyPyPy- β -Dp **3** of Figure 1 are shown in Table 1. Brenowitz, M., Senear, D. F., Shea, M. A. & Ackers, G. K. describe a quantitative DNase footprint titration method for studying protein-DNA interactions. *Methods Enzymol.* **130**, 132-

181 (1986); The K_d values were determined by quantitative DNase I footprint titration experiments: on a 3' 32 P-labeled 250-bp DNA fragment containing the target sites, 5'-TGGACA-3' and 5'-TGGTCA-3' which differ by a single A•T base pair in the fourth position. The DNase footprint gels are shown in Figure 3.

5

TABLE 1 Equilibrium dissociation constants*			
Polyamide†	5'-TGGTCA-3'	5'-TGGACA-3'	K_{rel}^\ddagger
1 Py/Py	 $K_d = 0.077$ nM	 $K_d = 0.15$ nM	2.0
2 Py/Hp	 $K_d = 15$ nM	 $K_d = 0.83$ nM	0.06
3 Hp/Py	 $K_d = 0.48$ nM	 $K_d = 37$ nM	77

*The reported dissociation constants are the average values obtained from three DNase I footprint titration experiments. The standard deviation for each data set is less than 15% of the reported number. Assays were carried out in the presence of 10 mM Tris•HCl, 10 mM KCl, 10 mM MgCl₂, and 5 mM CaCl₂ at pH 7.0 and 22 °C.

†Ring pairing opposite T•A and A•T in the fourth position.

‡Calculated as $K_d(5'-TGGACA-3')/K_d(5'-TGGTCA-3')$.

Based on the pairing rules for polyamide-DNA complexes both of these sequences are a match for control polyamide 1 which places a Py/Py pairing opposite A•T and T•A at both sites. It was determined that polyamide 1 (Py/Py) binds to 5'-TGGTCA-3' and 5'-TGGACA-3' within a factor of 2 ($K_d = 0.077$ or 0.15 nM respectively). In contrast, polyamide 2 (Py/Hp) binds to 5'-TGGTCA-3' and 5'-TGGACA-3' with dissociation constants which differ by a factor of 18 ($K_d = 15$ nM and 0.83 nM respectively). By reversing the pairing in polyamide 3 (Hp/Py) the dissociation constants differ again in the opposite direction by a factor of 77 ($K_d = 0.48$ nM and 37 nM respectively). Control experiments performed on separate DNA fragments; reveal that neither a 5'-TGGCA-3' or a 5'-TGGCA-3' site is bound by polyamide 2 or 3 at concentrations ≤ 100 nM, indicating that the Hp/Py and Py/Hp ring pairings do not bind opposite G•C or C•G.

15

20

The specificity of polyamides 2 and 3 for sites which differ by a single A•T/T•A base pair results from small chemical changes. Replacing the Py/Py pair in 1 with a Py/Hp pairing as in 2, a single substitution of C3-OH for C3-H, destabilizes interaction with 5'-TGGTCA-3' by 191-fold, a free energy difference of 3.1 kcal mol⁻¹. Interaction of 2 with 5'-TGGACA-3' is destabilized only 6-fold relative to 1, a free energy difference of 1.1 kcal mol⁻¹. Similarly,

replacing the Py/Py pair in **1** with Hp/Py as in **3** destabilizes interaction with 5'-TGGACA-3' by 252-fold, a free energy difference of 3.2 kcal mol⁻¹. Interaction of **3** with 5'TGGICA-3' is destabilized only 6-fold relative to **1**, a free energy difference of 1.0 kcal mol⁻¹.

5 The polyamides of this invention provide for coded targeting of predetermined DNA sequences with affinity and specificity comparable to sequence-specific DNA binding proteins. Hp, Im, and Py polyamides complete the minor groove recognition code using three aromatic amino acids which combine to form four ring pairings (Im/Py, Py/Im, Hp/Py, and Py/Hp) which complement the four Watson-Crick base pairs, as shown in TABLE 2. There are a possible 240
10 four base pair sequences which contain at least 1 A•T or T•A base pair and therefore can advantageously use an Hp/Py, or Py/Hp carboxamide binding. Polyamides binding to any of these sequences can be designed in accordance with the code of TABLE 2.

TABLE 2 Pairing code for minor groove recognition*

Pair	G•C	C•G	T•A	A•T
Im/Py	+	-	-	-
Py/Im	-	+	-	-
Hp/Py	-	-	+	-
Py/Hp	-	-	-	+

* favored (+), disfavored (-)

For certain G•C rich sequences the affinity of polyamide•DNA complexes may be enhanced by substitution of an Im/β pair for Im/Py at G•C and β/Im for Py/Im at C•G. At A•T and T•A base pairs, either a Py/β, β/Py, Hp/β, β/Hp, and β/β may be used. The alternate aliphatic/aromatic amino acid pairing code is described in Table 3.

TABLE 3 Aliphatic/ Aromatic substitution for ring pairings*

Pair	Substitution
Im/Py	Im/β
Py/Im	β/Im
Hp/Py	Py/β, β/Py, Hp/β, β/β
Py/Hp	Py/β, β/Py, β/Hp, β/β

U. S. Patent 5,578,444 describes numerous promoter region targeting sequences from which base pair sequences for targeting a polyamide can be identified.

PCT U.S. 97/003332 describes methods for synthesis of polyamides which are suitable for preparing polyamides of this invention. The use of β -alanine in place of a pyrrole amino acid in the synthetic methods provides aromatic/aliphatic pairing (Im/ β , β /Im, Hp/ β , β /Hp, Py/ β , and β /Py) and aliphatic/aliphatic pairing (β / β) substitution. The use of γ -aminobutyric acid, or a substituted γ -aminobutyric acid such as (R)-2,4 diaminobutyric acid, provides for preferred hairpin turns. The following examples illustrate the synthesis of polyamides of the present invention.

The process of designing a preferred polyamide molecule $X_1X_2X_3X_4\text{-}\gamma\text{-}X_5X_6X_7X_8$ comprising eight aromatic amino acid residues of this invention is shown schematically in Figure 5. The polyamide design process provides a method for designing an eight carboxamide residue molecule comprising four carboxamide binding pairs for detection and binding of a target six base pair 5'-WNNNNW-3' sequence in the minor groove of double stranded DNA. The design process identifies an appropriate polyamide ligand for recognition of a predetermined 6-bp, 5'-WNNNNW-3' sequence with subnanomolar affinity and >10-fold specificity versus mismatch sites. Trauger, J.W., Baird, E. E. Dervan, P.B. describes the recognition of DNA by designed ligands at subnanomolar concentrations. *Nature* **382**, 559-561 (1996).

In order to prepare a polyamide molecule specific for an identified six base pair sequence of double stranded DNA, a user starts the 8-ring polyamide design process that implements the minor groove recognition pairing code summarized in Table 2 above. In the design process a 5'-WNNNNW-3' sequence was identified. In a preferred embodiment, the identified sequence was located within a gene promoter. U. S. Patent 5,578,444 describes numerous promoter region targeting sequences from which target six base pair sequences for targeting a polyamide can be identified. The identified sequence was then defined as 5'-W**abcd**W-3' in a stepwise process wherein **a**, **b**, **c**, and **d**, were sequentially and independently defined as A, G, C, or T. The structure of the polyamide molecule was then correspondingly defined by sequentially choosing antiparallel carboxamide binding pairs according to the minor groove pairing code summarized in Table 2 above. Thus, if **a** was G, then X_1 was defined as Im, and X_8 was defined as Py. If **a** was C, then X_1 was defined as Py, and X_8 was defined as Im. If **a** was T, then X_1 was defined

as Hp, and X8 was defined as Py. If **a** was A, then X1 was defined as Py, and X8 was defined as Hp.

Similarly, **b** was defined as A, G, C, or T and corresponding carboxamide binding pairs were defined. According to the same rules, if **b** was G, then X2 was defined as Im, and X7 was defined as Py. If **b** was C, then X2 was defined as Py, and X7 was defined as Im. Likewise, if **b** was T, then X2 was defined as Hp, and X7 was defined as Py. If **b** was A, then X2 was defined as Py, and X7 was defined as Hp.

The next step was to define **c** as A, G, C, or T and then define corresponding carboxamide binding pairs. Following the same rules, if **c** was G, then X3 was defined as Im, and X6 was defined as Py. If **c** was C, then X3 was defined as Py, and X6 was defined as Im. Similarly, if **c** was T, then X3 was defined as Hp, and X6 was defined as Py. If **c** was A, then X3 was defined as Py, and X6 was defined as Hp. Lastly, **d** was defined as A, G, C, or T and the last corresponding carboxamide binding pair was defined. According to above rules, if **d** was G, then X4 was defined as Im, and X5 was defined as Py. If **d** was C, then X4 was defined as Py, and X5 was defined as Im. If **d** was T, then X4 was defined as Hp, and X5 was defined as Py. If **d** was A, then X4 was defined as Py, and X5 was defined as Hp.

With all eight carboxamide residues that participate in binding pairs now defined, the designed polyamide X₁X₂X₃X₄- γ -X₅X₆X₇X₈ suitable for binding to the identified sequence was synthesized using known techniques. Baird, E. E. & Dervan, P. B. describes the solid phase synthesis of polyamides containing imidazole and pyrrole amino acids. *J. Am. Chem. Soc.* **118**, 6141-6146 (1996); *also see* PCT US 97/003332.

The binding affinity of the synthesized polyamide to the identified sequence was determined using a quantitative DNase footprint titration method for studying protein-DNA interactions described by Brenowitz, M., Senear, D. F., Shea, M. A. & Ackers, G. K., *Methods Enzymol.* **130**, 132-181 (1986). If the affinity of the synthesized polyamide at the target site was not subnanomolar affinity then adding a β -alanine (process A) was considered in order to optimize the exact positions of the binding pairs of aromatic amino acids. If the affinity of the said polyamide at said target site was subnanomolar affinity then the sequence specificity of the polyamide versus mismatch sequences was determined. If the specificity versus mismatch sites

was not > 10-fold specificity then adding a β -alanine (process A schematically shown in Figure 6) was considered, in order to optimize the positions of the aromatic amino acids in relationship to the base pairs in the minor groove. Specificity of the polyamide molecule for the target identified sequence versus mismatch sequence sites of greater than 10-fold was considered a successful result of design process.

The 256 polyamide molecules comprising four carboxamide binding pairs that were designed using this method are useful for binding to the 256 target 5'-NNNN-3' core sequences, and are listed in Tables 4-11. A corresponding polyamide molecule was designed for each DNA sequence (1-240) and (G1-G16) using the process outlined above and shown schematically in Figure 5.

If the synthesized polyamide molecule did not bind to the target identified sequence with subnanomolar affinity or if the synthesized polyamide molecule did not exhibit a specificity for the target identified sequence versus mismatch sequence sites of greater than 10-fold, the option of substituting an aliphatic amino acid residues for one of the carboxamide residues was considered. The preferred aliphatic amino acid residue is β -alanine. At least one aliphatic amino acid residue such as a β -alanine residue provided some flexibility to the central portion of the polyamide molecule, acting as a "spring" to permit optimization of the hydrogen bonding between the carboxamide binding pairs and the nucleotide bases of the double stranded DNA.

In general, it was not found to be advantageous to replace either member of the terminal carboxamide binding pair, X1/X8, with β -alanine. Similarly, β -alanine was not substituted for members of the binding pair, X4/X5, adjacent to the γ hairpin residue. β -alanine residues were not substituted for N-methylimidazole residues. The use of β -alanine in place of a pyrrole or 3-hydroxypyrrole amino acid residue provides aromatic/aliphatic pairing (Im/ β , β /Im, Hp/ β , β /Hp, Py/ β , and β /Py) and aliphatic/aliphatic pairing (β / β) substitution.

The method for selecting which pyrrole amino acid to substitute with β -alanine is schematically illustrated in Figure 6. Selective placement of an aliphatic β -alanine (β) residue paired with either a pyrrole (Py), 3-hydroxypyrrole (Hp), or imidazole (Im) aromatic amino acid or another β -alanine residue is found to compensate for sequence composition effects to improve

recognition and binding of the minor groove of DNA by pyrrole-imidazole polyamides of the present invention. If an all-ring polyamide has been found to have an affinity which is not subnanomolar, or a specificity versus mismatch sequences which is less than 10-fold it may be caused by DNA sequence-composition effects which can be reduced by replacement of an aromatic amino acid with an aliphatic β -alanine residue. In a polyamide molecule that comprises four binding pairs it is only beneficial to place β -alanine in positions X₂, X₃, X₆, and X₇. No more than two β -alanine residues may be placed within a single hairpin structure. No more than a single β -residue may be placed within each individual polyamide subunit, e.g., if X₂ is replaced with β -alanine, then X₃ cannot be replaced.

These rules and others were implemented in the method schematically illustrated in Figure 6. This process is suitable for the refinement of the design polyamide comprising four binding pairs that has been designed by the method illustrated in Figure 5, but which lacks subnanomolar affinity or greater than 10-fold specificity at the identified target DNA sequence. As in the basic design method, the designed polyamides are synthesized and the affinity and specificity of binding to the target DNA were determined.

For a given polyamide molecule X₁X₂X₃X₄- γ -X₅X₆X₇X₈ there are five possible outcomes for the process of substituting a β -alanine residue for an aromatic amino acid residue. First, there may be no position at which it is possible to add a β -alanine residue; in such case, a better binding affinity or selectivity can be sought in the design and synthesis of a polyamide with five or six carboxamide binding pairs, described below. Second, the process may result in a derivative which contains a single β -alanine substitution (such derivatives are numbered according to the parent numbering scheme such that a single β -derivative of compound 5 would be called 5 β), which is sufficient to produce subnanomolar binding affinity and >10-fold specificity, and at which point the process is deemed complete.

Third, the process of Figure 5 may result in a polyamide which contains a single β -alanine substitution which is not sufficient to produce subnanomolar binding affinity and >10-fold specificity, but where there are no additional positions in which it is possible to substitute a β -alanine residue, and in such a case a polyamide with five or six carboxamide binding pairs, should be designed and synthesized, as described below. Fourth, the process of Figure 5 may

result in a polyamide that contains a single β -alanine substitution that is not sufficient to produce subnanomolar binding affinity and >10-fold specificity, but where there is an additional position for β -alanine substitution that does produce a polyamide with the criterion level of affinity and selectivity and therefore the design process is deemed complete. Polyamides that were designed by the process that produces polyamide molecules that contain two β -alanine residues are labeled β 2 in Tables 12-19.

A fifth possibility is that substitution at a second position by the method illustrated in Figure 6 with a second β -alanine residue is not sufficient to produce a polyamide having the subnanomolar binding affinity and >10-fold specificity, and a polyamide with five or six carboxamide binding pairs, should be designed and synthesized, as described below. Tables 12-19 list polyamides corresponding to sequences 1-240 and G1-G16 which contain either one or two β -alanine residues.

TABLE 4: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1) 5'-W G T T T W-3'	ImHpHpHp- γ -PyPyPyPy
5	2) 5'-W G T T A W-3'	ImHpHpPy- γ -HpPyPyPy
	3) 5'-W G T T G W-3'	ImHpHpIm- γ -PyPyPyPy
	4) 5'-W G T T C W-3'	ImHpHpPy- γ -ImPyPyPy
	5) 5'-W G T A T W-3'	ImHpPyHp- γ -PyHpPyPy
	6) 5'-W G T A A W-3'	ImHpPyPy- γ -HpHpPyPy
10	7) 5'-W G T A G W-3'	ImHpPyIm- γ -PyHpPyPy
	8) 5'-W G T A C W-3'	ImHpPyPy- γ -ImHpPyPy
	9) 5'-W G T G T W-3'	ImHpImHp- γ -PyPyPyPy
	10) 5'-W G T G A W-3'	ImHpImPy- γ -HpPyPyPy
	11) 5'-W G T G G W-3'	ImHpImIm- γ -PyPyPyPy
	12) 5'-W G T G C W-3'	ImHpImPy- γ -ImPyPyPy
	13) 5'-W G T C T W-3'	ImHpPyHp- γ -PyImPyPy
	14) 5'-W G T C A W-3'	ImHpPyPy- γ -HpImPyPy
	15) 5'-W G T C G W-3'	ImHpPyIm- γ -PyImPyPy
	16) 5'-W G T C C W-3'	ImHpPyPy- γ -ImImPyPy
20	17) 5'-W G A T T W-3'	ImPyHpHp- γ -PyPyHpPy
	18) 5'-W G A T A W-3'	ImPyHpPy- γ -HpPyHpPy
	19) 5'-W G A T G W-3'	ImPyHpIm- γ -PyPyHpPy
	20) 5'-W G A T C W-3'	ImPyHpPy- γ -ImPyHpPy
	21) 5'-W G A A T W-3'	ImPyPyHp- γ -PyHpHpPy
25	22) 5'-W G A A A W-3'	ImPyPyPy- γ -HpHpHpPy
	23) 5'-W G A A G W-3'	ImPyPyIm- γ -PyHpHpPy
	24) 5'-W G A A C W-3'	ImPyPyPy- γ -ImHpHpPy
	25) 5'-W G A G T W-3'	ImPyImHp- γ -PyPyHpPy
	26) 5'-W G A G A W-3'	ImPyImPy- γ -HpPyHpPy
30	27) 5'-W G A G G W-3'	ImPyImIm- γ -PyPyHpPy
	28) 5'-W G A G C W-3'	ImPyImPy- γ -ImPyHpPy
	29) 5'-W G A C T W-3'	ImPyPyHp- γ -PyImHpPy
	30) 5'-W G A C A W-3'	ImPyPyPy- γ -HpImHpPy
	31) 5'-W G A C G W-3'	ImPyPyIm- γ -PyImHpPy
35	32) 5'-W G A C C W-3'	ImPyPyPy- γ -ImImHpPy

TABLE 5: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	33) 5'-W G G T T W-3'	ImImHpHp- γ -PyPyPyPy
5	34) 5'-W G G T A W-3'	ImImHpPy- γ -HpPyPyPy
	35) 5'-W G G T G W-3'	ImImHpIm- γ -PyPyPyPy
	36) 5'-W G G T C W-3'	ImImHpPy- γ -ImPyPyPy
	37) 5'-W G G A T W-3'	ImImPyHp- γ -PyHpPyPy
	38) 5'-W G G A A W-3'	ImImPyPy- γ -HpHpPyPy
10	39) 5'-W G G A G W-3'	ImImPyIm- γ -PyHpPyPy
	40) 5'-W G G A C W-3'	ImImPyPy- γ -ImHpPyPy
	41) 5'-W G G G T W-3'	ImImImHp- γ -PyPyPyPy
	42) 5'-W G G G A W-3'	ImImImPy- γ -HpPyPyPy
	43) 5'-W G G C T W-3'	ImImPyHp- γ -PyImPyPy
	44) 5'-W G G C A W-3'	ImImPyPy- γ -HpImPyPy
	45) 5'-W G C T T W-3'	ImPyHpHp- γ -PyPyImPy
	46) 5'-W G C T A W-3'	ImPyHpPy- γ -HpPyImPy
	47) 5'-W G C T G W-3'	ImPyHpIm- γ -PyPyImPy
	48) 5'-W G C T C W-3'	ImPyHpPy- γ -ImPyImPy
20	49) 5'-W G C A T W-3'	ImPyPyHp- γ -PyHpImPy
	50) 5'-W G C A A W-3'	ImPyPyPy- γ -HpHpImPy
	51) 5'-W G C A G W-3'	ImPyPyIm- γ -PyHpImPy
	52) 5'-W G C A C W-3'	ImPyPyPy- γ -ImHpImPy
	53) 5'-W G C G T W-3'	ImPyImHp- γ -PyPyImPy
25	54) 5'-W G C G A W-3'	ImPyImPy- γ -HpPyImPy
	55) 5'-W G C C T W-3'	ImPyPyHp- γ -PyImImPy
	56) 5'-W G C C A W-3'	ImPyPyPy- γ -HpImImPy
	G1) 5'-W G G G G W-3'	ImImImIm- γ -PyPyPyPy
	G2) 5'-W G G G C W-3'	ImImImPy- γ -ImPyPyPy
30	G3) 5'-W G G C G W-3'	ImImPyIm- γ -PyImPyPy
	G4) 5'-W G G C C W-3'	ImImPyPy- γ -ImImPyPy
	G5) 5'-W G C G G W-3'	ImPyImIm- γ -PyPyImPy
	G6) 5'-W G C G C W-3'	ImPyImPy- γ -ImPyImPy
	G7) 5'-W G C C G W-3'	ImPyPyIm- γ -PyImImPy
35	G8) 5'-W G C C C W-3'	ImPyPyPy- γ -ImImImPy

TABLE 6: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	57) 5'-W T T T T W-3'	HpHpHpHp- γ -PyPyPyPy
5	58) 5'-W T T T A W-3'	HpHpHpPy- γ -HpPyPyPy
	59) 5'-W T T T G W-3'	HpHpHpIm- γ -PyPyPyPy
	60) 5'-W T T T C W-3'	HpHpHpPy- γ -ImPyPyPy
	61) 5'-W T T A T W-3'	HpHpPyHp- γ -PyHpPyPy
	62) 5'-W T T A A W-3'	HpHpPyPy- γ -HpHpPyPy
10	63) 5'-W T T A G W-3'	HpHpPyIm- γ -PyHpPyPy
	64) 5'-W T T A C W-3'	HpHpPyPy- γ -ImHpPyPy
	65) 5'-W T T G T W-3'	HpHpImHp- γ -PyPyPyPy
	66) 5'-W T T G A W-3'	HpHpImPy- γ -HpPyPyPy
	67) 5'-W T T G G W-3'	HpHpImIm- γ -PyPyPyPy
	68) 5'-W T T G C W-3'	HpHpImPy- γ -ImPyPyPy
	69) 5'-W T T C T W-3'	HpHpPyHp- γ -PyImPyPy
	70) 5'-W T T C A W-3'	HpHpPyPy- γ -HpImPyPy
	71) 5'-W T T C G W-3'	HpHpPyIm- γ -PyImPyPy
	72) 5'-W T T C C W-3'	HpHpPyPy- γ -ImImPyPy
20	73) 5'-W T A T T W-3'	HpPyHpHp- γ -PyPyHpPy
	74) 5'-W T A T A W-3'	HpPyHpPy- γ -HpPyHpPy
	75) 5'-W T A T G W-3'	HpPyHpIm- γ -PyPyHpPy
	76) 5'-W T A T C W-3'	HpPyHpPy- γ -ImPyHpPy
	77) 5'-W T A A T W-3'	HpPyPyHp- γ -PyHpHpPy
25	78) 5'-W T A A A W-3'	HpPyPyPy- γ -HpHpHpPy
	79) 5'-W T A A G W-3'	HpPyPyIm- γ -PyHpHpPy
	80) 5'-W T A A C W-3'	HpPyPyPy- γ -ImHpHpPy
	81) 5'-W T A G T W-3'	HpPyImHp- γ -PyPyHpPy
	82) 5'-W T A G A W-3'	HpPyImPy- γ -HpPyHpPy
30	83) 5'-W T A G G W-3'	HpPyImIm- γ -PyPyHpPy
	84) 5'-W T A G C W-3'	HpPyImPy- γ -ImPyHpPy
	85) 5'-W T A C T W-3'	HpPyPyHp- γ -PyImHpPy
	86) 5'-W T A C A W-3'	HpPyPyPy- γ -HpImHpPy
	87) 5'-W T A C G W-3'	HpPyPyIm- γ -PyImHpPy
35	88) 5'-W T A C C W-3'	HpPyPyPy- γ -ImImHpPy

TABLE 7: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	89) 5'-W T G T T W-3'	HpImHpHp- γ -PyPyPyPy
5	90) 5'-W T G T A W-3'	HpImHpPy- γ -HpPyPyPy
	91) 5'-W T G T G W-3'	HpImHpIm- γ -PyPyPyPy
	92) 5'-W T G T C W-3'	HpImHpPy- γ -ImPyPyPy
	93) 5'-W T G A T W-3'	HpImPyHp- γ -PyHpPyPy
	94) 5'-W T G A A W-3'	HpImPyPy- γ -HpHpPyPy
10	95) 5'-W T G A G W-3'	HpImPyIm- γ -PyHpPyPy
	96) 5'-W T G A C W-3'	HpImPyPy- γ -ImHpPyPy
	97) 5'-W T G G T W-3'	HpImImHp- γ -PyPyPyPy
	98) 5'-W T G G A W-3'	HpImImPy- γ -HpPyPyPy
	99) 5'-W T G C T W-3'	HpImPyHp- γ -PyImPyPy
	100) 5'-W T G C A W-3'	HpImPyPy- γ -HpImPyPy
	101) 5'-W T G G G W-3'	HpImImIm- γ -PyPyPyPy
	102) 5'-W T G G C W-3'	HpImImPy- γ -ImPyPyPy
	103) 5'-W T G C G W-3'	HpImPyIm- γ -PyImPyPy
	104) 5'-W T G C C W-3'	HpImPyPy- γ -ImImPyPy
	105) 5'-W T C T T W-3'	HpPyHpHp- γ -PyPyImPy
	106) 5'-W T C T A W-3'	HpPyHpPy- γ -HpPyImPy
	107) 5'-W T C T G W-3'	HpPyHpIm- γ -PyPyImPy
	108) 5'-W T C T C W-3'	HpPyHpPy- γ -ImPyImPy
	109) 5'-W T C A T W-3'	HpPyPyHp- γ -PyHpImPy
25	110) 5'-W T C A A W-3'	HpPyPyPy- γ -HpHpImPy
	111) 5'-W T C A G W-3'	HpPyPyIm- γ -PyHpImPy
	112) 5'-W T C A C W-3'	HpPyPyPy- γ -ImHpImPy
	113) 5'-W T C G T W-3'	HpPyImHp- γ -PyPyImPy
	114) 5'-W T C G A W-3'	HpPyImPy- γ -HpPyImPy
30	115) 5'-W T C C T W-3'	HpPyPyHp- γ -PyImImPy
	116) 5'-W T C C A W-3'	HpPyPyPy- γ -HpImImPy
	117) 5'-W T C G G W-3'	HpPyImIm- γ -PyPyImPy
	118) 5'-W T C G C W-3'	HpPyImPy- γ -ImPyImPy
	119) 5'-W T C C G W-3'	HpPyPyIm- γ -PyImImPy
35	120) 5'-W T C C C W-3'	HpPyPyPy- γ -ImImImPy

TABLE 8: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	121) 5'-W A T T T W-3'	PyHpHpHp- γ -PyPyPyHp
5	122) 5'-W A T T A W-3'	PyHpHpPy- γ -HpPyPyHp
	123) 5'-W A T T G W-3'	PyHpHpIm- γ -PyPyPyHp
	124) 5'-W A T T C W-3'	PyHpHpPy- γ -ImPyPyHp
	125) 5'-W A T A T W-3'	PyHpPyHp- γ -PyHpPyHp
	126) 5'-W A T A A W-3'	PyHpPyPy- γ -HpHpPyHp
10	127) 5'-W A T A G W-3'	PyHpPyIm- γ -PyHpPyHp
	128) 5'-W A T A C W-3'	PyHpPyPy- γ -ImHpPyHp
	129) 5'-W A T G T W-3'	PyHpImHp- γ -PyPyPyHp
	130) 5'-W A T G A W-3'	PyHpImPy- γ -HpPyPyHp
	131) 5'-W A T G G W-3'	PyHpImIm- γ -PyPyPyHp
	132) 5'-W A T G C W-3'	PyHpImPy- γ -ImPyPyHp
	133) 5'-W A T C T W-3'	PyHpPyHp- γ -PyImPyHp
	134) 5'-W A T C A W-3'	PyHpPyPy- γ -HpImPyHp
	135) 5'-W A T C G W-3'	PyHpPyIm- γ -PyImPyHp
	136) 5'-W A T C C W-3'	PyHpPyPy- γ -ImImPyHp
20	137) 5'-W A A T T W-3'	PyPyHpHp- γ -PyPyHpHp
	138) 5'-W A A T A W-3'	PyPyHpPy- γ -HpPyHpHp
	139) 5'-W A A T G W-3'	PyPyHpIm- γ -PyPyHpHp
	140) 5'-W A A T C W-3'	PyPyHpPy- γ -ImPyHpHp
	141) 5'-W A A A T W-3'	PyPyPyHp- γ -PyHpHpHp
25	142) 5'-W A A A A W-3'	PyPyPyPy- γ -HpHpHpHp
	143) 5'-W A A A G W-3'	PyPyPyIm- γ -PyHpHpHp
	144) 5'-W A A A C W-3'	PyPyPyPy- γ -ImHpHpHp
	145) 5'-W A A G T W-3'	PyPyImHp- γ -PyPyHpHp
	146) 5'-W A A G A W-3'	PyPyImPy- γ -HpPyHpHp
30	147) 5'-W A A G G W-3'	PyPyImIm- γ -PyPyHpHp
	148) 5'-W A A G C W-3'	PyPyImPy- γ -ImPyHpHp
	149) 5'-W A A C T W-3'	PyPyPyHp- γ -PyImHpHp
	150) 5'-W A A C A W-3'	PyPyPyPy- γ -HpImHpHp
	151) 5'-W A A C G W-3'	PyPyPyIm- γ -PyImHpHp
35	152) 5'-W A A C C W-3'	PyPyPyPy- γ -ImImHpHp

TABLE 9: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WASNNW-3'

	DNA sequence	aromatic amino acid sequence
	153) 5'-W A G T T W-3'	PyImHpHp- γ -PyPyPyHp
5	154) 5'-W A G T A W-3'	PyImHpPy- γ -HpPyPyHp
	155) 5'-W A G T G W-3'	PyImHpIm- γ -PyPyPyHp
	156) 5'-W A G T C W-3'	PyImHpPy- γ -ImPyPyHp
	157) 5'-W A G A T W-3'	PyImPyHp- γ -PyHpPyHp
	158) 5'-W A G A A W-3'	PyImPyPy- γ -HpHpPyHp
10	159) 5'-W A G A G W-3'	PyImPyIm- γ -PyHpPyHp
	160) 5'-W A G A C W-3'	PyImPyPy- γ -ImHpPyHp
	161) 5'-W A G G T W-3'	PyImImHp- γ -PyPyPyHp
	162) 5'-W A G G A W-3'	PyImImPy- γ -HpPyPyHp
	163) 5'-W A G C T W-3'	PyImPyHp- γ -PyImPyHp
	164) 5'-W A G C A W-3'	PyImPyPy- γ -HpImPyHp
	165) 5'-W A G G G W-3'	PyImImIm- γ -PyPyPyHp
	166) 5'-W A G G C W-3'	PyImImPy- γ -ImPyPyHp
	167) 5'-W A G C G W-3'	PyImPyIm- γ -PyImPyHp
	168) 5'-W A G C C W-3'	PyImPyPy- γ -ImImPyHp
20	169) 5'-W A C T T W-3'	PyPyHpHp- γ -PyPyImHp
	170) 5'-W A C T A W-3'	PyPyHpPy- γ -HpPyImHp
	171) 5'-W A C T G W-3'	PyPyHpIm- γ -PyPyImHp
	172) 5'-W A C T C W-3'	PyPyHpPy- γ -ImPyImHp
	173) 5'-W A C A T W-3'	PyPyPyHp- γ -PyHpImHp
25	174) 5'-W A C A A W-3'	PyPyPyPy- γ -HpHpImHp
	175) 5'-W A C A G W-3'	PyPyPyIm- γ -PyHpImHp
	176) 5'-W A C A C W-3'	PyPyPyPy- γ -ImHpImHp
	177) 5'-W A C G T W-3'	PyPyImHp- γ -PyPyImHp
	178) 5'-W A C G A W-3'	PyPyImPy- γ -HpPyImHp
30	179) 5'-W A C C T W-3'	PyPyPyHp- γ -PyImImHp
	180) 5'-W A C C A W-3'	PyPyPyPy- γ -HpImImHp
	181) 5'-W A C G G W-3'	PyPyImIm- γ -PyPyImHp
	182) 5'-W A C G C W-3'	PyPyImPy- γ -ImPyImHp
	183) 5'-W A C C G W-3'	PyPyPyIm- γ -PyImImHp
35	184) 5'-W A C C C W-3'	PyPyPyPy- γ -ImImImHp

TABLE 10: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	185) 5'-W C T T T W-3'	PyHpHpHp- γ -PyPyPyIm
5	186) 5'-W C T T A W-3'	PyHpHpPy- γ -HpPyPyIm
	187) 5'-W C T T G W-3'	PyHpHpIm- γ -PyPyPyIm
	188) 5'-W C T T C W-3'	PyHpHpPy- γ -ImPyPyIm
	189) 5'-W C T A T W-3'	PyHpPyHp- γ -PyHpPyIm
	190) 5'-W C T A A W-3'	PyHpPyPy- γ -HpHpPyIm
10	191) 5'-W C T A G W-3'	PyHpPyIm- γ -PyHpPyIm
	192) 5'-W C T A C W-3'	PyHpPyPy- γ -ImHpPyIm
	193) 5'-W C T G T W-3'	PyHpImHp- γ -PyPyPyIm
	194) 5'-W C T G A W-3'	PyHpImPy- γ -HpPyPyIm
	195) 5'-W C T G G W-3'	PyHpImIm- γ -PyPyPyIm
15	196) 5'-W C T G C W-3'	PyHpImPy- γ -ImPyPyIm
	197) 5'-W C T C T W-3'	PyHpPyHp- γ -PyImPyIm
	198) 5'-W C T C A W-3'	PyHpPyPy- γ -HpImPyIm
	199) 5'-W C T C G W-3'	PyHpPyIm- γ -PyImPyIm
	200) 5'-W C T C C W-3'	PyHpPyPy- γ -ImImPyIm
20	201) 5'-W C A T T W-3'	PyPyHpHp- γ -PyPyHpIm
	202) 5'-W C A T A W-3'	PyPyHpPy- γ -HpPyHpIm
	203) 5'-W C A T G W-3'	PyPyHpIm- γ -PyPyHpIm
	204) 5'-W C A T C W-3'	PyPyHpPy- γ -ImPyHpIm
	205) 5'-W C A A T W-3'	PyPyPyHp- γ -PyHpHpIm
25	206) 5'-W C A A A W-3'	PyPyPyPy- γ -HpHpHpIm
	207) 5'-W C A A G W-3'	PyPyPyIm- γ -PyHpHpIm
	208) 5'-W C A A C W-3'	PyPyPyPy- γ -ImHpHpIm
	209) 5'-W C A G T W-3'	PyPyImHp- γ -PyPyHpIm
	210) 5'-W C A G A W-3'	PyPyImPy- γ -HpPyHpIm
30	211) 5'-W C A G G W-3'	PyPyImIm- γ -PyPyHpIm
	212) 5'-W C A G C W-3'	PyPyImPy- γ -ImPyHpIm
	213) 5'-W C A C T W-3'	PyPyPyHp- γ -PyImHpIm
	214) 5'-W C A C A W-3'	PyPyPyPy- γ -HpImHpIm
	215) 5'-W C A C G W-3'	PyPyPyIm- γ -PyImHpIm
35	216) 5'-W C A C C W-3'	PyPyPyPy- γ -ImImHpIm

TABLE 11: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	217) 5'-W C G T T W-3'	PyImHpHp- γ -PyPyPyIm
5	218) 5'-W C G T A W-3'	PyImHpPy- γ -HpPyPyIm
	219) 5'-W C G T G W-3'	PyImHpIm- γ -PyPyPyIm
	220) 5'-W C G T C W-3'	PyImHpPy- γ -ImPyPyIm
	221) 5'-W C G A T W-3'	PyImPyHp- γ -PyHpPyIm
	222) 5'-W C G A A W-3'	PyImPyPy- γ -HpHpPyIm
10	223) 5'-W C G A G W-3'	PyImPyIm- γ -PyHpPyIm
	224) 5'-W C G A C W-3'	PyImPyPy- γ -ImHpPyIm
	225) 5'-W C G G T W-3'	PyImImHp- γ -PyPyPyIm
	226) 5'-W C G G A W-3'	PyImImPy- γ -HpPyPyIm
	227) 5'-W C G C T W-3'	PyImPyHp- γ -PyImPyIm
	228) 5'-W C G C A W-3'	PyImPyPy- γ -HpImPyIm
	229) 5'-W C C T T W-3'	PyPyHpHp- γ -PyPyImIm
	230) 5'-W C C T A W-3'	PyPyHpPy- γ -HpPyImIm
	231) 5'-W C C T G W-3'	PyPyHpIm- γ -PyPyImIm
	232) 5'-W C C T C W-3'	PyPyHpPy- γ -ImPyImIm
20	233) 5'-W C C A T W-3'	PyPyPyHp- γ -PyHpImIm
	234) 5'-W C C A A W-3'	PyPyPyPy- γ -HpHpImIm
	235) 5'-W C C A G W-3'	PyPyPyIm- γ -PyHpImIm
	236) 5'-W C C A C W-3'	PyPyPyPy- γ -ImHpImIm
	237) 5'-W C C G T W-3'	PyPyImHp- γ -PyPyImIm
25	238) 5'-W C C G A W-3'	PyPyImPy- γ -HpPyImIm
	239) 5'-W C C C T W-3'	PyPyPyHp- γ -PyImImIm
	240) 5'-W C C C A W-3'	PyPyPyPy- γ -HpImImIm
	G9) 5'-W C G G G W-3'	PyImImIm- γ -PyPyPyIm
	G10) 5'-W C G G C W-3'	PyImImPy- γ -ImPyPyIm
30	G11) 5'-W C G C G W-3'	PyImPyIm- γ -PyImPyIm
	G12) 5'-W C G C C W-3'	PyImPyPy- γ -ImImPyIm
	G13) 5'-W C C G G W-3'	PyPyImIm- γ -PyPyImIm
	G14) 5'-W C C G C W-3'	PyPyImPy- γ -ImPyImIm
	G15) 5'-W C C C G W-3'	PyPyPyIm- γ -PyImImIm
35	G16) 5'-W C C C C W-3'	PyPyPyPy- γ -ImImImIm

TABLE 12: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WGWNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
5	3 β) 5'-W G T T G W-3'	ImHp- β -Im- γ -PyPyPyPy
	7 β) 5'-W G T A G W-3'	ImHp- β -Im- γ -PyHpPyPy
	9 β) 5'-W G T G T W-3'	Im- β -ImHp- γ -PyPyPyPy
	10 β) 5'-W G T G A W-3'	Im- β -ImPy- γ -HpPyPyPy
	11 β) 5'-W G T G G W-3'	Im- β -ImIm- γ -PyPyPyPy
10	12 β) 5'-W G T G C W-3'	Im- β -ImPy- γ -ImPyPyPy
	15 β) 5'-W G T C G W-3'	ImHp- β -Im- γ -PyImPyPy
	19 β) 5'-W G A T G W-3'	ImPy- β -Im- γ -PyPyHpPy
	23 β) 5'-W G A A G W-3'	ImPy- β -Im- γ -PyHpHpPy
	25 β) 5'-W G A G T W-3'	Im- β -ImHp- γ -PyPyHpPy
	26 β) 5'-W G A G A W-3'	Im- β -ImPy- γ -HpPyHpPy
	27 β) 5'-W G A G G W-3'	Im- β -ImIm- γ -PyPyHpPy
	28 β) 5'-W G A G C W-3'	Im- β -ImPy- γ -ImPyHpPy
	31 β) 5'-W G A C G W-3'	ImPy- β -Im- γ -PyImHpPy

TABLE 13: 8-ring Harpin Polyamides for recognition of 6-bp 5'-WGSNNW-3' with β -substitutions included

	DNA sequence	aromatic amino acid sequence
	35 β) 5'-W G G T G W-3'	ImIm- β -Im- γ -PyPyPyPy
5	39 β) 5'-W G G A G W-3'	ImIm- β -Im- γ -PyHpPyPy
	45 β) 5'-W G C T T W-3'	ImPyHpHp- γ -Py- β -ImPy
	46 β) 5'-W G C T A W-3'	ImPyHpPy- γ -Hp- β -ImPy
	47 β) 5'-W G C T G W-3'	ImPyHpIm- γ -Py- β -ImPy
	47 β 2) 5'-W G C T G W-3'	ImPy- β -Im- γ -Py- β -ImPy
10	48 β) 5'-W G C T C W-3'	ImPyHpPy- γ -Im- β -ImPy
	49 β) 5'-W G C A T W-3'	ImPyPyHp- γ -Py- β -ImPy
	50 β) 5'-W G C A A W-3'	ImPyPyPy- γ -Hp- β -ImPy
	51 β) 5'-W G C A G W-3'	ImPyPyIm- γ -Py- β -ImPy
	51 β 2) 5'-W G C A G W-3'	ImPy- β -Im- γ -Py- β -ImPy
15	52 β) 5'-W G C A C W-3'	ImPyPyPy- γ -Im- β -ImPy
	53 β) 5'-W G C G T W-3'	ImPyImHp- γ -Py- β -ImPy
	53 β 2) 5'-W G C G T W-3'	Im- β -ImHp- γ -Py- β -ImPy
	54 β) 5'-W G C G A W-3'	ImPyImPy- γ -Hp- β -ImPy
	54 β 2) 5'-W G C G A W-3'	Im- β -ImPy- γ -Hp- β -ImPy
20	G3 β) 5'-W G G C G W-3'	ImIm- β -Im- γ -PyImPyPy
	G5 β) 5'-W G C G G W-3'	ImPyImIm- γ -Py- β -ImPy
	G5 β 2) 5'-W G C G G W-3'	Im- β -ImIm- γ -Py- β -ImPy
	G6 β) 5'-W G C G C W-3'	ImPyImPy- γ -Im- β -ImPy
	G6 β 2) 5'-W G C G C W-3'	Im- β -ImPy- γ -Im- β -ImPy
25	G7 β) 5'-W G C C G W-3'	ImPy- β -Im- γ -PyImImPy

TABLE 14: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WTWNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
	59 β) 5'-W T T T G W-3'	HpHp- β -Im- γ -PyPyPyPy
5	63 β) 5'-W T T A G W-3'	HpHp- β -Im- γ -PyHpPyPy
	65 β) 5'-W T T G T W-3'	Hp- β -ImHp- γ -PyPyPyPy
	66 β) 5'-W T T G A W-3'	Hp- β -ImPy- γ -HpPyPyPy
	67 β) 5'-W T T G G W-3'	Hp- β -ImIm- γ -PyPyPyPy
	68 β) 5'-W T T G C W-3'	Hp- β -ImPy- γ -ImPyPyPy
10	71 β) 5'-W T T C G W-3'	HpHp- β -Im- γ -PyImPyPy
	75 β) 5'-W T A T G W-3'	HpPy- β -Im- γ -PyPyHpPy
	79 β) 5'-W T A A G W-3'	HpPy- β -Im- γ -PyHpHpPy
	81 β) 5'-W T A G T W-3'	Hp- β -ImHp- γ -PyPyHpPy
	82 β) 5'-W T A G A W-3'	Hp- β -ImPy- γ -HpPyHpPy
	83 β) 5'-W T A G G W-3'	Hp- β -ImIm- γ -PyPyHpPy
	84 β) 5'-W T A G C W-3'	Hp- β -ImPy- γ -ImPyHpPy
	87 β) 5'-W T A C G W-3'	HpPy- β -Im- γ -PyImHpPy

TABLE 15: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WTSNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
	91 β) 5'-W T G T G W-3'	HpIm- β -Im- γ -PyPyPyPy
5	95 β) 5'-W T G A G W-3'	HpIm- β -Im- γ -PyHpPyPy
	103 β) 5'-W T G C G W-3'	HpIm- β -Im- γ -PyImPyPy
	105 β) 5'-W T C T T W-3'	HpPyHpHp- γ -Py- β -ImPy
	106 β) 5'-W T C T A W-3'	HpPyHpPy- γ -Hp- β -ImPy
	107 β) 5'-W T C T G W-3'	HpPyHpIm- γ -Py- β -ImPy
10	107 β 2) 5'-W T C T G W-3'	HpPy- β -Im- γ -Py- β -ImPy
	108 β) 5'-W T C T C W-3'	HpPyHpPy- γ -Im- β -ImPy
	109 β) 5'-W T C A T W-3'	HpPyPyHp- γ -Py- β -ImPy
	110 β) 5'-W T C A A W-3'	HpPyPyPy- γ -Hp- β -ImPy
	111 β) 5'-W T C A G W-3'	HpPyPyIm- γ -Py- β -ImPy
15	111 β 2) 5'-W T C A G W-3'	HpPy- β -Im- γ -Py- β -ImPy
	112 β) 5'-W T C A C W-3'	HpPyPyPy- γ -Im- β -ImPy
	113 β) 5'-W T C G T W-3'	HpPyImHp- γ -Py- β -ImPy
	113 β 2) 5'-W T C G T W-3'	Hp- β -ImHp- γ -Py- β -ImPy
	114 β) 5'-W T C G A W-3'	HpPyImPy- γ -Hp- β -ImPy
20	114 β 2) 5'-W T C G A W-3'	Hp- β -ImPy- γ -Hp- β -ImPy
	117 β) 5'-W T C G G W-3'	HpPyImIm- γ -Py- β -ImPy
	117 β 2) 5'-W T C G G W-3'	Hp- β -ImIm- γ -Py- β -ImPy
	118 β) 5'-W T C G C W-3'	HpPyImPy- γ -Im- β -ImPy
	118 β 2) 5'-W T C G C W-3'	Hp- β -ImPy- γ -Im- β -ImPy
25	119 β) 5'-W T C C G W-3'	HpPy- β -Im- γ -PyImImPy

TABLE 16: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WAWNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
5	123 β) 5'-W A T T G W-3'	PyHp- β -Im- γ -PyPyPyHp
	127 β) 5'-W A T A G W-3'	PyHp- β -Im- γ -PyHpPyHp
	129 β) 5'-W A T G T W-3'	Py- β -ImHp- γ -PyPyPyHp
	130 β) 5'-W A T G A W-3'	Py- β -ImPy- γ -HpPyPyHp
	131 β) 5'-W A T G G W-3'	Py- β -ImIm- γ -PyPyPyHp
10	132 β) 5'-W A T G C W-3'	Py- β -ImPy- γ -ImPyPyHp
	135 β) 5'-W A T C G W-3'	PyHp- β -Im- γ -PyImPyHp
	139 β) 5'-W A A T G W-3'	PyPy- β -Im- γ -PyPyHpHp
	143 β) 5'-W A A A G W-3'	PyPy- β -Im- γ -PyHpHpHp
	145 β) 5'-W A A G T W-3'	Py- β -ImHp- γ -PyPyHpHp
	146 β) 5'-W A A G A W-3'	Py- β -ImPy- γ -HpPyHpHp
	147 β) 5'-W A A G G W-3'	Py- β -ImIm- γ -PyPyHpHp
	148 β) 5'-W A A G C W-3'	Py- β -ImPy- γ -ImPyHpHp
	151 β) 5'-W A A C G W-3'	PyPy- β -Im- γ -PyImHpHp

TABLE 17: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WASNNW-3' with β -substitutions included

	DNA sequence	aromatic amino acid sequence
	155 β) 5'-W A G T G W-3'	PyIm- β -Im- γ -PyPyPyHp
5	159 β) 5'-W A G A G W-3'	PyIm- β -Im- γ -PyHpPyHp
	167 β) 5'-W A G C G W-3'	PyIm- β -Im- γ -PyImPyHp
	169 β) 5'-W A C T T W-3'	PyPyHpHp- γ -Py- β -ImHp
	170 β) 5'-W A C T A W-3'	PyPyHpPy- γ -Hp- β -ImHp
	171 β) 5'-W A C T G W-3'	PyPyHpIm- γ -Py- β -ImHp
10	171 β 2) 5'-W A C T G W-3'	PyPy- β -Im- γ -Py- β -ImHp
	172 β) 5'-W A C T C W-3'	PyPyHpPy- γ -Im- β -ImHp
	173 β) 5'-W A C A T W-3'	PyPyPyHp- γ -Py- β -ImHp
	174 β) 5'-W A C A A W-3'	PyPyPyPy- γ -Hp- β -ImHp
	175 β) 5'-W A C A G W-3'	PyPyPyIm- γ -Py- β -ImHp
	175 β 2) 5'-W A C A G W-3'	PyPy- β -Im- γ -Py- β -ImHp
	176 β) 5'-W A C A C W-3'	PyPyPyPy- γ -Im- β -ImHp
	177 β) 5'-W A C G T W-3'	PyPyImHp- γ -Py- β -ImHp
	177 β 2) 5'-W A C G T W-3'	Py- β -ImHp- γ -Py- β -ImHp
	178 β) 5'-W A C G A W-3'	PyPyImPy- γ -Hp- β -ImHp
20	178 β 2) 5'-W A C G A W-3'	Py- β -ImPy- γ -Hp- β -ImHp
	181 β) 5'-W A C G G W-3'	PyPyImIm- γ -Py- β -ImHp
	181 β 2) 5'-W A C G G W-3'	Py- β -ImIm- γ -Py- β -ImHp
	182 β) 5'-W A C G C W-3'	PyPyImPy- γ -Im- β -ImHp
	182 β 2) 5'-W A C G C W-3'	Py- β -ImPy- γ -Im- β -ImHp
25	183 β 2) 5'-W A C C G W-3'	PyPy- β -Im- γ -PyImImHp

TABLE 18: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WCWNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
	185 β) 5'-W C T T T W-3'	PyHpHpHp- γ -PyPy- β -Im
5	186 β) 5'-W C T T A W-3'	PyHpHpPy- γ -HpPy- β -Im
	187 β) 5'-W C T T G W-3'	PyHpHpIm- γ -PyPy- β -Im
	187 β 2) 5'-W C T T G W-3'	PyHp- β -Im- γ -PyPy- β -Im
	188 β) 5'-W C T T C W-3'	PyHpHpPy- γ -ImPy- β -Im
	189 β) 5'-W C T A T W-3'	PyHpPyHp- γ -PyHp- β -Im
10	190 β) 5'-W C T A A W-3'	PyHpPyPy- γ -HpHp- β -Im
	191 β) 5'-W C T A G W-3'	PyHpPyIm- γ -PyHp- β -Im
	191 β 2) 5'-W C T A G W-3'	PyHp- β -Im- γ -PyHp- β -Im
	192 β) 5'-W C T A C W-3'	PyHpPyPy- γ -ImHp- β -Im
	193 β) 5'-W C T G T W-3'	PyHpImHp- γ -PyPy- β -Im
	193 β 2) 5'-W C T G T W-3'	Py- β -ImHp- γ -PyPy- β -Im
	194 β) 5'-W C T G A W-3'	PyHpImPy- γ -HpPy- β -Im
	194 β 2) 5'-W C T G A W-3'	Py- β -ImPy- γ -HpPy- β -Im
	195 β) 5'-W C T G G W-3'	PyHpImIm- γ -PyPy- β -Im
	195 β 2) 5'-W C T G G W-3'	Py- β -ImIm- γ -PyPy- β -Im
20	196 β) 5'-W C T G C W-3'	PyHpImPy- γ -ImPy- β -Im
	196 β 2) 5'-W C T G C W-3'	Py- β -ImPy- γ -ImPy- β -Im
	197 β) 5'-W C T C T W-3'	PyHpPyHp- γ -PyIm- β -Im
	198 β) 5'-W C T C A W-3'	PyHpPyPy- γ -HpIm- β -Im
	199 β) 5'-W C T C G W-3'	PyHpPyIm- γ -PyIm- β -Im
25	199 β 2) 5'-W C T C G W-3'	PyHp- β -Im- γ -PyIm- β -Im
	200 β) 5'-W C T C C W-3'	PyHpPyPy- γ -ImIm- β -Im
	201 β) 5'-W C A T T W-3'	PyPyHpHp- γ -PyPy- β -Im
	202 β) 5'-W C A T A W-3'	PyPyHpPy- γ -HpPy- β -Im
	203 β) 5'-W C A T G W-3'	PyPyHpIm- γ -PyPy- β -Im
30	203 β 2) 5'-W C A T G W-3'	PyPy- β -Im- γ -PyPy- β -Im
	204 β) 5'-W C A T C W-3'	PyPyHpPy- γ -ImPy- β -Im
	205 β) 5'-W C A A T W-3'	PyPyPyHp- γ -PyHp- β -Im
	206 β) 5'-W C A A A W-3'	PyPyPyPy- γ -HpHp- β -Im

TABLE 18 (cont): 8-ring Hairpin Polyamides for 6-bp 5'-WCWNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
	207 β) 5'-W C A A G W-3'	PyPyPyIm- γ -PyHp- β -Im
5	207 β 2) 5'-W C A A G W-3'	PyPy- β -Im- γ -PyHp- β -Im
	208 β) 5'-W C A A C W-3'	PyPyPyPy- γ -ImHp- β -Im
	209 β) 5'-W C A G T W-3'	PyPyImHp- γ -PyPy- β -Im
	209 β 2) 5'-W C A G T W-3'	Py- β -ImHp- γ -PyPy- β -Im
	210 β) 5'-W C A G A W-3'	PyPyImPy- γ -HpPy- β -Im
10	210 β 2) 5'-W C A G A W-3'	Py- β -ImPy- γ -HpPy- β -Im
	211 β) 5'-W C A G G W-3'	PyPyImIm- γ -PyPy- β -Im
	211 β 2) 5'-W C A G G W-3'	Py- β -ImIm- γ -PyPy- β -Im
	212 β) 5'-W C A G C W-3'	PyPyImPy- γ -ImPy- β -Im
	212 β 2) 5'-W C A G C W-3'	Py- β -ImPy- γ -ImPy- β -Im
	213 β) 5'-W C A C T W-3'	PyPyPyHp- γ -PyIm- β -Im
	214 β) 5'-W C A C A W-3'	PyPyPyPy- γ -HpIm- β -Im
	215 β) 5'-W C A C G W-3'	PyPyPyIm- γ -PyIm- β -Im
	215 β 2) 5'-W C A C G W-3'	PyPy- β -Im- γ -PyIm- β -Im
	216 β) 5'-W C A C C W-3'	PyPyPyPy- γ -ImIm- β -Im

TABLE 19: 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WCSNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
	217 β) 5'-W C G T T W-3'	PyImHpHp- γ -PyPy- β -Im
5	218 β) 5'-W C G T A W-3'	PyImHpPy- γ -HpPy- β -Im
	219 β) 5'-W C G T G W-3'	PyImHpIm- γ -PyPy- β -Im
	219 β 2) 5'-W C G T G W-3'	PyIm- β -Im- γ -PyPy- β -Im
	220 β) 5'-W C G T C W-3'	PyImHpPy- γ -ImPy- β -Im
	221 β) 5'-W C G A T W-3'	PyImPyHp- γ -PyHp- β -Im
10	222 β) 5'-W C G A A W-3'	PyImPyPy- γ -HpHp- β -Im
	223 β) 5'-W C G A G W-3'	PyImPyIm- γ -PyHp- β -Im
	223 β 2) 5'-W C G A G W-3'	PyIm- β -Im- γ -PyHp- β -Im
	224 β) 5'-W C G A C W-3'	PyImPyPy- γ -ImHp- β -Im
	225 β) 5'-W C G G T W-3'	PyImImHp- γ -PyPy- β -Im
	226 β) 5'-W C G G A W-3'	PyImImPy- γ -HpPy- β -Im
	227 β) 5'-W C G C T W-3'	PyImPyHp- γ -PyIm- β -Im
	228 β) 5'-W C G C A W-3'	PyImPyPy- γ -HpIm- β -Im
	229 β) 5'-W C C T T W-3'	PyPyHpHp- γ -Py- β -ImIm
	230 β) 5'-W C C T A W-3'	PyPyHpPy- γ -Hp- β -ImIm
20	231 β) 5'-W C C T G W-3'	PyPyHpIm- γ -Py- β -ImIm
	231 β 2) 5'-W C C T G W-3'	PyPy- β -Im- γ -Py- β -ImIm
	232 β) 5'-W C C T C W-3'	PyPyHpPy- γ -Im- β -ImIm
	233 β) 5'-W C C A T W-3'	PyPyPyHp- γ -Py- β -ImIm
	234 β) 5'-W C C A A W-3'	PyPyPyPy- γ -Hp- β -ImIm
25	235 β) 5'-W C C A G W-3'	PyPyPyIm- γ -Py- β -ImIm
	235 β 2) 5'-W C C A G W-3'	PyPy- β -Im- γ -Py- β -ImIm
	236 β) 5'-W C C A C W-3'	PyPyPyPy- γ -Im- β -ImIm
	237 β) 5'-W C C G T W-3'	PyPyImHp- γ -Py- β -ImIm
	237 β 2) 5'-W C C G T W-3'	Py- β -ImHp- γ -Py- β -ImIm
30	238 β) 5'-W C C G A W-3'	PyPyImPy- γ -Hp- β -ImIm
	238 β 2) 5'-W C C G A W-3'	Py- β -ImPy- γ -Hp- β -ImIm
	G9 β) 5'-W C G G G W-3'	PyImImIm- γ -PyPy- β -Im
	G10 β) 5'-W C G G C W-3'	PyImImPy- γ -ImPy- β -Im

TABLE 19 (cont): 8-ring Hairpin Polyamides for recognition of 6-bp 5'-WCSNNW-3' with β -substitutions included.

	DNA sequence	aromatic amino acid sequence
5	G11 β) 5'-W C G C G W-3'	PyImPyIm- γ -PyIm- β -Im
	G11 β 2) 5'-W C G C G W-3'	PyIm- β -Im- γ -PyIm- β -Im
	G12 β) 5'-W C G C C W-3'	PyImPyPy- γ -ImIm- β -Im
	G13 β) 5'-W C C G G W-3'	PyPyImIm- γ -Py- β -ImIm
	G13 β 2) 5'-W C C G G W-3'	Py- β -ImIm- γ -Py- β -ImIm
10	G14 β) 5'-W C C G C W-3'	PyPyImPy- γ -Im- β -ImIm
	G14 β 2) 5'-W C C G C W-3'	Py- β -ImPy- γ -Im- β -ImIm
	G15 β) 5'-W C C C G W-3'	PyPy- β -Im- γ -PyImImIm

If the process described above of designing a preferred polyamide molecule X₁X₂X₃X₄- γ -X₅X₆X₇X₈ comprising eight aromatic aminoacid residues does not produce a selective polyamide that binds to the target identified DNA sequence with subnanomolar affinity and with a selectivity over mismatch sequences of greater than a factor of ten, a polyamide molecule X₁X₂X₃X₄X₅- γ -X₆X₇X₈X₉X₁₀ having five carboxamide binding pairs can be designed that is selective for a seven base pair identified target 5'-WNNNNNW-3' sequence. The design and synthesis of the five binding pair polyamide is similar to that of the four binding pair polyamide X₁X₂X₃X₄- γ -X₅X₆X₇X₈ described above.

The polyamide design process, shown schematically in Figure 7 provides a method for designing a ten carboxamide residue molecule comprising five carboxamide binding pairs for selective detection and binding of a target seven base pair 5'-WNNNNNW-3' sequence in the minor groove of double stranded DNA. The design process identifies an appropriate polyamide ligand for recognition of a predetermined seven base pair, 5'-WNNNNNW-3' sequence with subnanomolar affinity and >10-fold specificity versus mismatch sites. Trauger, J.W., Baird, E. E. Dervan, P.B. describes the recognition of DNA by designed ligands at subnanomolar concentrations. *Nature* **382**, 559-561 (1996).

In order to prepare a polyamide molecule specific for an identified seven base pair sequence of double stranded DNA, a user starts the 10-ring hairpin design process that implements the minor groove recognition pairing code summarized in Table 2 above. In the

design process a 5'-WNNNNNW-3' sequence was identified. In a preferred embodiment, the identified sequence was located within a gene promoter. The identified sequence was then defined as 5'-W**abcde**W-3' in a stepwise process wherein **a**, **b**, **c**, **d**, and **e**, were sequentially and independently defined as A, G, C, or T. The structure of the polyamide molecule was then correspondingly defined by sequentially choosing antiparallel carboxamide binding pairs according to the minor groove pairing code summarized in Table 2 above. Thus, if **a** was G, then X₁ was defined as Im, and X₁₀ was defined as Py. If **a** was C, then X₁ was defined as Py, and X₁₀ was defined as Im. If **a** was T, then X₁ was defined as Hp, and X₁₀ was defined as Py. If **a** was A, then X₁ was defined as Py, and X₁₀ was defined as Hp.

Similarly, **b** was defined as A, G, C, or T and corresponding carboxamide binding pairs were defined. According to the same rules, if **b** was G, then X₂ was defined as Im, and X₉ was defined as Py. If **b** was C, then X₂ was defined as Py, and X₉ was defined as Im. Likewise, if **b** was T, then X₂ was defined as Hp, and X₉ was defined as Py. If **b** was A, then X₂ was defined as Py, and X₉ was defined as Hp.

The next step was to define **c** as A, G, C, or T and then define corresponding carboxamide binding pairs. Following the same rules, if **c** was G, then X₃ was defined as Im, and X₈ was defined as Py. If **c** was C, then X₃ was defined as Py, and X₈ was defined as Im. Similarly, if **c** was T, then X₃ was defined as Hp, and X₈ was defined as Py. If **c** was A, then X₃ was defined as Py, and X₈ was defined as Hp. Similarly, **d** was defined as A, G, C, or T and the corresponding carboxamide binding pair was defined. According to the above rules, if **d** was G, then X₄ was defined as Im, and X₇ was defined as Py. If **d** was C, then X₄ was defined as Py, and X₇ was defined as Im. If **d** was T, then X₄ was defined as Hp, and X₇ was defined as Py. If **d** was A, then X₄ was defined as Py, and X₇ was defined as Hp. Finally, **e** was defined as A, G, C, or T and the corresponding carboxamide binding pair was defined. According to the above rules, if **e** was G, then X₅ was defined as Im, and X₆ was defined as Py. If **e** was C, then X₅ was defined as Py, and X₆ was defined as Im. If **e** was T, then X₅ was defined as Hp, and X₆ was defined as Py. If **e** was A, then X₅ was defined as Py, and X₆ was defined as Hp.

With all ten carboxamide residues that participate in the binding pairs now defined, the designed polyamide X₁X₂X₃X₄X₅-γ-X₆X₇X₈X₉X₁₀ suitable for binding to the identified

sequence was synthesized using known techniques. Baird, E. E. & Dervan, P. B. describes the solid phase synthesis of polyamides containing imidazole and pyrrole amino acids. *J. Am. Chem. Soc.* **118**, 6141-6146 (1996); also see PCT US 97/003332.

5 The binding affinity of the synthesized polyamide to the identified sequence was determined using a quantitative DNase footprint titration method for studying protein-DNA interactions described by Brenowitz, M., Senear, D. F., Shea, M. A. & Ackers, G. K., *Methods Enzymol.* **130**, 132-181 (1986). If the affinity of the synthesized polyamide at the target site was not subnanomolar affinity then substituting at least one β -alanine residue for a pyrrole or 3-
10 hydroxypyrrole residue was considered in order to optimize the exact positions of the binding pairs of aromatic amino acids. If the affinity of the polyamide at the target site was subnanomolar affinity then the sequence specificity of the polyamide versus mismatch sequences was determined. If the specificity versus mismatch sites was not > 10-fold specificity then adding a β -alanine (shown schematically in Figure 8) was considered, in order to optimize the positions of the aromatic amino acids in relationship to the base pairs in the minor groove. Specificity of the polyamide molecule for the target identified sequence versus mismatch sequence sites of greater than 10-fold was considered a successful result of design process.

20 The 1024 polyamide molecules comprising five carboxamide binding pairs that were designed using this method are useful for binding to the 1024 target 5'-NNNNN-3' core sequences, and are listed in Tables 20-51. A corresponding polyamide molecule was designed for each DNA sequence (241-1232) and (G17-G48) using the process outlined above and shown schematically in Figure 7.

25 If the synthesized polyamide molecule did not bind to the target identified sequence with subnanomolar affinity or if the synthesized polyamide molecule did not exhibit a specificity for the target identified sequence versus mismatch sequence sites of greater than 10-fold, the option of substituting an aliphatic amino acid residue for one of the carboxamide residues was considered. The preferred aliphatic amino acid residue is β -alanine. At least one aliphatic
30 amino acid residue such as a β -alanine residue provided some flexibility to the central portion of the polyamide molecule, acting as a "spring" to permit optimization of the hydrogen bonding between the carboxamide binding pairs and the nucleotide bases of the double stranded DNA.

In general, it was not found to be advantageous to replace either member of the terminal carboxamide binding pair, X₁/X₁₀, with β -alanine. Similarly, β -alanine was not substituted for members of the binding pair, X₅/X₆, adjacent to the γ hairpin residue. β -alanine residues were not substituted for N-methylimidazole residues. The use of β -alanine in place of a pyrrole or 3-hydroxypyrrole amino acid residue provides aromatic/aliphatic pairing (Im/ β , β /Im, Hp/ β , β /Hp, Py/ β , and β /Py) and aliphatic/aliphatic pairing (β / β) substitution.

The method for selecting which pyrrole amino acid to substitute with β -alanine is schematically illustrated in Figure 8. Selective placement of an aliphatic β -alanine (β) residue paired with either a pyrrole (Py), 3-hydroxypyrrole (Hp), or imidazole (Im) aromatic amino acid or another β -alanine residue is found to compensate for sequence composition effects to improve recognition and binding of the minor groove of DNA by pyrrole-imidazole polyamides of the present invention. If an all-ring polyamide has been found to have an affinity which is not subnanomolar, or a specificity versus mismatch sequences which is less than 10-fold it may be caused by DNA sequence-composition effects which can be reduced by replacement of an aromatic amino acid with an aliphatic β -alanine residue. In a polyamide molecule that comprises five binding pairs it is only beneficial to place β -alanine in positions X₂, X₃, X₄, X₇, X₈, and X₉. No more than two β -alanine residues may be placed within a single hairpin structure. No more than a single β -residue may be placed within each individual polyamide subunit, e.g., if X₂ is replaced with β -alanine, X₃ or X₄ cannot be replaced as well.

These rules and others were implemented in the method schematically illustrated in Figure 8. This process is suitable for the refinement of the design polyamide comprising five binding pairs that has been designed by the method illustrated in Figure 7, but which lacks subnanomolar affinity or greater than 10-fold specificity at the identified target DNA sequence. As in the basic design method, the designed polyamides are synthesized and the affinity and specificity of binding to the target DNA were determined.

As discussed above, for a given 10-ring polyamide molecule there are six possible outcomes for the process of substituting a β -alanine residue for an aromatic amino acid residue. First, there may be no position at which it is possible to add a β -alanine residue; in such case, a better binding affinity or selectivity can be sought in the design and synthesis of a polyamide

with four or six carboxamide binding pairs, described below. Second, the process may result in a derivative which contains a single β -alanine substitution (such derivatives are numbered according to the parent numbering scheme such that a single β -derivative of compound 5 would be called 5 β), which is sufficient to produce subnanomolar binding affinity and >10-fold specificity, and at which point the process is deemed complete.

Third, the process of Figure 8 may result in a polyamide which contains a single β -alanine substitution which is not sufficient to produce subnanomolar binding affinity and >10-fold specificity, but where there are no additional positions in which it is possible to substitute a β -alanine residue, and in such a case a paired β -alanine residue should be added as described in Figure 9 and text below. Fourth, the process of Figure 7 may result in a polyamide that contains a single β -alanine substitution that is not sufficient to produce subnanomolar binding affinity and >10-fold specificity, but where there is an additional position for β -alanine substitution that does produce a polyamide with the criterion level of affinity and selectivity. Tables 52-83 list polyamide compounds 241 β -1232 β and G17 β -G48 β , corresponding to DNA sequences 241-1232 and G17 – G48, that contain one or two β -alanine residues.

A fifth possibility is that substitution at a second position by the method illustrated in Figure 9 with a paired β -alanine residue is not sufficient to produce a polyamide having the subnanomolar binding affinity and >10-fold specificity, and a polyamide with four or six carboxamide binding pairs, should be designed and synthesized, as described below. Finally, the design process may result in a polyamide that has a paired β -alanine substitution that is sufficient to produce subnanomolar binding affinity and >10-fold specificity, and therefore the design process is deemed complete. Tables 52-83 list polyamide compounds 241 β -1232 β and G17 β -G48 β , corresponding to DNA sequences 241-1232 and G17 – G48, that contain one or two β -alanine residues. In addition, Tables 52-83 list polyamides corresponding to sequences (241-1232) and (G17-G48) labeled (241 β p-1232 β p) and (G17 β p-G48 β p) that contain paired β/β residues added by the process described in Figure 9.

TABLE 20: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	241) 5'-W G G T T T W-3'	ImImHpHpHp- γ -PyPyPyPyPy
5	242) 5'-W G G T T A W-3'	ImImHpHpPy- γ -HpPyPyPyPy
	243) 5'-W G G T T G W-3'	ImImHpHpIm- γ -PyPyPyPyPy
	244) 5'-W G G T T C W-3'	ImImHpHpPy- γ -ImPyPyPyPy
	245) 5'-W G G T A T W-3'	ImImHpPyHp- γ -PyHpPyPyPy
	246) 5'-W G G T A A W-3'	ImImHpPyPy- γ -HpHpPyPyPy
10	247) 5'-W G G T A G W-3'	ImImHpPyIm- γ -PyHpPyPyPy
	248) 5'-W G G T A C W-3'	ImImHpPyPy- γ -ImHpPyPyPy
	249) 5'-W G G T G T W-3'	ImImHpImHp- γ -PyPyPyPyPy
	250) 5'-W G G T G A W-3'	ImImHpImPy- γ -HpPyPyPyPy
	251) 5'-W G G T G G W-3'	ImImHpImIm- γ -PyPyPyPyPy
	252) 5'-W G G T G C W-3'	ImImHpImPy- γ -ImPyPyPyPy
	253) 5'-W G G T C T W-3'	ImImHpPyHp- γ -PyImPyPyPy
	254) 5'-W G G T C A W-3'	ImImHpPyPy- γ -HpImPyPyPy
	255) 5'-W G G T C G W-3'	ImImHpPyIm- γ -PyImPyPyPy
	256) 5'-W G G T C C W-3'	ImImHpPyPy- γ -ImImPyPyPy
	257) 5'-W G G A T T W-3'	ImImPyHpHp- γ -PyPyHpPyPy
	258) 5'-W G G A T A W-3'	ImImPyHpPy- γ -HpPyHpPyPy
	259) 5'-W G G A T G W-3'	ImImPyHpIm- γ -PyPyHpPyPy
	260) 5'-W G G A T C W-3'	ImImPyHpPy- γ -ImPyHpPyPy
	261) 5'-W G G A A T W-3'	ImImPyPyHp- γ -PyHpHpPyPy
25	262) 5'-W G G A A A W-3'	ImImPyPyPy- γ -HpHpHpPyPy
	263) 5'-W G G A A G W-3'	ImImPyPyIm- γ -PyHpHpPyPy
	264) 5'-W G G A A C W-3'	ImImPyPyPy- γ -ImHpHpPyPy
	265) 5'-W G G A G T W-3'	ImImPyImHp- γ -PyPyHpPyPy
	266) 5'-W G G A G A W-3'	ImImPyImPy- γ -HpPyHpPyPy
30	267) 5'-W G G A G G W-3'	ImImPyImIm- γ -PyPyHpPyPy
	268) 5'-W G G A G C W-3'	ImImPyImPy- γ -ImPyHpPyPy
	269) 5'-W G G A C T W-3'	ImImPyPyHp- γ -PyImHpPyPy
	270) 5'-W G G A C A W-3'	ImImPyPyPy- γ -HpImHpPyPy
	271) 5'-W G G A C G W-3'	ImImPyPyIm- γ -PyImHpPyPy
35	272) 5'-W G G A C C W-3'	ImImPyPyPy- γ -ImImHpPyPy

TABLE 21: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	273) 5'-W G G G T T W-3'	ImImImHpHp- γ -PyPyPyPyPy
	274) 5'-W G G G T A W-3'	ImImImHpPy- γ -HpPyPyPyPy
	275) 5'-W G G G T G W-3'	ImImImHpIm- γ -PyPyPyPyPy
	276) 5'-W G G G T C W-3'	ImImImHpPy- γ -ImPyPyPyPy
	277) 5'-W G G G A T W-3'	ImImImPyHp- γ -PyHpPyPyPy
10	278) 5'-W G G G A A W-3'	ImImImPyPy- γ -HpHpPyPyPy
	279) 5'-W G G G A G W-3'	ImImImPyIm- γ -PyHpPyPyPy
	280) 5'-W G G G A C W-3'	ImImImPyPy- γ -ImHpPyPyPy
	281) 5'-W G G G G T W-3'	ImImImImHp- γ -PyPyPyPyPy
	282) 5'-W G G G G A W-3'	ImImImImPy- γ -HpPyPyPyPy
	283) 5'-W G G G C T W-3'	ImImImPyHp- γ -PyImPyPyPy
	284) 5'-W G G G C A W-3'	ImImImPyPy- γ -HpImPyPyPy
	285) 5'-W G G C T T W-3'	ImImPyHpHp- γ -PyPyImPyPy
	286) 5'-W G G C T A W-3'	ImImPyHpPy- γ -HpPyImPyPy
	287) 5'-W G G C T G W-3'	ImImPyHpIm- γ -PyPyImPyPy
	288) 5'-W G G C T C W-3'	ImImPyHpPy- γ -ImPyImPyPy
20	289) 5'-W G G C A T W-3'	ImImPyPyHp- γ -PyHpImPyPy
	290) 5'-W G G C A A W-3'	ImImPyPyPy- γ -HpHpImPyPy
	291) 5'-W G G C A G W-3'	ImImPyPyIm- γ -PyHpImPyPy
	292) 5'-W G G C A C W-3'	ImImPyPyPy- γ -ImHpImPyPy
	293) 5'-W G G C G T W-3'	ImImPyImHp- γ -PyPyImPyPy
25	294) 5'-W G G C G A W-3'	ImImPyImPy- γ -HpPyImPyPy
	295) 5'-W G G C C T W-3'	ImImPyPyHp- γ -PyImImPyPy
	296) 5'-W G G C C A W-3'	ImImPyPyPy- γ -HpImImPyPy
	G17) 5'-W G G G G G W-3'	ImImImImIm- γ -PyPyPyPyPy
	G18) 5'-W G G G G C W-3'	ImImImImPy- γ -ImPyPyPyPy
30	G19) 5'-W G G G C G W-3'	ImImImPyIm- γ -PyImPyPyPy
	G20) 5'-W G G G C C W-3'	ImImImPyPy- γ -ImImPyPyPy
	G21) 5'-W G G C G G W-3'	ImImPyImIm- γ -PyPyImPyPy
	G22) 5'-W G G C G C W-3'	ImImPyImPy- γ -ImPyImPyPy
	G23) 5'-W G G C C G W-3'	ImImPyPyIm- γ -PyImImPyPy
35	G24) 5'-W G G C C C W-3'	ImImPyPyPy- γ -ImImImPyPy

TABLE 22: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGTWNNW-3'

		DNA sequence	aromatic amino acid sequence
5	297)	5'-W G T T T T W-3'	ImHpHpHpHp- γ -PyPyPyPyPy
	298)	5'-W G T T T A W-3'	ImHpHpHpPy- γ -HpPyPyPyPy
	299)	5'-W G T T T G W-3'	ImHpHpHpIm- γ -PyPyPyPyPy
	300)	5'-W G T T T C W-3'	ImHpHpHpPy- γ -ImPyPyPyPy
	301)	5'-W G T T A T W-3'	ImHpHpPyHp- γ -PyHpPyPyPy
10	302)	5'-W G T T A A W-3'	ImHpHpPyPy- γ -HpHpPyPyPy
	303)	5'-W G T T A G W-3'	ImHpHpPyIm- γ -PyHpPyPyPy
	304)	5'-W G T T A C W-3'	ImHpHpPyPy- γ -ImHpPyPyPy
	305)	5'-W G T T G T W-3'	ImHpHpImHp- γ -PyPyPyPyPy
	306)	5'-W G T T G A W-3'	ImHpHpImPy- γ -HpPyPyPyPy
	307)	5'-W G T T G G W-3'	ImHpHpImIm- γ -PyPyPyPyPy
	308)	5'-W G T T G C W-3'	ImHpHpImPy- γ -ImPyPyPyPy
	309)	5'-W G T T C T W-3'	ImHpHpPyHp- γ -PyImPyPyPy
	310)	5'-W G T T C A W-3'	ImHpHpPyPy- γ -HpImPyPyPy
	311)	5'-W G T T C G W-3'	ImHpHpPyIm- γ -PyImPyPyPy
20	312)	5'-W G T T C C W-3'	ImHpHpPyPy- γ -ImImPyPyPy
	313)	5'-W G T A T T W-3'	ImHpPyHpHp- γ -PyPyHpPyPy
	314)	5'-W G T A T A W-3'	ImHpPyHpPy- γ -HpPyHpPyPy
	315)	5'-W G T A T G W-3'	ImHpPyHpIm- γ -PyPyHpPyPy
	316)	5'-W G T A T C W-3'	ImHpPyHpPy- γ -ImPyHpPyPy
	317)	5'-W G T A A T W-3'	ImHpPyPyHp- γ -PyHpHpPyPy
	318)	5'-W G T A A A W-3'	ImHpPyPyPy- γ -HpHpHpPyPy
	319)	5'-W G T A A G W-3'	ImHpPyPyIm- γ -PyHpHpPyPy
	320)	5'-W G T A A C W-3'	ImHpPyPyPy- γ -ImHpHpPyPy
	321)	5'-W G T A G T W-3'	ImHpPyImHp- γ -PyPyHpPyPy
30	322)	5'-W G T A G A W-3'	ImHpPyImPy- γ -HpPyHpPyPy
	323)	5'-W G T A G G W-3'	ImHpPyImIm- γ -PyPyHpPyPy
	324)	5'-W G T A G C W-3'	ImHpPyImPy- γ -ImPyHpPyPy
	325)	5'-W G T A C T W-3'	ImHpPyPyHp- γ -PyImHpPyPy
	326)	5'-W G T A C A W-3'	ImHpPyPyPy- γ -HpImHpPyPy
	327)	5'-W G T A C G W-3'	ImHpPyPyIm- γ -PyImHpPyPy
	328)	5'-W G T A C C W-3'	ImHpPyPyPy- γ -ImImHpPyPy
35			

TABLE 23: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	329) 5'-W G T G T T W-3'	ImHpImHpHp- γ -PyPyPyPyPy
5	330) 5'-W G T G T A W-3'	ImHpImHpPy- γ -HpPyPyPyPy
	331) 5'-W G T G T G W-3'	ImHpImHpIm- γ -PyPyPyPyPy
	332) 5'-W G T G T C W-3'	ImHpImHpPy- γ -ImPyPyPyPy
	333) 5'-W G T G A T W-3'	ImHpImPyHp- γ -PyHpPyPyPy
	334) 5'-W G T G A A W-3'	ImHpImPyPy- γ -HpHpPyPyPy
10	335) 5'-W G T G A G W-3'	ImHpImPyIm- γ -PyHpPyPyPy
	336) 5'-W G T G A C W-3'	ImHpImPyPy- γ -ImHpPyPyPy
	337) 5'-W G T G G T W-3'	ImHpImImHp- γ -PyPyPyPyPy
	338) 5'-W G T G G A W-3'	ImHpImImPy- γ -HpPyPyPyPy
	339) 5'-W G T G C T W-3'	ImHpImPyHp- γ -PyImPyPyPy
	340) 5'-W G T G C A W-3'	ImHpImPyPy- γ -HpImPyPyPy
	341) 5'-W G T G G G W-3'	ImHpImImIm- γ -PyPyPyPyPy
	342) 5'-W G T G G C W-3'	ImHpImImPy- γ -ImPyPyPyPy
	343) 5'-W G T G C G W-3'	ImHpImPyIm- γ -PyImPyPyPy
	344) 5'-W G T G C C W-3'	ImHpImPyPy- γ -ImImPyPyPy
	345) 5'-W G T C T T W-3'	ImHpPyHpHp- γ -PyPyImPyPy
	346) 5'-W G T C T A W-3'	ImHpPyHpPy- γ -HpPyImPyPy
	347) 5'-W G T C T G W-3'	ImHpPyHpIm- γ -PyPyImPyPy
	348) 5'-W G T C T C W-3'	ImHpPyHpPy- γ -ImPyImPyPy
	349) 5'-W G T C A T W-3'	ImHpPyPyHp- γ -PyHpImPyPy
25	350) 5'-W G T C A A W-3'	ImHpPyPyPy- γ -HpHpImPyPy
	351) 5'-W G T C A G W-3'	ImHpPyPyIm- γ -PyHpImPyPy
	352) 5'-W G T C A C W-3'	ImHpPyPyPy- γ -ImHpImPyPy
	353) 5'-W G T C G T W-3'	ImHpPyImHp- γ -PyPyImPyPy
	354) 5'-W G T C G A W-3'	ImHpPyImPy- γ -HpPyImPyPy
30	355) 5'-W G T C C T W-3'	ImHpPyPyHp- γ -PyImImPyPy
	356) 5'-W G T C C A W-3'	ImHpPyPyPy- γ -HpImImPyPy
	357) 5'-W G T C G G W-3'	ImHpPyImIm- γ -PyPyImPyPy
	358) 5'-W G T C G C W-3'	ImHpPyImPy- γ -ImPyImPyPy
	359) 5'-W G T C C G W-3'	ImHpPyPyIm- γ -PyImImPyPy
35	360) 5'-W G T C C C W-3'	ImHpPyPyPy- γ -ImImImPyPy

TABLE 24: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	361) 5'-W G A T T T W-3'	ImPyHpHpHp- γ -PyPyPyHpPy
5	362) 5'-W G A T T A W-3'	ImPyHpHpPy- γ -HpPyPyHpPy
	363) 5'-W G A T T G W-3'	ImPyHpHpIm- γ -PyPyPyHpPy
	364) 5'-W G A T T C W-3'	ImPyHpHpPy- γ -ImPyPyHpPy
	365) 5'-W G A T A T W-3'	ImPyHpPyHp- γ -PyHpPyHpPy
	366) 5'-W G A T A A W-3'	ImPyHpPyPy- γ -HpHpPyHpPy
10	367) 5'-W G A T A G W-3'	ImPyHpPyIm- γ -PyHpPyHpPy
	368) 5'-W G A T A C W-3'	ImPyHpPyPy- γ -ImHpPyHpPy
	369) 5'-W G A T G T W-3'	ImPyHpImHp- γ -PyPyPyHpPy
	370) 5'-W G A T G A W-3'	ImPyHpImPy- γ -HpPyPyHpPy
	371) 5'-W G A T G G W-3'	ImPyHpImIm- γ -PyPyPyHpPy
	372) 5'-W G A T G C W-3'	ImPyHpImPy- γ -ImPyPyHpPy
	373) 5'-W G A T C T W-3'	ImPyHpPyHp- γ -PyImPyHpPy
	374) 5'-W G A T C A W-3'	ImPyHpPyPy- γ -HpImPyHpPy
	375) 5'-W G A T C G W-3'	ImPyHpPyIm- γ -PyImPyHpPy
	376) 5'-W G A T C C W-3'	ImPyHpPyPy- γ -ImImPyHpPy
20	377) 5'-W G A A T T W-3'	ImPyPyHpHp- γ -PyPyHpHpPy
	378) 5'-W G A A T A W-3'	ImPyPyHpPy- γ -HpPyHpHpPy
	379) 5'-W G A A T G W-3'	ImPyPyHpIm- γ -PyPyHpHpPy
	380) 5'-W G A A T C W-3'	ImPyPyHpPy- γ -ImPyHpHpPy
	381) 5'-W G A A A T W-3'	ImPyPyPyHp- γ -PyHpHpHpPy
25	382) 5'-W G A A A A W-3'	ImPyPyPyPy- γ -HpHpHpHpPy
	383) 5'-W G A A A G W-3'	ImPyPyPyIm- γ -PyHpHpHpPy
	384) 5'-W G A A A C W-3'	ImPyPyPyPy- γ -ImHpHpHpPy
	385) 5'-W G A A G T W-3'	ImPyPyImHp- γ -PyPyHpHpPy
	386) 5'-W G A A G A W-3'	ImPyPyImPy- γ -HpPyHpHpPy
30	387) 5'-W G A A G G W-3'	ImPyPyImIm- γ -PyPyHpHpPy
	388) 5'-W G A A G C W-3'	ImPyPyImPy- γ -ImPyHpHpPy
	389) 5'-W G A A C T W-3'	ImPyPyPyHp- γ -PyImHpHpPy
	390) 5'-W G A A C A W-3'	ImPyPyPyPy- γ -HpImHpHpPy
	391) 5'-W G A A C G W-3'	ImPyPyPyIm- γ -PyImHpHpPy
35	392) 5'-W G A A C C W-3'	ImPyPyPyPy- γ -ImImHpHpPy

TABLE 25: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGASNNW-3'

	DNA sequence	aromatic amino acid sequence
5	393) 5'-W G A G T T W-3'	ImPyImHpHp-γ-PyPyPyHpPy
	394) 5'-W G A G T A W-3'	ImPyImHpPy-γ-HpPyPyHpPy
	395) 5'-W G A G T G W-3'	ImPyImHpIm-γ-PyPyPyHpPy
	396) 5'-W G A G T C W-3'	ImPyImHpPy-γ-ImPyPyHpPy
	397) 5'-W G A G A T W-3'	ImPyImPyHp-γ-PyHpPyHpPy
10	398) 5'-W G A G A A W-3'	ImPyImPyPy-γ-HpHpPyHpPy
	399) 5'-W G A G A G W-3'	ImPyImPyIm-γ-PyHpPyHpPy
	400) 5'-W G A G A C W-3'	ImPyImPyPy-γ-ImHpPyHpPy
	401) 5'-W G A G G T W-3'	ImPyImImHp-γ-PyPyPyHpPy
	402) 5'-W G A G G A W-3'	ImPyImImPy-γ-HpPyPyHpPy
	403) 5'-W G A G C T W-3'	ImPyImPyHp-γ-PyImPyHpPy
	404) 5'-W G A G C A W-3'	ImPyImPyPy-γ-HpImPyHpPy
	405) 5'-W G A G G G W-3'	ImPyImImIm-γ-PyPyPyHpPy
	406) 5'-W G A G G C W-3'	ImPyImImPy-γ-ImPyPyHpPy
	407) 5'-W G A G C G W-3'	ImPyImPyIm-γ-PyImPyHpPy
20	408) 5'-W G A G C C W-3'	ImPyImPyPy-γ-ImImPyHpPy
	409) 5'-W G A C T T W-3'	ImPyPyHpHp-γ-PyPyImHpPy
	410) 5'-W G A C T A W-3'	ImPyPyHpPy-γ-HpPyImHpPy
	411) 5'-W G A C T G W-3'	ImPyPyHpIm-γ-PyPyImHpPy
	412) 5'-W G A C T C W-3'	ImPyPyHpPy-γ-ImPyImHpPy
	413) 5'-W G A C A T W-3'	ImPyPyPyHp-γ-PyHpImHpPy
	414) 5'-W G A C A A W-3'	ImPyPyPyPy-γ-HpHpImHpPy
	415) 5'-W G A C A G W-3'	ImPyPyPyIm-γ-PyHpImHpPy
	416) 5'-W G A C A C W-3'	ImPyPyPyPy-γ-ImHpImHpPy
	417) 5'-W G A C G T W-3'	ImPyPyImHp-γ-PyPyImHpPy
30	418) 5'-W G A C G A W-3'	ImPyPyImPy-γ-HpPyImHpPy
	419) 5'-W G A C C T W-3'	ImPyPyPyHp-γ-PyImImHpPy
	420) 5'-W G A C C A W-3'	ImPyPyPyPy-γ-HpImImHpPy
	421) 5'-W G A C G G W-3'	ImPyPyImIm-γ-PyPyImHpPy
	422) 5'-W G A C G C W-3'	ImPyPyImPy-γ-ImPyImHpPy
	423) 5'-W G A C C G W-3'	ImPyPyPyIm-γ-PyImImHpPy
	424) 5'-W G A C C C W-3'	ImPyPyPyPy-γ-ImImImHpPy

TABLE 26: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	425) 5'-W G C T T T W-3'	ImPyHpHpHp-γ-PyPyPyImPy
5	426) 5'-W G C T T A W-3'	ImPyHpHpPy-γ-HpPyPyImPy
	427) 5'-W G C T T G W-3'	ImPyHpHpIm-γ-PyPyPyImPy
	428) 5'-W G C T T C W-3'	ImPyHpHpPy-γ-ImPyPyImPy
	429) 5'-W G C T A T W-3'	ImPyHpPyHp-γ-PyHpPyImPy
	430) 5'-W G C T A A W-3'	ImPyHpPyPy-γ-HpHpPyImPy
10	431) 5'-W G C T A G W-3'	ImPyHpPyIm-γ-PyHpPyImPy
	432) 5'-W G C T A C W-3'	ImPyHpPyPy-γ-ImHpPyImPy
	433) 5'-W G C T G T W-3'	ImPyHpImHp-γ-PyPyPyImPy
	434) 5'-W G C T G A W-3'	ImPyHpImPy-γ-HpPyPyImPy
	435) 5'-W G C T G G W-3'	ImPyHpImIm-γ-PyPyPyImPy
	436) 5'-W G C T G C W-3'	ImPyHpImPy-γ-ImPyPyImPy
	437) 5'-W G C T C T W-3'	ImPyHpPyHp-γ-PyImPyImPy
	438) 5'-W G C T C A W-3'	ImPyHpPyPy-γ-HpImPyImPy
	439) 5'-W G C T C G W-3'	ImPyHpPyIm-γ-PyImPyImPy
	440) 5'-W G C T C C W-3'	ImPyHpPyPy-γ-ImImPyImPy
	441) 5'-W G C A T T W-3'	ImPyPyHpHp-γ-PyPyHpImPy
	442) 5'-W G C A T A W-3'	ImPyPyHpPy-γ-HpPyHpImPy
	443) 5'-W G C A T G W-3'	ImPyPyHpIm-γ-PyPyHpImPy
	444) 5'-W G C A T C W-3'	ImPyPyHpPy-γ-ImPyHpImPy
	445) 5'-W G C A A T W-3'	ImPyPyPyHp-γ-PyHpHpImPy
25	446) 5'-W G C A A A W-3'	ImPyPyPyPy-γ-HpHpHpImPy
	447) 5'-W G C A A G W-3'	ImPyPyPyIm-γ-PyHpHpImPy
	448) 5'-W G C A A C W-3'	ImPyPyPyPy-γ-ImHpHpImPy
	449) 5'-W G C A G T W-3'	ImPyPyImHp-γ-PyPyHpImPy
	450) 5'-W G C A G A W-3'	ImPyPyImPy-γ-HpPyHpImPy
30	451) 5'-W G C A G G W-3'	ImPyPyImIm-γ-PyPyHpImPy
	452) 5'-W G C A G C W-3'	ImPyPyImPy-γ-ImPyHpImPy
	453) 5'-W G C A C T W-3'	ImPyPyPyHp-γ-PyImHpImPy
	454) 5'-W G C A C A W-3'	ImPyPyPyPy-γ-HpImHpImPy
	455) 5'-W G C A C G W-3'	ImPyPyPyIm-γ-PyImHpImPy
35	456) 5'-W G C A C C W-3'	ImPyPyPyPy-γ-ImImHpImPy

TABLE 27: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	457) 5'-W G C G T T W-3'	ImPyImHpHp-γ-PyPyPyImPy
	458) 5'-W G C G T A W-3'	ImPyImHpPy-γ-HpPyPyImPy
	459) 5'-W G C G T G W-3'	ImPyImHpIm-γ-PyPyPyImPy
	460) 5'-W G C G T C W-3'	ImPyImHpPy-γ-ImPyPyImPy
	461) 5'-W G C G A T W-3'	ImPyImPyHp-γ-PyHpPyImPy
10	462) 5'-W G C G A A W-3'	ImPyImPyPy-γ-HpHpPyImPy
	463) 5'-W G C G A G W-3'	ImPyImPyIm-γ-PyHpPyImPy
	464) 5'-W G C G A C W-3'	ImPyImPyPy-γ-ImHpPyImPy
	465) 5'-W G C G G T W-3'	ImPyImImHp-γ-PyPyPyImPy
	466) 5'-W G C G G A W-3'	ImPyImImPy-γ-HpPyPyImPy
	467) 5'-W G C G C T W-3'	ImPyImPyHp-γ-PyImPyImPy
	468) 5'-W G C G C A W-3'	ImPyImPyPy-γ-HpImPyImPy
	469) 5'-W G C C T T W-3'	ImPyPyHpHp-γ-PyPyImImPy
	470) 5'-W G C C T A W-3'	ImPyPyHpPy-γ-HpPyImImPy
	471) 5'-W G C C T G W-3'	ImPyPyHpIm-γ-PyPyImImPy
20	472) 5'-W G C C T C W-3'	ImPyPyHpPy-γ-ImPyImImPy
	473) 5'-W G C C A T W-3'	ImPyPyPyHp-γ-PyHpImImPy
	474) 5'-W G C C A A W-3'	ImPyPyPyPy-γ-HpHpImImPy
	475) 5'-W G C C A G W-3'	ImPyPyPyIm-γ-PyHpImImPy
	476) 5'-W G C C A C W-3'	ImPyPyPyPy-γ-ImHpImImPy
	477) 5'-W G C C G T W-3'	ImPyPyImHp-γ-PyPyImImPy
	478) 5'-W G C C G A W-3'	ImPyPyImPy-γ-HpPyImImPy
25	479) 5'-W G C C C T W-3'	ImPyPyPyHp-γ-PyImImImPy
	480) 5'-W G C C C A W-3'	ImPyPyPyPy-γ-HpImImImPy
	G25) 5'-W G C G G G W-3'	ImPyImImIm-γ-PyPyPyImPy
	G26) 5'-W G C G G C W-3'	ImPyImImPy-γ-ImPyPyImPy
	G27) 5'-W G C G C G W-3'	ImPyImPyIm-γ-PyImPyImPy
30	G28) 5'-W G C G C C W-3'	ImPyImPyPy-γ-ImImPyImPy
	G29) 5'-W G C C G G W-3'	ImPyPyImIm-γ-PyPyImImPy
	G30) 5'-W G C C G C W-3'	ImPyPyImPy-γ-ImPyImImPy
	G31) 5'-W G C C C G W-3'	ImPyPyPyIm-γ-PyImImImPy
	G32) 5'-W G C C C C W-3'	ImPyPyPyPy-γ-ImImImImPy

TABLE 28: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	481) 5'-W C G T T T W-3'	PyImHpHpHp-γ-PyPyPyPyIm
	482) 5'-W C G T T A W-3'	PyImHpHpPy-γ-HpPyPyPyIm
	483) 5'-W C G T T G W-3'	PyImHpHpIm-γ-PyPyPyPyIm
	484) 5'-W C G T T C W-3'	PyImHpHpPy-γ-ImPyPyPyIm
	485) 5'-W C G T A T W-3'	PyImHpPyHp-γ-PyHpPyPyIm
10	486) 5'-W C G T A A W-3'	PyImHpPyPy-γ-HpHpPyPyIm
	487) 5'-W C G T A G W-3'	PyImHpPyIm-γ-PyHpPyPyIm
	488) 5'-W C G T A C W-3'	PyImHpPyPy-γ-ImHpPyPyIm
	489) 5'-W C G T G T W-3'	PyImHpImHp-γ-PyPyPyPyIm
	490) 5'-W C G T G A W-3'	PyImHpImPy-γ-HpPyPyPyIm
	491) 5'-W C G T G G W-3'	PyImHpImIm-γ-PyPyPyPyIm
	492) 5'-W C G T G C W-3'	PyImHpImPy-γ-ImPyPyPyIm
	493) 5'-W C G T C T W-3'	PyImHpPyHp-γ-PyImPyPyIm
	494) 5'-W C G T C A W-3'	PyImHpPyPy-γ-HpImPyPyIm
	495) 5'-W C G T C G W-3'	PyImHpPyIm-γ-PyImPyPyIm
20	496) 5'-W C G T C C W-3'	PyImHpPyPy-γ-ImImPyPyIm
	497) 5'-W C G A T T W-3'	PyImPyHpHp-γ-PyPyHpPyIm
	498) 5'-W C G A T A W-3'	PyImPyHpPy-γ-HpPyHpPyIm
	499) 5'-W C G A T G W-3'	PyImPyHpIm-γ-PyPyHpPyIm
	500) 5'-W C G A T C W-3'	PyImPyHpPy-γ-ImPyHpPyIm
	501) 5'-W C G A A T W-3'	PyImPyPyHp-γ-PyHpHpPyIm
	502) 5'-W C G A A A W-3'	PyImPyPyPy-γ-HpHpHpPyIm
	503) 5'-W C G A A G W-3'	PyImPyPyIm-γ-PyHpHpPyIm
	504) 5'-W C G A A C W-3'	PyImPyPyPy-γ-ImHpHpPyIm
	505) 5'-W C G A G T W-3'	PyImPyImHp-γ-PyPyHpPyIm
25	506) 5'-W C G A G A W-3'	PyImPyImPy-γ-HpPyHpPyIm
	507) 5'-W C G A G G W-3'	PyImPyImIm-γ-PyPyHpPyIm
	508) 5'-W C G A G C W-3'	PyImPyImPy-γ-ImPyHpPyIm
	509) 5'-W C G A C T W-3'	PyImPyPyHp-γ-PyImHpPyIm
	510) 5'-W C G A C A W-3'	PyImPyPyPy-γ-HpImHpPyIm
30	511) 5'-W C G A C G W-3'	PyImPyPyIm-γ-PyImHpPyIm
	512) 5'-W C G A C C W-3'	PyImPyPyPy-γ-ImImHpPyIm

TABLE 29: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	513) 5'-W C G G T T W-3'	PyImImHpHp-γ-PyPyPyPyIm
	514) 5'-W C G G T A W-3'	PyImImHpPy-γ-HpPyPyPyIm
	515) 5'-W C G G T G W-3'	PyImImHpIm-γ-PyPyPyPyIm
	516) 5'-W C G G T C W-3'	PyImImHpPy-γ-ImPyPyPyIm
	517) 5'-W C G G A T W-3'	PyImImPyHp-γ-PyHpPyPyIm
10	518) 5'-W C G G A A W-3'	PyImImPyPy-γ-HpHpPyPyIm
	519) 5'-W C G G A G W-3'	PyImImPyIm-γ-PyHpPyPyIm
	520) 5'-W C G G A C W-3'	PyImImPyPy-γ-ImHpPyPyIm
	521) 5'-W C G G G T W-3'	PyImImImHp-γ-PyPyPyPyIm
	522) 5'-W C G G G A W-3'	PyImImImPy-γ-HpPyPyPyIm
	523) 5'-W C G G C T W-3'	PyImImPyHp-γ-PyImPyPyIm
	524) 5'-W C G G C A W-3'	PyImImPyPy-γ-HpImPyPyIm
	525) 5'-W C G C T T W-3'	PyImPyHpHp-γ-PyPyImPyIm
	526) 5'-W C G C T A W-3'	PyImPyHpPy-γ-HpPyImPyIm
	527) 5'-W C G C T G W-3'	PyImPyHpIm-γ-PyPyImPyIm
20	528) 5'-W C G C T C W-3'	PyImPyHpPy-γ-ImPyImPyIm
	529) 5'-W C G C A T W-3'	PyImPyPyHp-γ-PyHpImPyIm
	530) 5'-W C G C A A W-3'	PyImPyPyPy-γ-HpHpImPyIm
	531) 5'-W C G C A G W-3'	PyImPyPyIm-γ-PyHpImPyIm
	532) 5'-W C G C A C W-3'	PyImPyPyPy-γ-ImHpImPyIm
	533) 5'-W C G C G T W-3'	PyImPyImHp-γ-PyPyImPyIm
	534) 5'-W C G C G A W-3'	PyImPyImPy-γ-HpPyImPyIm
	535) 5'-W C G C C T W-3'	PyImPyPyHp-γ-PyImImPyIm
	536) 5'-W C G C C A W-3'	PyImPyPyPy-γ-HpImImPyIm
	G33) 5'-W C G G G G W-3'	PyImImImIm-γ-PyPyPyPyIm
30	G34) 5'-W C G G G C W-3'	PyImImImPy-γ-ImPyPyPyIm
	G35) 5'-W C G G C G W-3'	PyImImPyIm-γ-PyImPyPyIm
	G36) 5'-W C G G C C W-3'	PyImImPyPy-γ-ImImPyPyIm
	G37) 5'-W C G C G G W-3'	PyImPyImIm-γ-PyPyImPyIm
	G38) 5'-W C G C G C W-3'	PyImPyImPy-γ-ImPyImPyIm
35	G39) 5'-W C G C C G W-3'	PyImPyPyIm-γ-PyImImPyIm
	G40) 5'-W C G C C C W-3'	PyImPyPyPy-γ-ImImImPyIm

TABLE 30: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	537) 5'-W C T T T T W-3'	PyHpHpHpHp-γ-PyPyPyPyIm
5	538) 5'-W C T T T A W-3'	PyHpHpHpPy-γ-HpPyPyPyIm
	539) 5'-W C T T T G W-3'	PyHpHpHpIm-γ-PyPyPyPyIm
	540) 5'-W C T T T C W-3'	PyHpHpHpPy-γ-ImPyPyPyIm
	541) 5'-W C T T A T W-3'	PyHpHpPyHp-γ-PyHpPyPyIm
	542) 5'-W C T T A A W-3'	PyHpHpPyPy-γ-HpHpPyPyIm
10	543) 5'-W C T T A G W-3'	PyHpHpPyIm-γ-PyHpPyPyIm
	544) 5'-W C T T A C W-3'	PyHpHpPyPy-γ-ImHpPyPyIm
	545) 5'-W C T T G T W-3'	PyHpHpImHp-γ-PyPyPyPyIm
	546) 5'-W C T T G A W-3'	PyHpHpImPy-γ-HpPyPyPyIm
	547) 5'-W C T T G G W-3'	PyHpHpImIm-γ-PyPyPyPyIm
	548) 5'-W C T T G C W-3'	PyHpHpImPy-γ-ImPyPyPyIm
	549) 5'-W C T T C T W-3'	PyHpHpPyHp-γ-PyImPyPyIm
	550) 5'-W C T T C A W-3'	PyHpHpPyPy-γ-HpImPyPyIm
	551) 5'-W C T T C G W-3'	PyHpHpPyIm-γ-PyImPyPyIm
	552) 5'-W C T T C C W-3'	PyHpHpPyPy-γ-ImImPyPyIm
20	553) 5'-W C T A T T W-3'	PyHpPyHpHp-γ-PyPyHpPyIm
	554) 5'-W C T A T A W-3'	PyHpPyHpPy-γ-HpPyHpPyIm
	555) 5'-W C T A T G W-3'	PyHpPyHpIm-γ-PyPyHpPyIm
	556) 5'-W C T A T C W-3'	PyHpPyHpPy-γ-ImPyHpPyIm
	557) 5'-W C T A A T W-3'	PyHpPyPyHp-γ-PyHpHpPyIm
25	558) 5'-W C T A A A W-3'	PyHpPyPyPy-γ-HpHpHpPyIm
	559) 5'-W C T A A G W-3'	PyHpPyPyIm-γ-PyHpHpPyIm
	560) 5'-W C T A A C W-3'	PyHpPyPyPy-γ-ImHpHpPyIm
	561) 5'-W C T A G T W-3'	PyHpPyImHp-γ-PyPyHpPyIm
	562) 5'-W C T A G A W-3'	PyHpPyImPy-γ-HpPyHpPyIm
30	563) 5'-W C T A G G W-3'	PyHpPyImIm-γ-PyPyHpPyIm
	564) 5'-W C T A G C W-3'	PyHpPyImPy-γ-ImPyHpPyIm
	565) 5'-W C T A C T W-3'	PyHpPyPyHp-γ-PyImHpPyIm
	566) 5'-W C T A C A W-3'	PyHpPyPyPy-γ-HpImHpPyIm
	567) 5'-W C T A C G W-3'	PyHpPyPyIm-γ-PyImHpPyIm
35	568) 5'-W C T A C C W-3'	PyHpPyPyPy-γ-ImImHpPyIm

TABLE 31: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	569) 5'-W C T G T T W-3'	PyHpImHpHp- γ -PyPyPyPyIm
5	570) 5'-W C T G T A W-3'	PyHpImHpPy- γ -HpPyPyPyIm
	571) 5'-W C T G T G W-3'	PyHpImHpIm- γ -PyPyPyPyIm
	572) 5'-W C T G T C W-3'	PyHpImHpPy- γ -ImPyPyPyIm
	573) 5'-W C T G A T W-3'	PyHpImPyHp- γ -PyHpPyPyIm
	574) 5'-W C T G A A W-3'	PyHpImPyPy- γ -HpHpPyPyIm
10	575) 5'-W C T G A G W-3'	PyHpImPyIm- γ -PyHpPyPyIm
	576) 5'-W C T G A C W-3'	PyHpImPyPy- γ -ImHpPyPyIm
	577) 5'-W C T G G T W-3'	PyHpImImHp- γ -PyPyPyPyIm
	578) 5'-W C T G G A W-3'	PyHpImImPy- γ -HpPyPyPyIm
	579) 5'-W C T G C T W-3'	PyHpImPyHp- γ -PyImPyPyIm
	580) 5'-W C T G C A W-3'	PyHpImPyPy- γ -HpImPyPyIm
	581) 5'-W C T G G G W-3'	PyHpImImIm- γ -PyPyPyPyIm
	582) 5'-W C T G G C W-3'	PyHpImImPy- γ -ImPyPyPyIm
	583) 5'-W C T G C G W-3'	PyHpImPyIm- γ -PyImPyPyIm
	584) 5'-W C T G C C W-3'	PyHpImPyPy- γ -ImImPyPyIm
20	585) 5'-W C T C T T W-3'	PyHpPyHpHp- γ -PyPyImPyIm
	586) 5'-W C T C T A W-3'	PyHpPyHpPy- γ -HpPyImPyIm
	587) 5'-W C T C T G W-3'	PyHpPyHpIm- γ -PyPyImPyIm
	588) 5'-W C T C T C W-3'	PyHpPyHpPy- γ -ImPyImPyIm
	589) 5'-W C T C A T W-3'	PyHpPyPyHp- γ -PyHpImPyIm
25	590) 5'-W C T C A A W-3'	PyHpPyPyPy- γ -HpHpImPyIm
	591) 5'-W C T C A G W-3'	PyHpPyPyIm- γ -PyHpImPyIm
	592) 5'-W C T C A C W-3'	PyHpPyPyPy- γ -ImHpImPyIm
	593) 5'-W C T C G T W-3'	PyHpPyImHp- γ -PyPyImPyIm
	594) 5'-W C T C G A W-3'	PyHpPyImPy- γ -HpPyImPyIm
30	595) 5'-W C T C C T W-3'	PyHpPyPyHp- γ -PyImImPyIm
	596) 5'-W C T C C A W-3'	PyHpPyPyPy- γ -HpImImPyIm
	597) 5'-W C T C G G W-3'	PyHpPyImIm- γ -PyPyImPyIm
	598) 5'-W C T C G C W-3'	PyHpPyImPy- γ -ImPyImPyIm
	599) 5'-W C T C C G W-3'	PyHpPyPyIm- γ -PyImImPyIm
35	600) 5'-W C T C C C W-3'	PyHpPyPyPy- γ -ImImImPyIm

TABLE 32: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCAWNNW-3'

		DNA sequence	aromatic amino acid sequence
5	601)	5'-W C A T T T W-3'	PyPyHpHpHp-γ-PyPyPyHpIm
	602)	5'-W C A T T A W-3'	PyPyHpHpPy-γ-HpPyPyHpIm
	603)	5'-W C A T T G W-3'	PyPyHpHpIm-γ-PyPyPyHpIm
	604)	5'-W C A T T C W-3'	PyPyHpHpPy-γ-ImPyPyHpIm
	605)	5'-W C A T A T W-3'	PyPyHpPyHp-γ-PyHpPyHpIm
10	606)	5'-W C A T A A W-3'	PyPyHpPyPy-γ-HpHpPyHpIm
	607)	5'-W C A T A G W-3'	PyPyHpPyIm-γ-PyHpPyHpIm
	608)	5'-W C A T A C W-3'	PyPyHpPyPy-γ-ImHpPyHpIm
	609)	5'-W C A T G T W-3'	PyPyHpImHp-γ-PyPyPyHpIm
	610)	5'-W C A T G A W-3'	PyPyHpImPy-γ-HpPyPyHpIm
20	611)	5'-W C A T G G W-3'	PyPyHpImIm-γ-PyPyPyHpIm
	612)	5'-W C A T G C W-3'	PyPyHpImPy-γ-ImPyPyHpIm
	613)	5'-W C A T C T W-3'	PyPyHpPyHp-γ-PyImPyHpIm
	614)	5'-W C A T C A W-3'	PyPyHpPyPy-γ-HpImPyHpIm
	615)	5'-W C A T C G W-3'	PyPyHpPyIm-γ-PyImPyHpIm
25	616)	5'-W C A T C C W-3'	PyPyHpPyPy-γ-ImImPyHpIm
	617)	5'-W C A A T T W-3'	PyPyPyHpHp-γ-PyPyHpHpIm
	618)	5'-W C A A T A W-3'	PyPyPyHpPy-γ-HpPyHpHpIm
	619)	5'-W C A A T G W-3'	PyPyPyHpIm-γ-PyPyHpHpIm
	620)	5'-W C A A T C W-3'	PyPyPyHpPy-γ-ImPyHpHpIm
30	621)	5'-W C A A A T W-3'	PyPyPyPyHp-γ-PyHpHpHpIm
	622)	5'-W C A A A A W-3'	PyPyPyPyPy-γ-HpHpHpHpIm
	623)	5'-W C A A A G W-3'	PyPyPyPyIm-γ-PyHpHpHpIm
	624)	5'-W C A A A C W-3'	PyPyPyPyPy-γ-ImHpHpHpIm
	625)	5'-W C A A G T W-3'	PyPyPyImHp-γ-PyPyHpHpIm
35	626)	5'-W C A A G A W-3'	PyPyPyImPy-γ-HpPyHpHpIm
	627)	5'-W C A A G G W-3'	PyPyPyImIm-γ-PyPyHpHpIm
	628)	5'-W C A A G C W-3'	PyPyPyImPy-γ-ImPyHpHpIm
	629)	5'-W C A A C T W-3'	PyPyPyPyHp-γ-PyImHpHpIm
	630)	5'-W C A A C A W-3'	PyPyPyPyPy-γ-HpImHpHpIm
	631)	5'-W C A A C G W-3'	PyPyPyPyIm-γ-PyImHpHpIm
	632)	5'-W C A A C C W-3'	PyPyPyPyPy-γ-ImImHpHpIm

TABLE 33: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCASNNW-3'

	DNA sequence	aromatic amino acid sequence
	633) 5'-W C A G T T W-3'	PyPyImHpHp-γ-PyPyPyHpIm
5	634) 5'-W C A G T A W-3'	PyPyImHpPy-γ-HpPyPyHpIm
	635) 5'-W C A G T G W-3'	PyPyImHpIm-γ-PyPyPyHpIm
	636) 5'-W C A G T C W-3'	PyPyImHpPy-γ-ImPyPyHpIm
	637) 5'-W C A G A T W-3'	PyPyImPyHp-γ-PyHpPyHpIm
	638) 5'-W C A G A A W-3'	PyPyImPyPy-γ-HpHpPyHpIm
10	639) 5'-W C A G A G W-3'	PyPyImPyIm-γ-PyHpPyHpIm
	640) 5'-W C A G A C W-3'	PyPyImPyPy-γ-ImHpPyHpIm
	641) 5'-W C A G G T W-3'	PyPyImImHp-γ-PyPyPyHpIm
	642) 5'-W C A G G A W-3'	PyPyImImPy-γ-HpPyPyHpIm
	643) 5'-W C A G C T W-3'	PyPyImPyHp-γ-PyImPyHpIm
	644) 5'-W C A G C A W-3'	PyPyImPyPy-γ-HpImPyHpIm
	645) 5'-W C A G G G W-3'	PyPyImImIm-γ-PyPyPyHpIm
	646) 5'-W C A G G C W-3'	PyPyImImPy-γ-ImPyPyHpIm
	647) 5'-W C A G C G W-3'	PyPyImPyIm-γ-PyImPyHpIm
	648) 5'-W C A G C C W-3'	PyPyImPyPy-γ-ImImPyHpIm
	649) 5'-W C A C T T W-3'	PyPyPyHpHp-γ-PyPyImHpIm
	650) 5'-W C A C T A W-3'	PyPyPyHpPy-γ-HpPyImHpIm
	651) 5'-W C A C T G W-3'	PyPyPyHpIm-γ-PyPyImHpIm
	652) 5'-W C A C T C W-3'	PyPyPyHpPy-γ-ImPyImHpIm
	653) 5'-W C A C A T W-3'	PyPyPyPyHp-γ-PyHpImHpIm
25	654) 5'-W C A C A A W-3'	PyPyPyPyPy-γ-HpHpImHpIm
	655) 5'-W C A C A G W-3'	PyPyPyPyIm-γ-PyHpImHpIm
	656) 5'-W C A C A C W-3'	PyPyPyPyPy-γ-ImHpImHpIm
	657) 5'-W C A C G T W-3'	PyPyPyImHp-γ-PyPyImHpIm
	658) 5'-W C A C G A W-3'	PyPyPyImPy-γ-HpPyImHpIm
30	659) 5'-W C A C C T W-3'	PyPyPyPyHp-γ-PyImImHpIm
	660) 5'-W C A C C A W-3'	PyPyPyPyPy-γ-HpImImHpIm
	661) 5'-W C A C G G W-3'	PyPyPyImIm-γ-PyPyImHpIm
	662) 5'-W C A C G C W-3'	PyPyPyImPy-γ-ImPyImHpIm
	663) 5'-W C A C C G W-3'	PyPyPyPyIm-γ-PyImImHpIm
35	664) 5'-W C A C C C W-3'	PyPyPyPyPy-γ-ImImImHpIm

TABLE 34: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	665) 5'-W C C T T T W-3'	PyPyHpHpHp-γ-PyPyPyImIm
5	666) 5'-W C C T T A W-3'	PyPyHpHpPy-γ-HpPyPyImIm
	667) 5'-W C C T T G W-3'	PyPyHpHpIm-γ-PyPyPyImIm
	668) 5'-W C C T T C W-3'	PyPyHpHpPy-γ-ImPyPyImIm
	669) 5'-W C C T A T W-3'	PyPyHpPyHp-γ-PyHpPyImIm
	670) 5'-W C C T A A W-3'	PyPyHpPyPy-γ-HpHpPyImIm
10	671) 5'-W C C T A G W-3'	PyPyHpPyIm-γ-PyHpPyImIm
	672) 5'-W C C T A C W-3'	PyPyHpPyPy-γ-ImHpPyImIm
	673) 5'-W C C T G T W-3'	PyPyHpImHp-γ-PyPyPyImIm
	674) 5'-W C C T G A W-3'	PyPyHpImPy-γ-HpPyPyImIm
	675) 5'-W C C T G G W-3'	PyPyHpImIm-γ-PyPyPyImIm
15	676) 5'-W C C T G C W-3'	PyPyHpImPy-γ-ImPyPyImIm
	677) 5'-W C C T C T W-3'	PyPyHpPyHp-γ-PyImPyImIm
	678) 5'-W C C T C A W-3'	PyPyHpPyPy-γ-HpImPyImIm
	679) 5'-W C C T C G W-3'	PyPyHpPyIm-γ-PyImPyImIm
	680) 5'-W C C T C C W-3'	PyPyHpPyPy-γ-ImImPyImIm
20	681) 5'-W C C A T T W-3'	PyPyPyHpHp-γ-PyPyHpImIm
	682) 5'-W C C A T A W-3'	PyPyPyHpPy-γ-HpPyHpImIm
	683) 5'-W C C A T G W-3'	PyPyPyHpIm-γ-PyPyHpImIm
	684) 5'-W C C A T C W-3'	PyPyPyHpPy-γ-ImPyHpImIm
	685) 5'-W C C A A T W-3'	PyPyPyPyHp-γ-PyHpHpImIm
25	686) 5'-W C C A A A W-3'	PyPyPyPyPy-γ-HpHpHpImIm
	687) 5'-W C C A A G W-3'	PyPyPyPyIm-γ-PyHpHpImIm
	688) 5'-W C C A A C W-3'	PyPyPyPyPy-γ-ImHpHpImIm
	689) 5'-W C C A G T W-3'	PyPyPyImHp-γ-PyPyHpImIm
	690) 5'-W C C A G A W-3'	PyPyPyImPy-γ-HpPyHpImIm
30	691) 5'-W C C A G G W-3'	PyPyPyImIm-γ-PyPyHpImIm
	692) 5'-W C C A G C W-3'	PyPyPyImPy-γ-ImPyHpImIm
	693) 5'-W C C A C T W-3'	PyPyPyPyHp-γ-PyImHpImIm
	694) 5'-W C C A C A W-3'	PyPyPyPyPy-γ-HpImHpImIm
	695) 5'-W C C A C G W-3'	PyPyPyPyIm-γ-PyImHpImIm
35	696) 5'-W C C A C C W-3'	PyPyPyPyPy-γ-ImImHpImIm

TABLE 35: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	697) 5'-W C C G T T W-3'	PyPyImHpHp- γ -PyPyPyImIm
5	698) 5'-W C C G T A W-3'	PyPyImHpPy- γ -HpPyPyImIm
	699) 5'-W C C G T G W-3'	PyPyImHpIm- γ -PyPyPyImIm
	700) 5'-W C C G T C W-3'	PyPyImHpPy- γ -ImPyPyImIm
	701) 5'-W C C G A T W-3'	PyPyImPyHp- γ -PyHpPyImIm
	702) 5'-W C C G A A W-3'	PyPyImPyPy- γ -HpHpPyImIm
10	703) 5'-W C C G A G W-3'	PyPyImPyIm- γ -PyHpPyImIm
	704) 5'-W C C G A C W-3'	PyPyImPyPy- γ -ImHpPyImIm
	705) 5'-W C C G G T W-3'	PyPyImImHp- γ -PyPyPyImIm
	706) 5'-W C C G G A W-3'	PyPyImImPy- γ -HpPyPyImIm
	707) 5'-W C C G C T W-3'	PyPyImPyHp- γ -PyImPyImIm
	708) 5'-W C C G C A W-3'	PyPyImPyPy- γ -HpImPyImIm
	709) 5'-W C C C T T W-3'	PyPyPyHpHp- γ -PyPyImImIm
	710) 5'-W C C C T A W-3'	PyPyPyHpPy- γ -HpPyImImIm
	711) 5'-W C C C T G W-3'	PyPyPyHpIm- γ -PyPyImImIm
	712) 5'-W C C C T C W-3'	PyPyPyHpPy- γ -ImPyImImIm
20	713) 5'-W C C C A T W-3'	PyPyPyPyHp- γ -PyHpImImIm
	714) 5'-W C C C A A W-3'	PyPyPyPyPy- γ -HpHpImImIm
	715) 5'-W C C C A G W-3'	PyPyPyPyIm- γ -PyHpImImIm
	716) 5'-W C C C A C W-3'	PyPyPyPyPy- γ -ImHpImImIm
	717) 5'-W C C C G T W-3'	PyPyPyImHp- γ -PyPyImImIm
25	718) 5'-W C C C G A W-3'	PyPyPyImPy- γ -HpPyImImIm
	719) 5'-W C C C C T W-3'	PyPyPyPyHp- γ -PyImImImIm
	720) 5'-W C C C C A W-3'	PyPyPyPyPy- γ -HpImImImIm
	G41) 5'-W C C G G G W-3'	PyPyImImIm- γ -PyPyPyImIm
	G42) 5'-W C C G G C W-3'	PyPyImImPy- γ -ImPyPyImIm
30	G43) 5'-W C C G C G W-3'	PyPyImPyIm- γ -PyImPyImIm
	G44) 5'-W C C G C C W-3'	PyPyImPyPy- γ -ImImPyImIm
	G45) 5'-W C C C G G W-3'	PyPyPyImIm- γ -PyPyImImIm
	G46) 5'-W C C C G C W-3'	PyPyPyImPy- γ -ImPyImImIm
	G47) 5'-W C C C C G W-3'	PyPyPyPyIm- γ -PyImImImIm
35	G48) 5'-W C C C C C W-3'	PyPyPyPyPy- γ -ImImImImIm

TABLE 36: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	721) 5'-W A G T T T W-3'	PyImHpHpHp- γ -PyPyPyPyHp
5	722) 5'-W A G T T A W-3'	PyImHpHpPy- γ -HpPyPyPyHp
	723) 5'-W A G T T G W-3'	PyImHpHpIm- γ -PyPyPyPyHp
	724) 5'-W A G T T C W-3'	PyImHpHpPy- γ -ImPyPyPyHp
	725) 5'-W A G T A T W-3'	PyImHpPyHp- γ -PyHpPyPyHp
	726) 5'-W A G T A A W-3'	PyImHpPyPy- γ -HpHpPyPyHp
10	727) 5'-W A G T A G W-3'	PyImHpPyIm- γ -PyHpPyPyHp
	728) 5'-W A G T A C W-3'	PyImHpPyPy- γ -ImHpPyPyHp
	729) 5'-W A G T G T W-3'	PyImHpImHp- γ -PyPyPyPyHp
	730) 5'-W A G T G A W-3'	PyImHpImPy- γ -HpPyPyPyHp
	731) 5'-W A G T G G W-3'	PyImHpImIm- γ -PyPyPyPyHp
	732) 5'-W A G T G C W-3'	PyImHpImPy- γ -ImPyPyPyHp
	733) 5'-W A G T C T W-3'	PyImHpPyHp- γ -PyImPyPyHp
	734) 5'-W A G T C A W-3'	PyImHpPyPy- γ -HpImPyPyHp
	735) 5'-W A G T C G W-3'	PyImHpPyIm- γ -PyImPyPyHp
	736) 5'-W A G T C C W-3'	PyImHpPyPy- γ -ImImPyPyHp
20	737) 5'-W A G A T T W-3'	PyImPyHpHp- γ -PyPyHpPyHp
	738) 5'-W A G A T A W-3'	PyImPyHpPy- γ -HpPyHpPyHp
	739) 5'-W A G A T G W-3'	PyImPyHpIm- γ -PyPyHpPyHp
	740) 5'-W A G A T C W-3'	PyImPyHpPy- γ -ImPyHpPyHp
	741) 5'-W A G A A T W-3'	PyImPyPyHp- γ -PyHpHpPyHp
25	742) 5'-W A G A A A W-3'	PyImPyPyPy- γ -HpHpHpPyHp
	743) 5'-W A G A A G W-3'	PyImPyPyIm- γ -PyHpHpPyHp
	744) 5'-W A G A A C W-3'	PyImPyPyPy- γ -ImHpHpPyHp
	745) 5'-W A G A G T W-3'	PyImPyImHp- γ -PyPyHpPyHp
	746) 5'-W A G A G A W-3'	PyImPyImPy- γ -HpPyHpPyHp
30	747) 5'-W A G A G G W-3'	PyImPyImIm- γ -PyPyHpPyHp
	748) 5'-W A G A G C W-3'	PyImPyImPy- γ -ImPyHpPyHp
	749) 5'-W A G A C T W-3'	PyImPyPyHp- γ -PyImHpPyHp
	750) 5'-W A G A C A W-3'	PyImPyPyPy- γ -HpImHpPyHp
	751) 5'-W A G A C G W-3'	PyImPyPyIm- γ -PyImHpPyHp
35	752) 5'-W A G A C C W-3'	PyImPyPyPy- γ -ImImHpPyHp

TABLE 37: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	753) 5'-W A G G T T W-3'	PyImImHpHp- γ -PyPyPyPyHp
5	754) 5'-W A G G T A W-3'	PyImImHpPy- γ -HpPyPyPyHp
	755) 5'-W A G G T G W-3'	PyImImHpIm- γ -PyPyPyPyHp
	756) 5'-W A G G T C W-3'	PyImImHpPy- γ -ImPyPyPyHp
	757) 5'-W A G G A T W-3'	PyImImPyHp- γ -PyHpPyPyHp
	758) 5'-W A G G A A W-3'	PyImImPyPy- γ -HpHpPyPyHp
10	759) 5'-W A G G A G W-3'	PyImImPyIm- γ -PyHpPyPyHp
	760) 5'-W A G G A C W-3'	PyImImPyPy- γ -ImHpPyPyHp
	761) 5'-W A G G G T W-3'	PyImImImHp- γ -PyPyPyPyHp
	762) 5'-W A G G G A W-3'	PyImImImPy- γ -HpPyPyPyHp
	763) 5'-W A G G C T W-3'	PyImImPyHp- γ -PyImPyPyHp
	764) 5'-W A G G C A W-3'	PyImImPyPy- γ -HpImPyPyHp
	765) 5'-W A G C T T W-3'	PyImPyHpHp- γ -PyPyImPyHp
	766) 5'-W A G C T A W-3'	PyImPyHpPy- γ -HpPyImPyHp
	767) 5'-W A G C T G W-3'	PyImPyHpIm- γ -PyPyImPyHp
	768) 5'-W A G C T C W-3'	PyImPyHpPy- γ -ImPyImPyHp
20	769) 5'-W A G C A T W-3'	PyImPyPyHp- γ -PyHpImPyHp
	770) 5'-W A G C A A W-3'	PyImPyPyPy- γ -HpHpImPyHp
	771) 5'-W A G C A G W-3'	PyImPyPyIm- γ -PyHpImPyHp
	772) 5'-W A G C A C W-3'	PyImPyPyPy- γ -ImHpImPyHp
	773) 5'-W A G C G T W-3'	PyImPyImHp- γ -PyPyImPyHp
25	774) 5'-W A G C G A W-3'	PyImPyImPy- γ -HpPyImPyHp
	775) 5'-W A G C C T W-3'	PyImPyPyHp- γ -PyImImPyHp
	776) 5'-W A G C C A W-3'	PyImPyPyPy- γ -HpImImPyHp
	777) 5'-W A G G G G W-3'	PyImImImIm- γ -PyPyPyPyHp
	778) 5'-W A G G G C W-3'	PyImImImPy- γ -ImPyPyPyHp
30	779) 5'-W A G G C G W-3'	PyImImPyIm- γ -PyImPyPyHp
	780) 5'-W A G G C C W-3'	PyImImPyPy- γ -ImImPyPyHp
	781) 5'-W A G C G G W-3'	PyImPyImIm- γ -PyPyImPyHp
	782) 5'-W A G C G C W-3'	PyImPyImPy- γ -ImPyImPyHp
	783) 5'-W A G C C G W-3'	PyImPyPyIm- γ -PyImImPyHp
35	784) 5'-W A G C C C W-3'	PyImPyPyPy- γ -ImImImPyHp

TABLE 38: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WATWNNW-3'

		DNA sequence	aromatic amino acid sequence
	785)	5'-W A T T T T W-3'	PyHpHpHpHp-γ-PyPyPyPyHp
5	786)	5'-W A T T T A W-3'	PyHpHpHpPy-γ-HpPyPyPyHp
	787)	5'-W A T T T G W-3'	PyHpHpHpIm-γ-PyPyPyPyHp
	788)	5'-W A T T T C W-3'	PyHpHpHpPy-γ-ImPyPyPyHp
	789)	5'-W A T T A T W-3'	PyHpHpPyHp-γ-PyHpPyPyHp
	790)	5'-W A T T A A W-3'	PyHpHpPyPy-γ-HpHpPyPyHp
10	791)	5'-W A T T A G W-3'	PyHpHpPyIm-γ-PyHpPyPyHp
	792)	5'-W A T T A C W-3'	PyHpHpPyPy-γ-ImHpPyPyHp
	793)	5'-W A T T G T W-3'	PyHpHpImHp-γ-PyPyPyPyHp
	794)	5'-W A T T G A W-3'	PyHpHpImPy-γ-HpPyPyPyHp
	795)	5'-W A T T G G W-3'	PyHpHpImIm-γ-PyPyPyPyHp
	796)	5'-W A T T G C W-3'	PyHpHpImPy-γ-ImPyPyPyHp
	797)	5'-W A T T C T W-3'	PyHpHpPyHp-γ-PyImPyPyHp
	798)	5'-W A T T C A W-3'	PyHpHpPyPy-γ-HpImPyPyHp
	799)	5'-W A T T C G W-3'	PyHpHpPyIm-γ-PyImPyPyHp
	800)	5'-W A T T C C W-3'	PyHpHpPyPy-γ-ImImPyPyHp
	801)	5'-W A T A T T W-3'	PyHpPyHpHp-γ-PyPyHpPyHp
	802)	5'-W A T A T A W-3'	PyHpPyHpPy-γ-HpPyHpPyHp
	803)	5'-W A T A T G W-3'	PyHpPyHpIm-γ-PyPyHpPyHp
	804)	5'-W A T A T C W-3'	PyHpPyHpPy-γ-ImPyHpPyHp
	805)	5'-W A T A A T W-3'	PyHpPyPyHp-γ-PyHpHpPyHp
25	806)	5'-W A T A A A W-3'	PyHpPyPyPy-γ-HpHpHpPyHp
	807)	5'-W A T A A G W-3'	PyHpPyPyIm-γ-PyHpHpPyHp
	808)	5'-W A T A A C W-3'	PyHpPyPyPy-γ-ImHpHpPyHp
	809)	5'-W A T A G T W-3'	PyHpPyImHp-γ-PyPyHpPyHp
	810)	5'-W A T A G A W-3'	PyHpPyImPy-γ-HpPyHpPyHp
30	811)	5'-W A T A G G W-3'	PyHpPyImIm-γ-PyPyHpPyHp
	812)	5'-W A T A G C W-3'	PyHpPyImPy-γ-ImPyHpPyHp
	813)	5'-W A T A C T W-3'	PyHpPyPyHp-γ-PyImHpPyHp
	814)	5'-W A T A C A W-3'	PyHpPyPyPy-γ-HpImHpPyHp
	815)	5'-W A T A C G W-3'	PyHpPyPyIm-γ-PyImHpPyHp
35	816)	5'-W A T A C C W-3'	PyHpPyPyPy-γ-ImImHpPyHp

TABLE 39: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WATSNW-3'

	DNA sequence	aromatic amino acid sequence
5	817) 5'-W A T G T T W-3'	PyHpImHpHp- γ -PyPyPyPyHp
	818) 5'-W A T G T A W-3'	PyHpImHpPy- γ -HpPyPyPyHp
	819) 5'-W A T G T G W-3'	PyHpImHpIm- γ -PyPyPyPyHp
	820) 5'-W A T G T C W-3'	PyHpImHpPy- γ -ImPyPyPyHp
	821) 5'-W A T G A T W-3'	PyHpImPyHp- γ -PyHpPyPyHp
10	822) 5'-W A T G A A W-3'	PyHpImPyPy- γ -HpHpPyPyHp
	823) 5'-W A T G A G W-3'	PyHpImPyIm- γ -PyHpPyPyHp
	824) 5'-W A T G A C W-3'	PyHpImPyPy- γ -ImHpPyPyHp
	825) 5'-W A T G G T W-3'	PyHpImImHp- γ -PyPyPyPyHp
	826) 5'-W A T G G A W-3'	PyHpImImPy- γ -HpPyPyPyHp
	827) 5'-W A T G C T W-3'	PyHpImPyHp- γ -PyImPyPyHp
	828) 5'-W A T G C A W-3'	PyHpImPyPy- γ -HpImPyPyHp
	829) 5'-W A T G G G W-3'	PyHpImImIm- γ -PyPyPyPyHp
	830) 5'-W A T G G C W-3'	PyHpImImPy- γ -ImPyPyPyHp
	831) 5'-W A T G C G W-3'	PyHpImPyIm- γ -PyImPyPyHp
20	832) 5'-W A T G C C W-3'	PyHpImPyPy- γ -ImImPyPyHp
	833) 5'-W A T C T T W-3'	PyHpPyHpHp- γ -PyPyImPyHp
	834) 5'-W A T C T A W-3'	PyHpPyHpPy- γ -HpPyImPyHp
	835) 5'-W A T C T G W-3'	PyHpPyHpIm- γ -PyPyImPyHp
	836) 5'-W A T C T C W-3'	PyHpPyHpPy- γ -ImPyImPyHp
	837) 5'-W A T C A T W-3'	PyHpPyPyHp- γ -PyHpImPyHp
	838) 5'-W A T C A A W-3'	PyHpPyPyPy- γ -HpHpImPyHp
	839) 5'-W A T C A G W-3'	PyHpPyPyIm- γ -PyHpImPyHp
	840) 5'-W A T C A C W-3'	PyHpPyPyPy- γ -ImHpImPyHp
	841) 5'-W A T C G T W-3'	PyHpPyImHp- γ -PyPyImPyHp
30	842) 5'-W A T C G A W-3'	PyHpPyImPy- γ -HpPyImPyHp
	843) 5'-W A T C C T W-3'	PyHpPyPyHp- γ -PyImImPyHp
	844) 5'-W A T C C A W-3'	PyHpPyPyPy- γ -HpImImPyHp
	845) 5'-W A T C G G W-3'	PyHpPyImIm- γ -PyPyImPyHp
	846) 5'-W A T C G C W-3'	PyHpPyImPy- γ -ImPyImPyHp
35	847) 5'-W A T C C G W-3'	PyHpPyPyIm- γ -PyImImPyHp
	848) 5'-W A T C C C W-3'	PyHpPyPyPy- γ -ImImImPyHp

TABLE 40: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	849) 5'-W A A T T T W-3'	PyPyHpHpHp-γ-PyPyPyHpHp
5	850) 5'-W A A T T A W-3'	PyPyHpHpPy-γ-HpPyPyHpHp
	851) 5'-W A A T T G W-3'	PyPyHpHpIm-γ-PyPyPyHpHp
	852) 5'-W A A T T C W-3'	PyPyHpHpPy-γ-ImPyPyHpHp
	853) 5'-W A A T A T W-3'	PyPyHpPyHp-γ-PyHpPyHpHp
	854) 5'-W A A T A A W-3'	PyPyHpPyPy-γ-HpHpPyHpHp
10	855) 5'-W A A T A G W-3'	PyPyHpPyIm-γ-PyHpPyHpHp
	856) 5'-W A A T A C W-3'	PyPyHpPyPy-γ-ImHpPyHpHp
	857) 5'-W A A T G T W-3'	PyPyHpImHp-γ-PyPyPyHpHp
	858) 5'-W A A T G A W-3'	PyPyHpImPy-γ-HpPyPyHpHp
	859) 5'-W A A T G G W-3'	PyPyHpImIm-γ-PyPyPyHpHp
	860) 5'-W A A T G C W-3'	PyPyHpImPy-γ-ImPyPyHpHp
	861) 5'-W A A T C T W-3'	PyPyHpPyHp-γ-PyImPyHpHp
	862) 5'-W A A T C A W-3'	PyPyHpPyPy-γ-HpImPyHpHp
	863) 5'-W A A T C G W-3'	PyPyHpPyIm-γ-PyImPyHpHp
	864) 5'-W A A T C C W-3'	PyPyHpPyPy-γ-ImImPyHpHp
20	865) 5'-W A A A T T W-3'	PyPyPyHpHp-γ-PyPyHpHpHp
	866) 5'-W A A A T A W-3'	PyPyPyHpPy-γ-HpPyHpHpHp
	867) 5'-W A A A T G W-3'	PyPyPyHpIm-γ-PyPyHpHpHp
	868) 5'-W A A A T C W-3'	PyPyPyHpPy-γ-ImPyHpHpHp
	869) 5'-W A A A A T W-3'	PyPyPyPyHp-γ-PyHpHpHpHp
25	870) 5'-W A A A A A W-3'	PyPyPyPyPy-γ-HpHpHpHpHp
	871) 5'-W A A A A G W-3'	PyPyPyPyIm-γ-PyHpHpHpHp
	872) 5'-W A A A A C W-3'	PyPyPyPyPy-γ-ImHpHpHpHp
	873) 5'-W A A A G T W-3'	PyPyPyImHp-γ-PyPyHpHpHp
	874) 5'-W A A A G A W-3'	PyPyPyImPy-γ-HpPyHpHpHp
30	875) 5'-W A A A G G W-3'	PyPyPyImIm-γ-PyPyHpHpHp
	876) 5'-W A A A G C W-3'	PyPyPyImPy-γ-ImPyHpHpHp
	877) 5'-W A A A C T W-3'	PyPyPyPyHp-γ-PyImHpHpHp
	878) 5'-W A A A C A W-3'	PyPyPyPyPy-γ-HpImHpHpHp
	879) 5'-W A A A C G W-3'	PyPyPyPyIm-γ-PyImHpHpHp
35	880) 5'-W A A A C C W-3'	PyPyPyPyPy-γ-ImImHpHpHp

TABLE 41: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAASNNW-3'

	DNA sequence	aromatic amino acid sequence
	881) 5'-W A A G T T W-3'	PyPyImHpHp- γ -PyPyPyHpHp
5	882) 5'-W A A G T A W-3'	PyPyImHpPy- γ -HpPyPyHpHp
	883) 5'-W A A G T G W-3'	PyPyImHpIm- γ -PyPyPyHpHp
	884) 5'-W A A G T C W-3'	PyPyImHpPy- γ -ImPyPyHpHp
	885) 5'-W A A G A T W-3'	PyPyImPyHp- γ -PyHpPyHpHp
	886) 5'-W A A G A A W-3'	PyPyImPyPy- γ -HpHpPyHpHp
10	887) 5'-W A A G A G W-3'	PyPyImPyIm- γ -PyHpPyHpHp
	888) 5'-W A A G A C W-3'	PyPyImPyPy- γ -ImHpPyHpHp
	889) 5'-W A A G G T W-3'	PyPyImImHp- γ -PyPyPyHpHp
	890) 5'-W A A G G A W-3'	PyPyImImPy- γ -HpPyPyHpHp
	891) 5'-W A A G C T W-3'	PyPyImPyHp- γ -PyImPyHpHp
	892) 5'-W A A G C A W-3'	PyPyImPyPy- γ -HpImPyHpHp
	893) 5'-W A A G G G W-3'	PyPyImImIm- γ -PyPyPyHpHp
	894) 5'-W A A G G C W-3'	PyPyImImPy- γ -ImPyPyHpHp
	895) 5'-W A A G C G W-3'	PyPyImPyIm- γ -PyImPyHpHp
	896) 5'-W A A G C C W-3'	PyPyImPyPy- γ -ImImPyHpHp
	897) 5'-W A A C T T W-3'	PyPyPyHpHp- γ -PyPyImHpHp
	898) 5'-W A A C T A W-3'	PyPyPyHpPy- γ -HpPyImHpHp
	899) 5'-W A A C T G W-3'	PyPyPyHpIm- γ -PyPyImHpHp
	900) 5'-W A A C T C W-3'	PyPyPyHpPy- γ -ImPyImHpHp
	901) 5'-W A A C A T W-3'	PyPyPyPyHp- γ -PyHpImHpHp
25	902) 5'-W A A C A A W-3'	PyPyPyPyPy- γ -HpHpImHpHp
	903) 5'-W A A C A G W-3'	PyPyPyPyIm- γ -PyHpImHpHp
	904) 5'-W A A C A C W-3'	PyPyPyPyPy- γ -ImHpImHpHp
	905) 5'-W A A C G T W-3'	PyPyPyImHp- γ -PyPyImHpHp
	906) 5'-W A A C G A W-3'	PyPyPyImPy- γ -HpPyImHpHp
30	907) 5'-W A A C C T W-3'	PyPyPyPyHp- γ -PyImImHpHp
	908) 5'-W A A C C A W-3'	PyPyPyPyPy- γ -HpImImHpHp
	909) 5'-W A A C G G W-3'	PyPyPyImIm- γ -PyPyImHpHp
	910) 5'-W A A C G C W-3'	PyPyPyImPy- γ -ImPyImHpHp
	911) 5'-W A A C C G W-3'	PyPyPyPyIm- γ -PyImImHpHp
35	912) 5'-W A A C C C W-3'	PyPyPyPyPy- γ -ImImImHpHp

TABLE 42: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	913) 5'-W A C T T T W-3'	PyPyHpHpHp-γ-PyPyPyImHp
	914) 5'-W A C T T A W-3'	PyPyHpHpPy-γ-HpPyPyImHp
	915) 5'-W A C T T G W-3'	PyPyHpHpIm-γ-PyPyPyImHp
	916) 5'-W A C T T C W-3'	PyPyHpHpPy-γ-ImPyPyImHp
	917) 5'-W A C T A T W-3'	PyPyHpPyHp-γ-PyHpPyImHp
10	918) 5'-W A C T A A W-3'	PyPyHpPyPy-γ-HpHpPyImHp
	919) 5'-W A C T A G W-3'	PyPyHpPyIm-γ-PyHpPyImHp
	920) 5'-W A C T A C W-3'	PyPyHpPyPy-γ-ImHpPyImHp
	921) 5'-W A C T G T W-3'	PyPyHpImHp-γ-PyPyPyImHp
	922) 5'-W A C T G A W-3'	PyPyHpImPy-γ-HpPyPyImHp
	923) 5'-W A C T G G W-3'	PyPyHpImIm-γ-PyPyPyImHp
	924) 5'-W A C T G C W-3'	PyPyHpImPy-γ-ImPyPyImHp
	925) 5'-W A C T C T W-3'	PyPyHpPyHp-γ-PyImPyImHp
	926) 5'-W A C T C A W-3'	PyPyHpPyPy-γ-HpImPyImHp
	927) 5'-W A C T C G W-3'	PyPyHpPyIm-γ-PyImPyImHp
20	928) 5'-W A C T C C W-3'	PyPyHpPyPy-γ-ImImPyImHp
	929) 5'-W A C A T T W-3'	PyPyPyHpHp-γ-PyPyHpImHp
	930) 5'-W A C A T A W-3'	PyPyPyHpPy-γ-HpPyHpImHp
	931) 5'-W A C A T G W-3'	PyPyPyHpIm-γ-PyPyHpImHp
	932) 5'-W A C A T C W-3'	PyPyPyHpPy-γ-ImPyHpImHp
	933) 5'-W A C A A T W-3'	PyPyPyPyHp-γ-PyHpHpImHp
	934) 5'-W A C A A A W-3'	PyPyPyPyPy-γ-HpHpHpImHp
	935) 5'-W A C A A G W-3'	PyPyPyPyIm-γ-PyHpHpImHp
	936) 5'-W A C A A C W-3'	PyPyPyPyPy-γ-ImHpHpImHp
	937) 5'-W A C A G T W-3'	PyPyPyImHp-γ-PyPyHpImHp
30	938) 5'-W A C A G A W-3'	PyPyPyImPy-γ-HpPyHpImHp
	939) 5'-W A C A G G W-3'	PyPyPyImIm-γ-PyPyHpImHp
	940) 5'-W A C A G C W-3'	PyPyPyImPy-γ-ImPyHpImHp
	941) 5'-W A C A C T W-3'	PyPyPyPyHp-γ-PyImHpImHp
	942) 5'-W A C A C A W-3'	PyPyPyPyPy-γ-HpImHpImHp
	943) 5'-W A C A C G W-3'	PyPyPyPyIm-γ-PyImHpImHp
	944) 5'-W A C A C C W-3'	PyPyPyPyPy-γ-ImImHpImHp

TABLE 43: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACSNW-3'

	DNA sequence	aromatic amino acid sequence
5	945) 5'-W A C G T T W-3'	PyPyImHpHp-γ-PyPyPyImHp
	946) 5'-W A C G T A W-3'	PyPyImHpPy-γ-HpPyPyImHp
	947) 5'-W A C G T G W-3'	PyPyImHpIm-γ-PyPyPyImHp
	948) 5'-W A C G T C W-3'	PyPyImHpPy-γ-ImPyPyImHp
	949) 5'-W A C G A T W-3'	PyPyImPyHp-γ-PyHpPyImHp
10	950) 5'-W A C G A A W-3'	PyPyImPyPy-γ-HpHpPyImHp
	951) 5'-W A C G A G W-3'	PyPyImPyIm-γ-PyHpPyImHp
	952) 5'-W A C G A C W-3'	PyPyImPyPy-γ-ImHpPyImHp
	953) 5'-W A C G G T W-3'	PyPyImImHp-γ-PyPyPyImHp
	954) 5'-W A C G G A W-3'	PyPyImImPy-γ-HpPyPyImHp
	955) 5'-W A C G C T W-3'	PyPyImPyHp-γ-PyImPyImHp
	956) 5'-W A C G C A W-3'	PyPyImPyPy-γ-HpImPyImHp
	957) 5'-W A C C T T W-3'	PyPyPyHpHp-γ-PyPyImImHp
	958) 5'-W A C C T A W-3'	PyPyPyHpPy-γ-HpPyImImHp
	959) 5'-W A C C T G W-3'	PyPyPyHpIm-γ-PyPyImImHp
20	960) 5'-W A C C T C W-3'	PyPyPyHpPy-γ-ImPyImImHp
	961) 5'-W A C C A T W-3'	PyPyPyPyHp-γ-PyHpImImHp
	962) 5'-W A C C A A W-3'	PyPyPyPyPy-γ-HpHpImImHp
	963) 5'-W A C C A G W-3'	PyPyPyPyIm-γ-PyHpImImHp
	964) 5'-W A C C A C W-3'	PyPyPyPyPy-γ-ImHpImImHp
	965) 5'-W A C C G T W-3'	PyPyPyImHp-γ-PyPyImImHp
	966) 5'-W A C C G A W-3'	PyPyPyImPy-γ-HpPyImImHp
	967) 5'-W A C C C T W-3'	PyPyPyPyHp-γ-PyImImImHp
	968) 5'-W A C C C A W-3'	PyPyPyPyPy-γ-HpImImImHp
	969) 5'-W A C G G G W-3'	PyPyImImIm-γ-PyPyPyImHp
30	970) 5'-W A C G G C W-3'	PyPyImImPy-γ-ImPyPyImHp
	971) 5'-W A C G C G W-3'	PyPyImPyIm-γ-PyImPyImHp
	972) 5'-W A C G C C W-3'	PyPyImPyPy-γ-ImImPyImHp
	973) 5'-W A C C G G W-3'	PyPyPyImIm-γ-PyPyImImHp
	974) 5'-W A C C G C W-3'	PyPyPyImPy-γ-ImPyImImHp
35	975) 5'-W A C C C G W-3'	PyPyPyPyIm-γ-PyImImImHp
	976) 5'-W A C C C C W-3'	PyPyPyPyPy-γ-ImImImImHp

TABLE 44: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTGWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	977) 5'-W T G T T T W-3'	HpImHpHpHp-γ-PyPyPyPyPy
	978) 5'-W T G T T A W-3'	HpImHpHpPy-γ-HpPyPyPyPy
	979) 5'-W T G T T G W-3'	HpImHpHpIm-γ-PyPyPyPyPy
	980) 5'-W T G T T C W-3'	HpImHpHpPy-γ-ImPyPyPyPy
	981) 5'-W T G T A T W-3'	HpImHpPyHp-γ-PyHpPyPyPy
10	982) 5'-W T G T A A W-3'	HpImHpPyPy-γ-HpHpPyPyPy
	983) 5'-W T G T A G W-3'	HpImHpPyIm-γ-PyHpPyPyPy
	984) 5'-W T G T A C W-3'	HpImHpPyPy-γ-ImHpPyPyPy
	985) 5'-W T G T G T W-3'	HpImHpImHp-γ-PyPyPyPyPy
	986) 5'-W T G T G A W-3'	HpImHpImPy-γ-HpPyPyPyPy
	987) 5'-W T G T G G W-3'	HpImHpImIm-γ-PyPyPyPyPy
	988) 5'-W T G T G C W-3'	HpImHpImPy-γ-ImPyPyPyPy
	989) 5'-W T G T C T W-3'	HpImHpPyHp-γ-PyImPyPyPy
	990) 5'-W T G T C A W-3'	HpImHpPyPy-γ-HpImPyPyPy
	991) 5'-W T G T C G W-3'	HpImHpPyIm-γ-PyImPyPyPy
20	992) 5'-W T G T C C W-3'	HpImHpPyPy-γ-ImImPyPyPy
	993) 5'-W T G A T T W-3'	HpImPyHpHp-γ-PyPyHpPyPy
	994) 5'-W T G A T A W-3'	HpImPyHpPy-γ-HpPyHpPyPy
	995) 5'-W T G A T G W-3'	HpImPyHpIm-γ-PyPyHpPyPy
	996) 5'-W T G A T C W-3'	HpImPyHpPy-γ-ImPyHpPyPy
	997) 5'-W T G A A T W-3'	HpImPyPyHp-γ-PyHpHpPyPy
	998) 5'-W T G A A A W-3'	HpImPyPyPy-γ-HpHpHpPyPy
	999) 5'-W T G A A G W-3'	HpImPyPyIm-γ-PyHpHpPyPy
	1000) 5'-W T G A A C W-3'	HpImPyPyPy-γ-ImHpHpPyPy
	1001) 5'-W T G A G T W-3'	HpImPyImHp-γ-PyPyHpPyPy
30	1002) 5'-W T G A G A W-3'	HpImPyImPy-γ-HpPyHpPyPy
	1003) 5'-W T G A G G W-3'	HpImPyImIm-γ-PyPyHpPyPy
	1004) 5'-W T G A G C W-3'	HpImPyImPy-γ-ImPyHpPyPy
	1005) 5'-W T G A C T W-3'	HpImPyPyHp-γ-PyImHpPyPy
	1006) 5'-W T G A C A W-3'	HpImPyPyPy-γ-HpImHpPyPy
35	1007) 5'-W T G A C G W-3'	HpImPyPyIm-γ-PyImHpPyPy
	1008) 5'-W T G A C C W-3'	HpImPyPyPy-γ-ImImHpPyPy

TABLE 45: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1009) 5'-W T G G T T W-3'	HpImImHpHp-γ-PyPyPyPyPy
5	1010) 5'-W T G G T A W-3'	HpImImHpPy-γ-HpPyPyPyPy
	1011) 5'-W T G G T G W-3'	HpImImHpIm-γ-PyPyPyPyPy
	1012) 5'-W T G G T C W-3'	HpImImHpPy-γ-ImPyPyPyPy
	1013) 5'-W T G G A T W-3'	HpImImPyHp-γ-PyHpPyPyPy
	1014) 5'-W T G G A A W-3'	HpImImPyPy-γ-HpHpPyPyPy
10	1015) 5'-W T G G A G W-3'	HpImImPyIm-γ-PyHpPyPyPy
	1016) 5'-W T G G A C W-3'	HpImImPyPy-γ-ImHpPyPyPy
	1017) 5'-W T G G G T W-3'	HpImImImHp-γ-PyPyPyPyPy
	1018) 5'-W T G G G A W-3'	HpImImImPy-γ-HpPyPyPyPy
	1019) 5'-W T G G C T W-3'	HpImImPyHp-γ-PyImPyPyPy
	1020) 5'-W T G G C A W-3'	HpImImPyPy-γ-HpImPyPyPy
	1021) 5'-W T G C T T W-3'	HpImPyHpHp-γ-PyPyImPyPy
	1022) 5'-W T G C T A W-3'	HpImPyHpPy-γ-HpPyImPyPy
	1023) 5'-W T G C T G W-3'	HpImPyHpIm-γ-PyPyImPyPy
	1024) 5'-W T G C T C W-3'	HpImPyHpPy-γ-ImPyImPyPy
20	1025) 5'-W T G C A T W-3'	HpImPyPyHp-γ-PyHpImPyPy
	1026) 5'-W T G C A A W-3'	HpImPyPyPy-γ-HpHpImPyPy
	1027) 5'-W T G C A G W-3'	HpImPyPyIm-γ-PyHpImPyPy
	1028) 5'-W T G C A C W-3'	HpImPyPyPy-γ-ImHpImPyPy
	1029) 5'-W T G C G T W-3'	HpImPyImHp-γ-PyPyImPyPy
25	1030) 5'-W T G C G A W-3'	HpImPyImPy-γ-HpPyImPyPy
	1031) 5'-W T G C C T W-3'	HpImPyPyHp-γ-PyImImPyPy
	1032) 5'-W T G C C A W-3'	HpImPyPyPy-γ-HpImImPyPy
	1033) 5'-W T G G G G W-3'	HpImImImIm-γ-PyPyPyPyPy
	1034) 5'-W T G G G C W-3'	HpImImImPy-γ-ImPyPyPyPy
30	1035) 5'-W T G G C G W-3'	HpImImPyIm-γ-PyImPyPyPy
	1036) 5'-W T G G C C W-3'	HpImImPyPy-γ-ImImPyPyPy
	1037) 5'-W T G C G G W-3'	HpImPyImIm-γ-PyPyImPyPy
	1038) 5'-W T G C G C W-3'	HpImPyImPy-γ-ImPyImPyPy
	1039) 5'-W T G C C G W-3'	HpImPyPyIm-γ-PyImImPyPy
35	1040) 5'-W T G C C C W-3'	HpImPyPyPy-γ-ImImImPyPy

TABLE 46: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1041) 5'-W T T T T T W-3'	HpHpHpHpHp- γ -PyPyPyPyPy
5	1042) 5'-W T T T T A W-3'	HpHpHpHpPy- γ -HpPyPyPyPy
	1043) 5'-W T T T T G W-3'	HpHpHpHpIm- γ -PyPyPyPyPy
	1044) 5'-W T T T T C W-3'	HpHpHpHpPy- γ -ImPyPyPyPy
	1045) 5'-W T T T A T W-3'	HpHpHpPyHp- γ -PyHpPyPyPy
	1046) 5'-W T T T A A W-3'	HpHpHpPyPy- γ -HpHpPyPyPy
10	1047) 5'-W T T T A G W-3'	HpHpHpPyIm- γ -PyHpPyPyPy
	1048) 5'-W T T T A C W-3'	HpHpHpPyPy- γ -ImHpPyPyPy
	1049) 5'-W T T T G T W-3'	HpHpHpImHp- γ -PyPyPyPyPy
	1050) 5'-W T T T G A W-3'	HpHpHpImPy- γ -HpPyPyPyPy
	1051) 5'-W T T T G G W-3'	HpHpHpImIm- γ -PyPyPyPyPy
	1052) 5'-W T T T G C W-3'	HpHpHpImPy- γ -ImPyPyPyPy
	1053) 5'-W T T T C T W-3'	HpHpHpPyHp- γ -PyImPyPyPy
	1054) 5'-W T T T C A W-3'	HpHpHpPyPy- γ -HpImPyPyPy
	1055) 5'-W T T T C G W-3'	HpHpHpPyIm- γ -PyImPyPyPy
	1056) 5'-W T T T C C W-3'	HpHpHpPyPy- γ -ImImPyPyPy
20	1057) 5'-W T T A T T W-3'	HpHpPyHpHp- γ -PyPyHpPyPy
	1058) 5'-W T T A T A W-3'	HpHpPyHpPy- γ -HpPyHpPyPy
	1059) 5'-W T T A T G W-3'	HpHpPyHpIm- γ -PyPyHpPyPy
	1060) 5'-W T T A T C W-3'	HpHpPyHpPy- γ -ImPyHpPyPy
	1061) 5'-W T T A A T W-3'	HpHpPyPyHp- γ -PyHpHpPyPy
25	1062) 5'-W T T A A A W-3'	HpHpPyPyPy- γ -HpHpHpPyPy
	1063) 5'-W T T A A G W-3'	HpHpPyPyIm- γ -PyHpHpPyPy
	1064) 5'-W T T A A C W-3'	HpHpPyPyPy- γ -ImHpHpPyPy
	1065) 5'-W T T A G T W-3'	HpHpPyImHp- γ -PyPyHpPyPy
	1066) 5'-W T T A G A W-3'	HpHpPyImPy- γ -HpPyHpPyPy
30	1067) 5'-W T T A G G W-3'	HpHpPyImIm- γ -PyPyHpPyPy
	1068) 5'-W T T A G C W-3'	HpHpPyImPy- γ -ImPyHpPyPy
	1069) 5'-W T T A C T W-3'	HpHpPyPyHp- γ -PyImHpPyPy
	1070) 5'-W T T A C A W-3'	HpHpPyPyPy- γ -HpImHpPyPy
	1071) 5'-W T T A C G W-3'	HpHpPyPyIm- γ -PyImHpPyPy
35	1072) 5'-W T T A C C W-3'	HpHpPyPyPy- γ -ImImHpPyPy

TABLE 47: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1073) 5'-W T T G T T W-3'	HpHpImHpHp- γ -PyPyPyPyPy
5	1074) 5'-W T T G T A W-3'	HpHpImHpPy- γ -HpPyPyPyPy
	1075) 5'-W T T G T G W-3'	HpHpImHpIm- γ -PyPyPyPyPy
	1076) 5'-W T T G T C W-3'	HpHpImHpPy- γ -ImPyPyPyPy
	1077) 5'-W T T G A T W-3'	HpHpImPyHp- γ -PyHpPyPyPy
	1078) 5'-W T T G A A W-3'	HpHpImPyPy- γ -HpHpPyPyPy
10	1079) 5'-W T T G A G W-3'	HpHpImPyIm- γ -PyHpPyPyPy
	1080) 5'-W T T G A C W-3'	HpHpImPyPy- γ -ImHpPyPyPy
	1081) 5'-W T T G G T W-3'	HpHpImImHp- γ -PyPyPyPyPy
	1082) 5'-W T T G G A W-3'	HpHpImImPy- γ -HpPyPyPyPy
	1083) 5'-W T T G C T W-3'	HpHpImPyHp- γ -PyImPyPyPy
	1084) 5'-W T T G C A W-3'	HpHpImPyPy- γ -HpImPyPyPy
	1085) 5'-W T T G G G W-3'	HpHpImImIm- γ -PyPyPyPyPy
	1086) 5'-W T T G G C W-3'	HpHpImImPy- γ -ImPyPyPyPy
	1087) 5'-W T T G C G W-3'	HpHpImPyIm- γ -PyImPyPyPy
	1088) 5'-W T T G C C W-3'	HpHpImPyPy- γ -ImImPyPyPy
20	1089) 5'-W T T C T T W-3'	HpHpPyHpHp- γ -PyPyImPyPy
	1090) 5'-W T T C T A W-3'	HpHpPyHpPy- γ -HpPyImPyPy
	1091) 5'-W T T C T G W-3'	HpHpPyHpIm- γ -PyPyImPyPy
	1092) 5'-W T T C T C W-3'	HpHpPyHpPy- γ -ImPyImPyPy
	1093) 5'-W T T C A T W-3'	HpHpPyPyHp- γ -PyHpImPyPy
25	1094) 5'-W T T C A A W-3'	HpHpPyPyPy- γ -HpHpImPyPy
	1095) 5'-W T T C A G W-3'	HpHpPyPyIm- γ -PyHpImPyPy
	1096) 5'-W T T C A C W-3'	HpHpPyPyPy- γ -ImHpImPyPy
	1097) 5'-W T T C G T W-3'	HpHpPyImHp- γ -PyPyImPyPy
	1098) 5'-W T T C G A W-3'	HpHpPyImPy- γ -HpPyImPyPy
30	1099) 5'-W T T C C T W-3'	HpHpPyPyHp- γ -PyImImPyPy
	1100) 5'-W T T C C A W-3'	HpHpPyPyPy- γ -HpImImPyPy
	1101) 5'-W T T C G G W-3'	HpHpPyImIm- γ -PyPyImPyPy
	1102) 5'-W T T C G C W-3'	HpHpPyImPy- γ -ImPyImPyPy
	1103) 5'-W T T C C G W-3'	HpHpPyPyIm- γ -PyImImPyPy
35	1104) 5'-W T T C C C W-3'	HpHpPyPyPy- γ -ImImImPyPy

TABLE 48: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1105) 5'-W T A T T T W-3'	HpPyHpHpHp- γ -PyPyPyHpPy
5	1106) 5'-W T A T T A W-3'	HpPyHpHpPy- γ -HpPyPyHpPy
	1107) 5'-W T A T T G W-3'	HpPyHpHpIm- γ -PyPyPyHpPy
	1108) 5'-W T A T T C W-3'	HpPyHpHpPy- γ -ImPyPyHpPy
	1109) 5'-W T A T A T W-3'	HpPyHpPyHp- γ -PyHpPyHpPy
	1110) 5'-W T A T A A W-3'	HpPyHpPyPy- γ -HpHpPyHpPy
10	1111) 5'-W T A T A G W-3'	HpPyHpPyIm- γ -PyHpPyHpPy
	1112) 5'-W T A T A C W-3'	HpPyHpPyPy- γ -ImHpPyHpPy
	1113) 5'-W T A T G T W-3'	HpPyHpImHp- γ -PyPyPyHpPy
	1114) 5'-W T A T G A W-3'	HpPyHpImPy- γ -HpPyPyHpPy
	1115) 5'-W T A T G G W-3'	HpPyHpImIm- γ -PyPyPyHpPy
	1116) 5'-W T A T G C W-3'	HpPyHpImPy- γ -ImPyPyHpPy
	1117) 5'-W T A T C T W-3'	HpPyHpPyHp- γ -PyImPyHpPy
	1118) 5'-W T A T C A W-3'	HpPyHpPyPy- γ -HpImPyHpPy
	1119) 5'-W T A T C G W-3'	HpPyHpPyIm- γ -PyImPyHpPy
	1120) 5'-W T A T C C W-3'	HpPyHpPyPy- γ -ImImPyHpPy
	1121) 5'-W T A A T T W-3'	HpPyPyHpHp- γ -PyPyHpHpPy
	1122) 5'-W T A A T A W-3'	HpPyPyHpPy- γ -HpPyHpHpPy
	1123) 5'-W T A A T G W-3'	HpPyPyHpIm- γ -PyPyHpHpPy
	1124) 5'-W T A A T C W-3'	HpPyPyHpPy- γ -ImPyHpHpPy
	1125) 5'-W T A A A T W-3'	HpPyPyPyHp- γ -PyHpHpHpPy
25	1126) 5'-W T A A A A W-3'	HpPyPyPyPy- γ -HpHpHpHpPy
	1127) 5'-W T A A A G W-3'	HpPyPyPyIm- γ -PyHpHpHpPy
	1128) 5'-W T A A A C W-3'	HpPyPyPyPy- γ -ImHpHpHpPy
	1129) 5'-W T A A G T W-3'	HpPyPyImHp- γ -PyPyHpHpPy
	1130) 5'-W T A A G A W-3'	HpPyPyImPy- γ -HpPyHpHpPy
30	1131) 5'-W T A A G G W-3'	HpPyPyImIm- γ -PyPyHpHpPy
	1132) 5'-W T A A G C W-3'	HpPyPyImPy- γ -ImPyHpHpPy
	1133) 5'-W T A A C T W-3'	HpPyPyPyHp- γ -PyImHpHpPy
	1134) 5'-W T A A C A W-3'	HpPyPyPyPy- γ -HpImHpHpPy
	1135) 5'-W T A A C G W-3'	HpPyPyPyIm- γ -PyImHpHpPy
35	1136) 5'-W T A A C C W-3'	HpPyPyPyPy- γ -ImImHpHpPy

TABLE 49: 10-ring Harpin Polyamides for recognition of 7-bp 5'-WTASNNW-3'

	DNA sequence	aromatic amino acid sequence
5	1137) 5'-W T A G T T W-3'	HpPyImHpHp- γ -PyPyPyHpPy
	1138) 5'-W T A G T A W-3'	HpPyImHpPy- γ -HpPyPyHpPy
	1139) 5'-W T A G T G W-3'	HpPyImHpIm- γ -PyPyPyHpPy
	1140) 5'-W T A G T C W-3'	HpPyImHpPy- γ -ImPyPyHpPy
	1141) 5'-W T A G A T W-3'	HpPyImPyHp- γ -PyHpPyHpPy
10	1142) 5'-W T A G A A W-3'	HpPyImPyPy- γ -HpHpPyHpPy
	1143) 5'-W T A G A G W-3'	HpPyImPyIm- γ -PyHpPyHpPy
	1144) 5'-W T A G A C W-3'	HpPyImPyPy- γ -ImHpPyHpPy
	1145) 5'-W T A G G T W-3'	HpPyImImHp- γ -PyPyPyHpPy
	1146) 5'-W T A G G A W-3'	HpPyImImPy- γ -HpPyPyHpPy
	1147) 5'-W T A G C T W-3'	HpPyImPyHp- γ -PyImPyHpPy
	1148) 5'-W T A G C A W-3'	HpPyImPyPy- γ -HpImPyHpPy
	1149) 5'-W T A G G G W-3'	HpPyImImIm- γ -PyPyPyHpPy
	1150) 5'-W T A G G C W-3'	HpPyImImPy- γ -ImPyPyHpPy
	1151) 5'-W T A G C G W-3'	HpPyImPyIm- γ -PyImPyHpPy
20	1152) 5'-W T A G C C W-3'	HpPyImPyPy- γ -ImImPyHpPy
	1153) 5'-W T A C T T W-3'	HpPyPyHpHp- γ -PyPyImHpPy
	1154) 5'-W T A C T A W-3'	HpPyPyHpPy- γ -HpPyImHpPy
	1155) 5'-W T A C T G W-3'	HpPyPyHpIm- γ -PyPyImHpPy
	1156) 5'-W T A C T C W-3'	HpPyPyHpPy- γ -ImPyImHpPy
	1157) 5'-W T A C A T W-3'	HpPyPyPyHp- γ -PyHpImHpPy
	1158) 5'-W T A C A A W-3'	HpPyPyPyPy- γ -HpHpImHpPy
25	1159) 5'-W T A C A G W-3'	HpPyPyPyIm- γ -PyHpImHpPy
	1160) 5'-W T A C A C W-3'	HpPyPyPyPy- γ -ImHpImHpPy
	1161) 5'-W T A C G T W-3'	HpPyPyImHp- γ -PyPyImHpPy
	1162) 5'-W T A C G A W-3'	HpPyPyImPy- γ -HpPyImHpPy
	1163) 5'-W T A C C T W-3'	HpPyPyPyHp- γ -PyImImHpPy
	1164) 5'-W T A C C A W-3'	HpPyPyPyPy- γ -HpImImHpPy
	1165) 5'-W T A C G G W-3'	HpPyPyImIm- γ -PyPyImHpPy
30	1166) 5'-W T A C G C W-3'	HpPyPyImPy- γ -ImPyImHpPy
	1167) 5'-W T A C C G W-3'	HpPyPyPyIm- γ -PyImImHpPy
	1168) 5'-W T A C C C W-3'	HpPyPyPyPy- γ -ImImImHpPy

TABLE 50: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	1169) 5'-W T C T T T W-3'	HpPyHpHpHp- γ -PyPyPyImPy
	1170) 5'-W T C T T A W-3'	HpPyHpHpPy- γ -HpPyPyImPy
	1171) 5'-W T C T T G W-3'	HpPyHpHpIm- γ -PyPyPyImPy
	1172) 5'-W T C T T C W-3'	HpPyHpHpPy- γ -ImPyPyImPy
	1173) 5'-W T C T A T W-3'	HpPyHpPyHp- γ -PyHpPyImPy
10	1174) 5'-W T C T A A W-3'	HpPyHpPyPy- γ -HpHpPyImPy
	1175) 5'-W T C T A G W-3'	HpPyHpPyIm- γ -PyHpPyImPy
	1176) 5'-W T C T A C W-3'	HpPyHpPyPy- γ -ImHpPyImPy
	1177) 5'-W T C T G T W-3'	HpPyHpImHp- γ -PyPyPyImPy
	1178) 5'-W T C T G A W-3'	HpPyHpImPy- γ -HpPyPyImPy
	1179) 5'-W T C T G G W-3'	HpPyHpImIm- γ -PyPyPyImPy
	1180) 5'-W T C T G C W-3'	HpPyHpImPy- γ -ImPyPyImPy
	1181) 5'-W T C T C T W-3'	HpPyHpPyHp- γ -PyImPyImPy
	1182) 5'-W T C T C A W-3'	HpPyHpPyPy- γ -HpImPyImPy
	1183) 5'-W T C T C G W-3'	HpPyHpPyIm- γ -PyImPyImPy
20	1184) 5'-W T C T C C W-3'	HpPyHpPyPy- γ -ImImPyImPy
	1185) 5'-W T C A T T W-3'	HpPyPyHpHp- γ -PyPyHpImPy
	1186) 5'-W T C A T A W-3'	HpPyPyHpPy- γ -HpPyHpImPy
	1187) 5'-W T C A T G W-3'	HpPyPyHpIm- γ -PyPyHpImPy
	1188) 5'-W T C A T C W-3'	HpPyPyHpPy- γ -ImPyHpImPy
	1189) 5'-W T C A A T W-3'	HpPyPyPyHp- γ -PyHpHpImPy
	1190) 5'-W T C A A A W-3'	HpPyPyPyPy- γ -HpHpHpImPy
	1191) 5'-W T C A A G W-3'	HpPyPyPyIm- γ -PyHpHpImPy
	1192) 5'-W T C A A C W-3'	HpPyPyPyPy- γ -ImHpHpImPy
	1193) 5'-W T C A G T W-3'	HpPyPyImHp- γ -PyPyHpImPy
30	1194) 5'-W T C A G A W-3'	HpPyPyImPy- γ -HpPyHpImPy
	1195) 5'-W T C A G G W-3'	HpPyPyImIm- γ -PyPyHpImPy
	1196) 5'-W T C A G C W-3'	HpPyPyImPy- γ -ImPyHpImPy
	1197) 5'-W T C A C T W-3'	HpPyPyPyHp- γ -PyImHpImPy
	1198) 5'-W T C A C A W-3'	HpPyPyPyPy- γ -HpImHpImPy
	1199) 5'-W T C A C G W-3'	HpPyPyPyIm- γ -PyImHpImPy
	1200) 5'-W T C A C C W-3'	HpPyPyPyPy- γ -ImImHpImPy

TABLE 51: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1201) 5'-W T C G T T W-3'	HpPyImHpHp- γ -PyPyPyImPy
5	1202) 5'-W T C G T A W-3'	HpPyImHpPy- γ -HpPyPyImPy
	1203) 5'-W T C G T G W-3'	HpPyImHpIm- γ -PyPyPyImPy
	1204) 5'-W T C G T C W-3'	HpPyImHpPy- γ -ImPyPyImPy
	1205) 5'-W T C G A T W-3'	HpPyImPyHp- γ -PyHpPyImPy
	1206) 5'-W T C G A A W-3'	HpPyImPyPy- γ -HpHpPyImPy
10	1207) 5'-W T C G A G W-3'	HpPyImPyIm- γ -PyHpPyImPy
	1208) 5'-W T C G A C W-3'	HpPyImPyPy- γ -ImHpPyImPy
	1209) 5'-W T C G G T W-3'	HpPyImImHp- γ -PyPyPyImPy
	1210) 5'-W T C G G A W-3'	HpPyImImPy- γ -HpPyPyImPy
	1211) 5'-W T C G C T W-3'	HpPyImPyHp- γ -PyImPyImPy
	1212) 5'-W T C G C A W-3'	HpPyImPyPy- γ -HpImPyImPy
	1213) 5'-W T C C T T W-3'	HpPyPyHpHp- γ -PyPyImImPy
	1214) 5'-W T C C T A W-3'	HpPyPyHpPy- γ -HpPyImImPy
	1215) 5'-W T C C T G W-3'	HpPyPyHpIm- γ -PyPyImImPy
	1216) 5'-W T C C T C W-3'	HpPyPyHpPy- γ -ImPyImImPy
20	1217) 5'-W T C C A T W-3'	HpPyPyPyHp- γ -PyHpImImPy
	1218) 5'-W T C C A A W-3'	HpPyPyPyPy- γ -HpHpImImPy
	1219) 5'-W T C C A G W-3'	HpPyPyPyIm- γ -PyHpImImPy
	1220) 5'-W T C C A C W-3'	HpPyPyPyPy- γ -ImHpImImPy
	1221) 5'-W T C C G T W-3'	HpPyPyImHp- γ -PyPyImImPy
25	1222) 5'-W T C C G A W-3'	HpPyPyImPy- γ -HpPyImImPy
	1223) 5'-W T C C C T W-3'	HpPyPyPyHp- γ -PyImImImPy
	1224) 5'-W T C C C A W-3'	HpPyPyPyPy- γ -HpImImImPy
	1225) 5'-W T C G G G W-3'	HpPyImImIm- γ -PyPyPyImPy
	1226) 5'-W T C G G C W-3'	HpPyImImPy- γ -ImPyPyImPy
30	1227) 5'-W T C G C G W-3'	HpPyImPyIm- γ -PyImPyImPy
	1228) 5'-W T C G C C W-3'	HpPyImPyPy- γ -ImImPyImPy
	1229) 5'-W T C C G G W-3'	HpPyPyImIm- γ -PyPyImImPy
	1230) 5'-W T C C G C W-3'	HpPyPyImPy- γ -ImPyImImPy
	1231) 5'-W T C C C G W-3'	HpPyPyPyIm- γ -PyImImImPy
35	1232) 5'-W T C C C C W-3'	HpPyPyPyPy- γ -ImImImImPy

TABLE 52: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGGWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	243 β) 5'-W G G T T G W-3'	ImIm- β -HpIm- γ -PyPyPyPyPy
5	243 β p) 5'-W G G T T G W-3'	ImIm- β -HpIm- γ -PyPy- β -PyPy
	247 β) 5'-W G G T A G W-3'	ImIm- β -PyIm- γ -PyHpPyPyPy
	247 β p) 5'-W G G T A G W-3'	ImIm- β -PyIm- γ -PyHp- β -PyPy
	249 β) 5'-W G G T G T W-3'	ImIm- β -ImHp- γ -PyPyPyPyPy
	249 β p) 5'-W G G T G T W-3'	ImIm- β -ImHp- γ -PyPy- β -PyPy
10	250 β) 5'-W G G T G A W-3'	ImIm- β -ImPy- γ -HpPyPyPyPy
	250 β p) 5'-W G G T G A W-3'	ImIm- β -ImPy- γ -HpPy- β -PyPy
	251 β) 5'-W G G T G G W-3'	ImIm- β -ImIm- γ -PyPyPyPyPy
	251 β p) 5'-W G G T G G W-3'	ImIm- β -ImIm- γ -PyPy- β -PyPy
	252 β) 5'-W G G T G C W-3'	ImIm- β -ImPy- γ -ImPyPyPyPy
	252 β p) 5'-W G G T G C W-3'	ImIm- β -ImPy- γ -ImPy- β -PyPy
	255 β) 5'-W G G T C G W-3'	ImIm- β -PyIm- γ -PyImPyPyPy
	255 β p) 5'-W G G T C G W-3'	ImIm- β -PyIm- γ -PyIm- β -PyPy
	259 β) 5'-W G G A T G W-3'	ImIm- β -HpIm- γ -PyPyHpPyPy
	259 β p) 5'-W G G A T G W-3'	ImIm- β -HpIm- γ -PyPy- β -PyPy
20	263 β) 5'-W G G A A G W-3'	ImIm- β -PyIm- γ -PyHpHpPyPy
	263 β p) 5'-W G G A A G W-3'	ImIm- β -PyIm- γ -PyHp- β -PyPy
	265 β) 5'-W G G A G T W-3'	ImIm- β -ImHp- γ -PyPyHpPyPy
	265 β p) 5'-W G G A G T W-3'	ImIm- β -ImHp- γ -PyPy- β -PyPy
	266 β) 5'-W G G A G A W-3'	ImIm- β -ImPy- γ -HpPyHpPyPy
25	266 β p) 5'-W G G A G A W-3'	ImIm- β -ImPy- γ -HpPy- β -PyPy
	267 β) 5'-W G G A G G W-3'	ImIm- β -ImIm- γ -PyPyHpPyPy
	267 β p) 5'-W G G A G G W-3'	ImIm- β -ImIm- γ -PyPy- β -PyPy
	268 β) 5'-W G G A G C W-3'	ImIm- β -ImPy- γ -ImPyHpPyPy
	268 β p) 5'-W G G A G C W-3'	ImIm- β -ImPy- γ -ImPy- β -PyPy
30	271 β) 5'-W G G A C G W-3'	ImIm- β -PyIm- γ -PyImHpPyPy
	271 β p) 5'-W G G A C G W-3'	ImIm- β -PyIm- γ -PyIm- β -PyPy

TABLE 53: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGGSNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	273 β) 5'-W G G G T T W-3'	ImImIm- β -Hp- γ -PyPyPyPyPy
5	273 β p) 5'-W G G G T T W-3'	ImImIm- β -Hp- γ -Py- β -PyPyPy
	274 β) 5'-W G G G T A W-3'	ImImIm- β -Py- γ -HpPyPyPyPy
	274 β p) 5'-W G G G T A W-3'	ImImIm- β -Py- γ -Hp- β -PyPyPy
	275 β) 5'-W G G G T G W-3'	ImImIm- β -Im- γ -PyPyPyPyPy
	275 β p) 5'-W G G G T G W-3'	ImImIm- β -Im- γ -Py- β -PyPyPy
10	276 β) 5'-W G G G T C W-3'	ImImIm- β -Py- γ -ImPyPyPyPy
	276 β p) 5'-W G G G T C W-3'	ImImIm- β -Py- γ -Im- β -PyPyPy
	277 β) 5'-W G G G A T W-3'	ImImIm- β -Hp- γ -PyHpPyPyPy
	277 β p) 5'-W G G G A T W-3'	ImImIm- β -Hp- γ -Py- β -PyPyPy
	278 β) 5'-W G G G A A W-3'	ImImIm- β -Py- γ -HpHpPyPyPy
	278 β p) 5'-W G G G A A W-3'	ImImIm- β -Py- γ -Hp- β -PyPyPy
	279 β) 5'-W G G G A G W-3'	ImImIm- β -Im- γ -PyHpPyPyPy
	279 β p) 5'-W G G G A G W-3'	ImImIm- β -Im- γ -Py- β -PyPyPy
	280 β) 5'-W G G G A C W-3'	ImImIm- β -Py- γ -ImHpPyPyPy
	280 β p) 5'-W G G G A C W-3'	ImImIm- β -Py- γ -Im- β -PyPyPy
	283 β) 5'-W G G G C T W-3'	ImImIm- β -Hp- γ -PyImPyPyPy
	284 β) 5'-W G G G C A W-3'	ImImIm- β -Py- γ -HpImPyPyPy
	285 β) 5'-W G G C T T W-3'	ImImPyHpHp- γ -Py- β -ImPyPy
	285 β p) 5'-W G G C T T W-3'	ImImPy- β -Hp- γ -Py- β -ImPyPy
	286 β) 5'-W G G C T A W-3'	ImImPyHpPy- γ -Hp- β -ImPyPy
25	286 β p) 5'-W G G C T A W-3'	ImImPy- β -Py- γ -Hp- β -ImPyPy
	287 β) 5'-W G G C T G W-3'	ImIm- β -HpIm- γ -Py- β -ImPyPy
	288 β) 5'-W G G C T C W-3'	ImImPyHpPy- γ -Im- β -ImPyPy
	288 β p) 5'-W G G C T C W-3'	ImImPy- β -Py- γ -Im- β -ImPyPy
	289 β) 5'-W G G C A T W-3'	ImImPyPyHp- γ -Py- β -ImPyPy
30	289 β p) 5'-W G G C A T W-3'	ImImPy- β -Hp- γ -Py- β -ImPyPy
	290 β) 5'-W G G C A A W-3'	ImImPyPyPy- γ -Hp- β -ImPyPy
	290 β p) 5'-W G G C A A W-3'	ImImPy- β -Py- γ -Hp- β -ImPyPy

TABLE 53 (cont.): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGGSNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	291 β) 5'-W G G C A G W-3'	ImIm- β -PyIm- γ -Py- β -ImPyPy
	292 β) 5'-W G G C A C W-3'	ImImPyPyPy- γ -Im- β -ImPyPy
5	292 β_p) 5'-W G G C A C W-3'	ImImPy- β -Py- γ -Im- β -ImPyPy
	293 β) 5'-W G G C G T W-3'	ImIm- β -ImHp- γ -Py- β -ImPyPy
	294 β) 5'-W G G C G A W-3'	ImIm- β -ImPy- γ -Hp- β -ImPyPy
	295 β) 5'-W G G C C T W-3'	ImImPyPyHp- γ -PyImIm- β -Py
	296 β) 5'-W G G C C A W-3'	ImImPyPyPy- γ -HpImIm- β -Py
10	G19 β) 5'-W G G G C G W-3'	ImImIm- β -Im- γ -PyImPyPyPy
	G20 β) 5'-W G G G C C W-3'	ImImIm- β -Py- γ -ImImPyPyPy
	G21 β) 5'-W G G C G G W-3'	ImIm- β -ImIm- γ -Py- β -ImPyPy
	G22 β) 5'-W G G C G C W-3'	ImIm- β -ImPy- γ -Im- β -ImPyPy
	G23 β) 5'-W G G C C G W-3'	ImIm- β -PyIm- γ -PyImIm- β -Py
	G24 β) 5'-W G G C C C W-3'	ImImPyPyPy- γ -ImImIm- β -Py

TABLE 54: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGTWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	299 β) 5'-W G T T T G W-3'	ImHp- β -HpIm- γ -PyPyPyPyPy
	299 β p) 5'-W G T T T G W-3'	ImHp- β -HpIm- γ -PyPy- β -PyPy
5	303 β) 5'-W G T T A G W-3'	ImHp- β -PyIm- γ -PyHpPyPyPy
	303 β p) 5'-W G T T A G W-3'	ImHp- β -PyIm- γ -PyHp- β -PyPy
	305 β) 5'-W G T T G T W-3'	ImHp- β -ImHp- γ -PyPyPyPyPy
	305 β p) 5'-W G T T G T W-3'	ImHp- β -ImHp- γ -PyPy- β -PyPy
	306 β) 5'-W G T T G A W-3'	ImHp- β -ImPy- γ -HpPyPyPyPy
10	306 β p) 5'-W G T T G A W-3'	ImHp- β -ImPy- γ -HpPy- β -PyPy
	307 β) 5'-W G T T G G W-3'	ImHp- β -ImIm- γ -PyPyPyPyPy
	307 β p) 5'-W G T T G G W-3'	ImHp- β -ImIm- γ -PyPy- β -PyPy
	308 β) 5'-W G T T G C W-3'	ImHp- β -ImPy- γ -ImPyPyPyPy
	308 β p) 5'-W G T T G C W-3'	ImHp- β -ImPy- γ -ImPy- β -PyPy
	311 β) 5'-W G T T C G W-3'	ImHp- β -PyIm- γ -PyImPyPyPy
	311 β p) 5'-W G T T C G W-3'	ImHp- β -PyIm- γ -PyIm- β -PyPy
	315 β) 5'-W G T A T G W-3'	ImHp- β -HpIm- γ -PyPyHpPyPy
	315 β p) 5'-W G T A T G W-3'	ImHp- β -HpIm- γ -PyPy- β -PyPy
	319 β) 5'-W G T A A G W-3'	ImHp- β -PyIm- γ -PyHpHpPyPy
20	319 β p) 5'-W G T A A G W-3'	ImHp- β -PyIm- γ -PyHp- β -PyPy
	321 β) 5'-W G T A G T W-3'	ImHp- β -ImHp- γ -PyPyHpPyPy
	321 β p) 5'-W G T A G T W-3'	ImHp- β -ImHp- γ -PyPy- β -PyPy
	322 β) 5'-W G T A G A W-3'	ImHp- β -ImPy- γ -HpPyHpPyPy
	322 β p) 5'-W G T A G A W-3'	ImHp- β -ImPy- γ -HpPy- β -PyPy
25	323 β) 5'-W G T A G G W-3'	ImHp- β -ImIm- γ -PyPyHpPyPy
	323 β p) 5'-W G T A G G W-3'	ImHp- β -ImIm- γ -PyPy- β -PyPy
	324 β) 5'-W G T A G C W-3'	ImHp- β -ImPy- γ -ImPyHpPyPy
	324 β p) 5'-W G T A G C W-3'	ImHp- β -ImPy- γ -ImPy- β -PyPy
	327 β) 5'-W G T A C G W-3'	ImHp- β -PyIm- γ -PyImHpPyPy
30	327 β p) 5'-W G T A C G W-3'	ImHp- β -PyIm- γ -PyIm- β -PyPy

TABLE 55: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGTSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	329 β) 5'-W G T G T T W-3'	Im- β -ImHpHp- γ -PyPyPyPyPy
5	329 β p) 5'-W G T G T T W-3'	Im- β -ImHpHp- γ -PyPyPy- β -Py
	330 β) 5'-W G T G T A W-3'	Im- β -ImHpPy- γ -HpPyPyPyPy
	330 β p) 5'-W G T G T A W-3'	Im- β -ImHpPy- γ -HpPyPy- β -Py
	331 β) 5'-W G T G T G W-3'	Im- β -ImHpIm- γ -PyPyPyPyPy
	331 β p) 5'-W G T G T G W-3'	Im- β -ImHpIm- γ -PyPyPy- β -Py
10	332 β) 5'-W G T G T C W-3'	Im- β -ImHpPy- γ -ImPyPyPyPy
	332 β p) 5'-W G T G T C W-3'	Im- β -ImHpPy- γ -ImPyPy- β -Py
	333 β) 5'-W G T G A T W-3'	Im- β -ImPyHp- γ -PyHpPyPyPy
	333 β p) 5'-W G T G A T W-3'	Im- β -ImPyHp- γ -PyHpPy- β -Py
	334 β) 5'-W G T G A A W-3'	Im- β -ImPyPy- γ -HpHpPyPyPy
	334 β p) 5'-W G T G A A W-3'	Im- β -ImPyPy- γ -HpHpPy- β -Py
	335 β) 5'-W G T G A G W-3'	Im- β -ImPyIm- γ -PyHpPyPyPy
	335 β p) 5'-W G T G A G W-3'	Im- β -ImPyIm- γ -PyHpPy- β -Py
	336 β) 5'-W G T G A C W-3'	Im- β -ImPyPy- γ -ImHpPyPyPy
	336 β p) 5'-W G T G A C W-3'	Im- β -ImPyPy- γ -ImHpPy- β -Py
20	337 β) 5'-W G T G G T W-3'	Im- β -ImImHp- γ -PyPyPyPyPy
	337 β p) 5'-W G T G G T W-3'	Im- β -ImImHp- γ -PyPyPy- β -Py
	338 β) 5'-W G T G G A W-3'	Im- β -ImImPy- γ -HpPyPyPyPy
	338 β p) 5'-W G T G G A W-3'	Im- β -ImImPy- γ -HpPyPy- β -Py
	339 β) 5'-W G T G C T W-3'	Im- β -ImPyHp- γ -PyImPyPyPy
25	339 β p) 5'-W G T G C T W-3'	Im- β -ImPyHp- γ -PyImPy- β -Py
	340 β) 5'-W G T G C A W-3'	Im- β -ImPyPy- γ -HpImPyPyPy
	340 β p) 5'-W G T G C A W-3'	Im- β -ImPyPy- γ -HpImPy- β -Py
	341 β) 5'-W G T G G G W-3'	Im- β -ImImIm- γ -PyPyPyPyPy
	341 β p) 5'-W G T G G G W-3'	Im- β -ImImIm- γ -PyPyPy- β -Py
30	342 β) 5'-W G T G G C W-3'	Im- β -ImImPy- γ -ImPyPyPyPy
	342 β p) 5'-W G T G G C W-3'	Im- β -ImImPy- γ -ImPyPy- β -Py
	343 β) 5'-W G T G C G W-3'	Im- β -ImPyIm- γ -PyImPyPyPy

TABLE 55 (cont.): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGTSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	343 β p) 5'-W G T G C G W-3'	Im- β -ImPyIm- γ -PyImPy- β -Py
	344 β) 5'-W G T G C C W-3'	Im- β -ImPyPy- γ -ImImPyPyPy
5	344 β p) 5'-W G T G C C W-3'	Im- β -ImPyPy- γ -ImImPy- β -Py
	345 β) 5'-W G T C T T W-3'	ImHpPyHpHp- γ -Py- β -ImPyPy
	345 β p) 5'-W G T C T T W-3'	ImHpPy- β -Hp- γ -Py- β -ImPyPy
	346 β) 5'-W G T C T A W-3'	ImHpPyHpPy- γ -Hp- β -ImPyPy
	346 β p) 5'-W G T C T A W-3'	ImHpPy- β -Py- γ -Hp- β -ImPyPy
10	347 β) 5'-W G T C T G W-3'	ImHp- β -HpIm- γ -Py- β -ImPyPy
	348 β) 5'-W G T C T C W-3'	ImHpPyHpPy- γ -Im- β -ImPyPy
	348 β p) 5'-W G T C T C W-3'	ImHpPy- β -Py- γ -Im- β -ImPyPy
	349 β) 5'-W G T C A T W-3'	ImHpPyPyHp- γ -Py- β -ImPyPy
	349 β p) 5'-W G T C A T W-3'	ImHpPyPyHp- γ -Py- β -ImPyPy
	350 β) 5'-W G T C A A W-3'	ImHpPyPyPy- γ -Hp- β -ImPyPy
	350 β p) 5'-W G T C A A W-3'	ImHpPy- β -Py- γ -Hp- β -ImPyPy
	351 β) 5'-W G T C A G W-3'	ImHp- β -PyIm- γ -Py- β -ImPyPy
	352 β) 5'-W G T C A C W-3'	ImHpPyPyPy- γ -Im- β -ImPyPy
	352 β p) 5'-W G T C A C W-3'	ImHpPy- β -Py- γ -Im- β -ImPyPy
	353 β) 5'-W G T C G T W-3'	ImHp- β -ImHp- γ -Py- β -ImPyPy
	354 β) 5'-W G T C G A W-3'	ImHp- β -ImPy- γ -Hp- β -ImPyPy
	355 β) 5'-W G T C C T W-3'	ImHpPyPyHp- γ -PyImIm- β -Py
	355 β p) 5'-W G T C C T W-3'	Im- β -PyPyHp- γ -PyImIm- β -Py
	356 β) 5'-W G T C C A W-3'	ImHpPyPyPy- γ -HpImIm- β -Py
25	356 β p) 5'-W G T C C A W-3'	Im- β -PyPyPy- γ -HpImIm- β -Py
	357 β) 5'-W G T C G G W-3'	ImHp- β -ImIm- γ -Py- β -ImPyPy
	358 β) 5'-W G T C G C W-3'	ImHp- β -ImPy- γ -Im- β -ImPyPy
	359 β) 5'-W G T C C G W-3'	ImHp- β -PyIm- γ -PyImIm- β -Py
	360 β) 5'-W G T C C C W-3'	ImHpPyPyPy- γ -ImImIm- β -Py
30	360 β p) 5'-W G T C C C W-3'	Im- β -PyPyPy- γ -ImImIm- β -Py

TABLE 56: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGAWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	363 β) 5'-W G A T T G W-3'	ImPy- β -HpIm- γ -PyPyPyHpPy
	363 β p) 5'-W G A T T G W-3'	ImPy- β -HpIm- γ -PyPy- β -HpPy
5	367 β) 5'-W G A T A G W-3'	ImPy- β -PyIm- γ -PyHpPyHpPy
	367 β p) 5'-W G A T A G W-3'	ImPy- β -PyIm- γ -PyHp- β -HpPy
	369 β) 5'-W G A T G T W-3'	ImPy- β -ImHp- γ -PyPyPyHpPy
	369 β p) 5'-W G A T G T W-3'	ImPy- β -ImHp- γ -PyPy- β -HpPy
	370 β) 5'-W G A T G A W-3'	ImPy- β -ImPy- γ -HpPyPyHpPy
10	370 β p) 5'-W G A T G A W-3'	ImPy- β -ImPy- γ -HpPy- β -HpPy
	371 β) 5'-W G A T G G W-3'	ImPy- β -ImIm- γ -PyPyPyHpPy
	371 β p) 5'-W G A T G G W-3'	ImPy- β -ImIm- γ -PyPy- β -HpPy
	372 β) 5'-W G A T G C W-3'	ImPy- β -ImPy- γ -ImPyPyHpPy
	372 β p) 5'-W G A T G C W-3'	ImPy- β -ImPy- γ -ImPy- β -HpPy
	375 β) 5'-W G A T C G W-3'	ImPy- β -PyIm- γ -PyImPyHpPy
	375 β p) 5'-W G A T C G W-3'	ImPy- β -PyIm- γ -PyIm- β -HpPy
	379 β) 5'-W G A A T G W-3'	ImPy- β -HpIm- γ -PyPyHpHpPy
	379 β p) 5'-W G A A T G W-3'	ImPy- β -HpIm- γ -PyPy- β -HpPy
	383 β) 5'-W G A A A G W-3'	ImPy- β -PyIm- γ -PyHpHpHpPy
20	383 β p) 5'-W G A A A G W-3'	ImPy- β -PyIm- γ -PyHp- β -HpPy
	385 β) 5'-W G A A G T W-3'	ImPy- β -ImHp- γ -PyPyHpHpPy
	385 β p) 5'-W G A A G T W-3'	ImPy- β -ImHp- γ -PyPy- β -HpPy
	386 β) 5'-W G A A G A W-3'	ImPy- β -ImPy- γ -HpPyHpHpPy
	386 β p) 5'-W G A A G A W-3'	ImPy- β -ImPy- γ -HpPy- β -HpPy
25	387 β) 5'-W G A A G G W-3'	ImPy- β -ImIm- γ -PyPyHpHpPy
	387 β p) 5'-W G A A G G W-3'	ImPy- β -ImIm- γ -PyPy- β -HpPy
	388 β) 5'-W G A A G C W-3'	ImPy- β -ImPy- γ -ImPyHpHpPy
	388 β p) 5'-W G A A G C W-3'	ImPy- β -ImPy- γ -ImPy- β -HpPy
	391 β) 5'-W G A A C G W-3'	ImPy- β -PyIm- γ -PyImHpHpPy
30	391 β p) 5'-W G A A C G W-3'	ImPy- β -PyIm- γ -PyIm- β -HpPy

TABLE 57: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	393 β) 5'-W G A G T T W-3'	Im- β -ImHpHp- γ -PyPyPyHpPy
	394 β p) 5'-W G A G T A W-3'	Im- β -ImHpPy- γ -HpPyPy- β -Py
5	395 β) 5'-W G A G T G W-3'	Im- β -ImHpIm- γ -PyPyPyHpPy
	395 β p) 5'-W G A G T G W-3'	Im- β -ImHpIm- γ -PyPyPy- β -Py
	396 β) 5'-W G A G T C W-3'	Im- β -ImHpPy- γ -ImPyPyHpPy
	396 β p) 5'-W G A G T C W-3'	Im- β -ImHpPy- γ -ImPyPy- β -Py
	397 β) 5'-W G A G A T W-3'	Im- β -ImPyHp- γ -PyHpPyHpPy
10	397 β p) 5'-W G A G A T W-3'	Im- β -ImPyHp- γ -PyHpPy- β -Py
	398 β) 5'-W G A G A A W-3'	Im- β -ImPyPy- γ -HpHpPyHpPy
	398 β p) 5'-W G A G A A W-3'	Im- β -ImPyPy- γ -HpHpPy- β -Py
	399 β) 5'-W G A G A G W-3'	Im- β -ImPyIm- γ -PyHpPyHpPy
	399 β p) 5'-W G A G A G W-3'	Im- β -ImPyIm- γ -PyHpPy- β -Py
	400 β) 5'-W G A G A C W-3'	Im- β -ImPyPy- γ -ImHpPyHpPy
	400 β p) 5'-W G A G A C W-3'	Im- β -ImPyPy- γ -ImHpPy- β -Py
	401 β) 5'-W G A G G T W-3'	Im- β -ImImHp- γ -PyPyPyHpPy
	401 β p) 5'-W G A G G T W-3'	Im- β -ImImHp- γ -PyPyPy- β -Py
	402 β) 5'-W G A G G A W-3'	Im- β -ImImPy- γ -HpPyPyHpPy
20	402 β p) 5'-W G A G G A W-3'	Im- β -ImImPy- γ -HpPyPy- β -Py
	403 β) 5'-W G A G C T W-3'	Im- β -ImPyHp- γ -PyImPyHpPy
	403 β p) 5'-W G A G C T W-3'	Im- β -ImPyHp- γ -PyImPy- β -Py
	404 β) 5'-W G A G C A W-3'	Im- β -ImPyPy- γ -HpImPyHpPy
	404 β p) 5'-W G A G C A W-3'	Im- β -ImPyPy- γ -HpImPy- β -Py
25	405 β) 5'-W G A G G G W-3'	Im- β -ImImIm- γ -PyPyPyHpPy
	405 β p) 5'-W G A G G G W-3'	Im- β -ImImIm- γ -PyPyPy- β -Py
	406 β) 5'-W G A G G C W-3'	Im- β -ImImPy- γ -ImPyPyHpPy
	406 β p) 5'-W G A G G C W-3'	Im- β -ImImPy- γ -ImPyPy- β -Py
	407 β) 5'-W G A G C G W-3'	Im- β -ImPyIm- γ -PyImPyHpPy
30	407 β p) 5'-W G A G C G W-3'	Im- β -ImPyIm- γ -PyImPy- β -Py
	408 β) 5'-W G A G C C W-3'	Im- β -ImPyPy- γ -ImImPyHpPy
	408 β p) 5'-W G A G C C W-3'	Im- β -ImPyPy- γ -ImImPy- β -Py

TABLE 57 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	409 β) 5'-W G A C T T W-3'	ImPyPyHpHp- γ -Py- β -ImHpPy
	409 β p) 5'-W G A C T T W-3'	ImPyPy- β -Hp- γ -Py- β -ImHpPy
5	410 β) 5'-W G A C T A W-3'	ImPyPyHpPy- γ -Hp- β -ImHpPy
	410 β p) 5'-W G A C T A W-3'	ImPyPy- β -Py- γ -Hp- β -ImHpPy
	411 β) 5'-W G A C T G W-3'	ImPy- β -HpIm- γ -Py- β -ImHpPy
	412 β) 5'-W G A C T C W-3'	ImPyPyHpPy- γ -Im- β -ImHpPy
	412 β p) 5'-W G A C T C W-3'	ImPyPy- β -Py- γ -Im- β -ImHpPy
10	413 β) 5'-W G A C A T W-3'	ImPyPyPyHp- γ -Py- β -ImHpPy
	413 β p) 5'-W G A C A T W-3'	ImPyPy- β -Hp- γ -Py- β -ImHpPy
	414 β) 5'-W G A C A A W-3'	ImPyPyPyPy- γ -Hp- β -ImHpPy
	414 β p) 5'-W G A C A A W-3'	ImPyPy- β -Py- γ -Hp- β -ImHpPy
	415 β) 5'-W G A C A G W-3'	ImPy- β -PyIm- γ -Py- β -ImHpPy
	416 β) 5'-W G A C A C W-3'	ImPyPyPyPy- γ -Im- β -ImHpPy
	416 β p) 5'-W G A C A C W-3'	ImPyPy- β -Py- γ -Im- β -ImHpPy
	417 β) 5'-W G A C G T W-3'	ImPy- β -ImHp- γ -Py- β -ImHpPy
	418 β) 5'-W G A C G A W-3'	ImPy- β -ImPy- γ -Hp- β -ImHpPy
	419 β) 5'-W G A C C T W-3'	Im- β -PyPyHp- γ -PyImIm- β -Py
	419 β p) 5'-W G A C C T W-3'	ImPyPyPyHp- γ -PyImIm- β -Py
	420 β) 5'-W G A C C A W-3'	Im- β -PyPyPy- γ -HpImIm- β -Py
	420 β p) 5'-W G A C C A W-3'	ImPyPyPyPy- γ -HpImIm- β -Py
	421 β) 5'-W G A C G G W-3'	ImPy- β -ImIm- γ -Py- β -ImHpPy
	422 β) 5'-W G A C G C W-3'	ImPy- β -ImPy- γ -Im- β -ImHpPy
25	423 β) 5'-W G A C C G W-3'	ImPy- β -PyIm- γ -PyImIm- β -Py
	424 β) 5'-W G A C C C W-3'	ImPyPyPyPy- γ -ImImIm- β -Py
	424 β p) 5'-W G A C C C W-3'	Im- β -PyPyPy- γ -ImImIm- β -Py

TABLE 58: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	425 β) 5'-W G C T T T W-3'	ImPyHpHpHp- γ -PyPy- β -ImPy
5	425 β p) 5'-W G C T T T W-3'	ImPy- β -HpHp- γ -PyPy- β -ImPy
	426 β) 5'-W G C T T A W-3'	ImPyHpHpPy- γ -HpPy- β -ImPy
	426 β p) 5'-W G C T T A W-3'	ImPy- β -HpPy- γ -HpPy- β -ImPy
	427 β) 5'-W G C T T G W-3'	ImPy- β -HpIm- γ -PyPy- β -ImPy
	428 β) 5'-W G C T T C W-3'	ImPyHpHpPy- γ -ImPy- β -ImPy
10	428 β p) 5'-W G C T T C W-3'	ImPy- β -HpPy- γ -ImPy- β -ImPy
	429 β) 5'-W G C T A T W-3'	ImPyHpPyHp- γ -PyHp- β -ImPy
	429 β p) 5'-W G C T A T W-3'	ImPy- β -PyHp- γ -PyHp- β -ImPy
	430 β) 5'-W G C T A A W-3'	ImPyHpPyPy- γ -HpHp- β -ImPy
	430 β p) 5'-W G C T A A W-3'	ImPy- β -PyPy- γ -HpHp- β -ImPy
	431 β) 5'-W G C T A G W-3'	ImPy- β -PyIm- γ -PyHp- β -ImPy
	432 β) 5'-W G C T A C W-3'	ImPyHpPyPy- γ -ImHp- β -ImPy
	432 β p) 5'-W G C T A C W-3'	ImPy- β -PyPy- γ -ImHp- β -ImPy
	433 β) 5'-W G C T G T W-3'	ImPy- β -ImHp- γ -PyPy- β -ImPy
	434 β) 5'-W G C T G A W-3'	ImPy- β -ImPy- γ -HpPy- β -ImPy
20	435 β) 5'-W G C T G G W-3'	ImPy- β -ImIm- γ -PyPy- β -ImPy
	436 β) 5'-W G C T G C W-3'	ImPy- β -ImPy- γ -ImPy- β -ImPy
	437 β) 5'-W G C T C T W-3'	ImPyHpPyHp- γ -PyIm- β -ImPy
	437 β p) 5'-W G C T C T W-3'	ImPy- β -PyHp- γ -PyIm- β -ImPy
	438 β) 5'-W G C T C A W-3'	ImPyHpPyPy- γ -HpIm- β -ImPy
25	438 β p) 5'-W G C T C A W-3'	ImPy- β -PyPy- γ -HpIm- β -ImPy
	439 β) 5'-W G C T C G W-3'	ImPy- β -PyIm- γ -PyIm- β -ImPy
	440 β) 5'-W G C T C C W-3'	ImPyHpPyPy- γ -ImIm- β -ImPy
	440 β p) 5'-W G C T C C W-3'	ImPy- β -PyPy- γ -ImIm- β -ImPy
	441 β) 5'-W G C A T T W-3'	ImPyPyHpHp- γ -PyPy- β -ImPy
30	441 β p) 5'-W G C A T T W-3'	ImPy- β -HpHp- γ -PyPy- β -ImPy
	442 β) 5'-W G C A T A W-3'	ImPyPyHpPy- γ -HpPy- β -ImPy
	442 β p) 5'-W G C A T A W-3'	ImPy- β -HpPy- γ -HpPy- β -ImPy
	443 β) 5'-W G C A T G W-3'	ImPy- β -HpIm- γ -PyPy- β -ImPy

TABLE 58 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	444 β) 5'-W G C A T C W-3'	ImPyPyHpPy- γ -ImPy- β -ImPy
	444 β p) 5'-W G C A T C W-3'	ImPy- β -HpPy- γ -ImPy- β -ImPy
5	445 β) 5'-W G C A A T W-3'	ImPyPyPyHpPy- γ -PyHp- β -ImPy
	445 β p) 5'-W G C A A T W-3'	ImPy- β -PyHp- γ -PyHp- β -ImPy
	446 β) 5'-W G C A A A W-3'	ImPyPyPyPy- γ -HpHp- β -ImPy
	446 β p) 5'-W G C A A A W-3'	ImPy- β -PyPy- γ -HpHp- β -ImPy
	447 β) 5'-W G C A A G W-3'	ImPy- β -PyIm- γ -PyHp- β -ImPy
10	448 β) 5'-W G C A A C W-3'	ImPyPyPyPy- γ -ImHp- β -ImPy
	448 β p) 5'-W G C A A C W-3'	ImPy- β -PyPy- γ -ImHp- β -ImPy
	449 β) 5'-W G C A G T W-3'	ImPy- β -ImHp- γ -PyPy- β -ImPy
	450 β) 5'-W G C A G A W-3'	ImPy- β -ImPy- γ -HpPy- β -ImPy
	451 β) 5'-W G C A G G W-3'	ImPy- β -ImIm- γ -PyPy- β -ImPy
15	452 β) 5'-W G C A G C W-3'	ImPy- β -ImPy- γ -ImPy- β -ImPy
	453 β) 5'-W G C A C T W-3'	ImPyPyPyHpPy- γ -PyIm- β -ImPy
	453 β p) 5'-W G C A C T W-3'	ImPy- β -PyHp- γ -PyIm- β -ImPy
	454 β) 5'-W G C A C A W-3'	ImPyPyPyPy- γ -HpIm- β -ImPy
	454 β p) 5'-W G C A C A W-3'	ImPy- β -PyPy- γ -HpIm- β -ImPy
20	455 β) 5'-W G C A C G W-3'	ImPy- β -PyIm- γ -PyIm- β -ImPy
	456 β) 5'-W G C A C C W-3'	ImPyPyPyPy- γ -ImIm- β -ImPy
	456 β p) 5'-W G C A C C W-3'	ImPy- β -PyPy- γ -ImIm- β -ImPy

TABLE 59: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	457 β) 5'-W G C G T T W-3'	Im- β -ImHpHp- γ -PyPy- β -ImPy
5	458 β) 5'-W G C G T A W-3'	Im- β -ImHpPy- γ -HpPy- β -ImPy
	459 β) 5'-W G C G T G W-3'	Im- β -ImHpIm- γ -PyPy- β -ImPy
	460 β) 5'-W G C G T C W-3'	Im- β -ImHpPy- γ -ImPy- β -ImPy
	461 β) 5'-W G C G A T W-3'	Im- β -ImPyHp- γ -PyHp- β -ImPy
	462 β) 5'-W G C G A A W-3'	Im- β -ImPyPy- γ -HpHp- β -ImPy
10	463 β) 5'-W G C G A G W-3'	Im- β -ImPyIm- γ -PyHp- β -ImPy
	464 β) 5'-W G C G A C W-3'	Im- β -ImPyPy- γ -ImHp- β -ImPy
	465 β) 5'-W G C G G T W-3'	Im- β -ImImHp- γ -PyPy- β -ImPy
	466 β) 5'-W G C G G A W-3'	Im- β -ImImPy- γ -HpPy- β -ImPy
	467 β) 5'-W G C G C T W-3'	Im- β -ImPyHp- γ -PyIm- β -ImPy
	468 β) 5'-W G C G C A W-3'	Im- β -ImPyPy- γ -HpIm- β -ImPy
	469 β) 5'-W G C C T T W-3'	ImPyPyHpHp- γ -Py- β -ImImPy
	469 β p) 5'-W G C C T T W-3'	ImPyPy- β -Hp- γ -Py- β -ImImPy
	470 β) 5'-W G C C T A W-3'	ImPyPyHpPy- γ -Hp- β -ImImPy
	470 β p) 5'-W G C C T A W-3'	ImPyPy- β -Py- γ -Hp- β -ImImPy
20	471 β) 5'-W G C C T G W-3'	ImPy- β -HpIm- γ -Py- β -ImImPy
	472 β) 5'-W G C C T C W-3'	ImPyPyHpPy- γ -Im- β -ImImPy
	472 β p) 5'-W G C C T C W-3'	ImPyPy- β -Py- γ -Im- β -ImImPy
	473 β) 5'-W G C C A T W-3'	ImPyPyPyHp- γ -Py- β -ImImPy
	473 β p) 5'-W G C C A T W-3'	ImPyPy- β -Hp- γ -Py- β -ImImPy
25	474 β) 5'-W G C C A A W-3'	ImPyPyPyPy- γ -Hp- β -ImImPy
	474 β p) 5'-W G C C A A W-3'	ImPyPy- β -Py- γ -Hp- β -ImImPy
	475 β) 5'-W G C C A G W-3'	ImPy- β -PyIm- γ -Py- β -ImImPy
	476 β) 5'-W G C C A C W-3'	ImPyPyPyPy- γ -Im- β -ImImPy
	476 β p) 5'-W G C C A C W-3'	ImPyPy- β -Py- γ -Im- β -ImImPy
30	477 β) 5'-W G C C G T W-3'	ImPy- β -ImHp- γ -Py- β -ImImPy
	478 β) 5'-W G C C G A W-3'	ImPy- β -ImPy- γ -Hp- β -ImImPy

TABLE 59 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WGCSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	G25 β) 5'-W G C G G G W-3'	Im- β -ImImIm- γ -PyPy- β -ImPy
	G26 β) 5'-W G C G G C W-3'	Im- β -ImImPy- γ -ImPy- β -ImPy
5	G27 β) 5'-W G C G C G W-3'	Im- β -ImPyIm- γ -PyIm- β -ImPy
	G28 β) 5'-W G C G C C W-3'	Im- β -ImPyPy- γ -ImIm- β -ImPy
	G29 β) 5'-W G C C G G W-3'	ImPy- β -ImIm- γ -Py- β -ImImPy
	G30 β) 5'-W G C C G C W-3'	ImPy- β -ImPy- γ -Im- β -ImImPy
	G31 β) 5'-W G C C C G W-3'	ImPy- β -PyIm- γ -PyImImImPy

10-11-2017 10:11:11 AM

TABLE 60: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	481 β) 5'-W C G T T T W-3'	PyImHpHpHp- γ -PyPy- β -PyIm
5	481 β p) 5'-W C G T T T W-3'	PyIm- β -HpHp- γ -PyPy- β -PyIm
	482 β) 5'-W C G T T A W-3'	PyImHpHpPy- γ -HpPy- β -PyIm
	482 β p) 5'-W C G T T A W-3'	PyIm- β -HpPy- γ -HpPy- β -PyIm
	483 β) 5'-W C G T T G W-3'	PyIm- β -HpIm- γ -PyPy- β -PyIm
	484 β) 5'-W C G T T C W-3'	PyImHpHpPy- γ -ImPy- β -PyIm
10	484 β p) 5'-W C G T T C W-3'	PyIm- β -HpPy- γ -ImPy- β -PyIm
	485 β) 5'-W C G T A T W-3'	PyImHpPyHp- γ -PyHp- β -PyIm
	485 β p) 5'-W C G T A T W-3'	PyIm- β -PyHp- γ -PyHp- β -PyIm
	486 β) 5'-W C G T A A W-3'	PyImHpPyPy- γ -HpHp- β -PyIm
	486 β p) 5'-W C G T A A W-3'	PyIm- β -PyPy- γ -HpHp- β -PyIm
15	487 β) 5'-W C G T A G W-3'	PyIm- β -PyIm- γ -PyHp- β -PyIm
	488 β) 5'-W C G T A C W-3'	PyImHpPyPy- γ -ImHp- β -PyIm
	488 β p) 5'-W C G T A C W-3'	PyIm- β -PyPy- γ -ImHp- β -PyIm
	489 β) 5'-W C G T G T W-3'	PyIm- β -ImHp- γ -PyPy- β -PyIm
	490 β) 5'-W C G T G A W-3'	PyIm- β -ImPy- γ -HpPy- β -PyIm
20	491 β) 5'-W C G T G G W-3'	PyIm- β -ImIm- γ -PyPy- β -PyIm
	492 β) 5'-W C G T G C W-3'	PyIm- β -ImPy- γ -ImPy- β -PyIm
	493 β) 5'-W C G T C T W-3'	PyImHpPyHp- γ -PyIm- β -PyIm
	493 β p) 5'-W C G T C T W-3'	PyIm- β -PyHp- γ -PyIm- β -PyIm
	494 β) 5'-W C G T C A W-3'	PyImHpPyPy- γ -HpIm- β -PyIm
25	494 β p) 5'-W C G T C A W-3'	PyIm- β -PyPy- γ -HpIm- β -PyIm
	495 β) 5'-W C G T C G W-3'	PyIm- β -PyIm- γ -PyIm- β -PyIm
	496 β) 5'-W C G T C C W-3'	PyImHpPyPy- γ -ImIm- β -PyIm
	496 β p) 5'-W C G T C C W-3'	PyIm- β -PyPy- γ -ImIm- β -PyIm
	497 β) 5'-W C G A T T W-3'	PyImPyHpHp- γ -PyPy- β -PyIm
30	497 β p) 5'-W C G A T T W-3'	PyIm- β -HpHp- γ -PyPy- β -PyIm
	498 β) 5'-W C G A T A W-3'	PyImPyHpPy- γ -HpPy- β -PyIm
	498 β p) 5'-W C G A T A W-3'	PyIm- β -HpPy- γ -HpPy- β -PyIm

TABLE 60 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	499 β) 5'-W C G A T G W-3'	PyIm- β -HpIm- γ -PyPy- β -PyIm
	500 β) 5'-W C G A T C W-3'	PyImPyHpPy- γ -ImPy- β -PyIm
5	500 β p) 5'-W C G A T C W-3'	PyIm- β -HpPy- γ -ImPy- β -PyIm
	501 β) 5'-W C G A A T W-3'	PyImPyPyHp- γ -PyHp- β -PyIm
	501 β p) 5'-W C G A A T W-3'	PyIm- β -PyHp- γ -PyHp- β -PyIm
	502 β) 5'-W C G A A A W-3'	PyImPyPyPy- γ -HpHp- β -PyIm
	502 β p) 5'-W C G A A A W-3'	PyIm- β -PyPy- γ -HpHp- β -PyIm
10	503 β) 5'-W C G A A G W-3'	PyIm- β -PyIm- γ -PyHp- β -PyIm
	504 β) 5'-W C G A A C W-3'	PyImPyPyPy- γ -ImHp- β -PyIm
	504 β p) 5'-W C G A A C W-3'	PyIm- β -PyPy- γ -ImHp- β -PyIm
	505 β) 5'-W C G A G T W-3'	PyIm- β -ImHp- γ -PyPy- β -PyIm
	506 β) 5'-W C G A G A W-3'	PyIm- β -ImPy- γ -HpPy- β -PyIm
	507 β) 5'-W C G A G G W-3'	PyIm- β -ImIm- γ -PyPy- β -PyIm
	508 β) 5'-W C G A G C W-3'	PyIm- β -ImPy- γ -ImPy- β -PyIm
	509 β) 5'-W C G A C T W-3'	PyImPyPyHp- γ -PyIm- β -PyIm
	509 β p) 5'-W C G A C T W-3'	PyIm- β -PyHp- γ -PyIm- β -PyIm
	510 β) 5'-W C G A C A W-3'	PyImPyPyPy- γ -HpIm- β -PyIm
20	510 β p) 5'-W C G A C A W-3'	PyIm- β -PyPy- γ -HpIm- β -PyIm
	511 β) 5'-W C G A C G W-3'	PyIm- β -PyIm- γ -PyIm- β -PyIm
	512 β) 5'-W C G A C C W-3'	PyImPyPyPy- γ -ImIm- β -PyIm
	512 β p) 5'-W C G A C C W-3'	PyIm- β -PyPy- γ -ImIm- β -PyIm

TABLE 61: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	513 β) 5'-W C G G T T W-3'	PyImIm- β -Hp- γ -PyPy- β -PyIm
5	514 β) 5'-W C G G T A W-3'	PyImIm- β -Py- γ -HpPy- β -PyIm
	515 β) 5'-W C G G T G W-3'	PyImIm- β -Im- γ -PyPy- β -PyIm
	516 β) 5'-W C G G T C W-3'	PyImIm- β -Py- γ -ImPy- β -PyIm
	517 β) 5'-W C G G A T W-3'	PyImIm- β -Hp- γ -PyHp- β -PyIm
	518 β) 5'-W C G G A A W-3'	PyImIm- β -Py- γ -HpHp- β -PyIm
10	519 β) 5'-W C G G A G W-3'	PyImIm- β -Im- γ -PyHp- β -PyIm
	520 β) 5'-W C G G A C W-3'	PyImIm- β -Py- γ -ImHp- β -PyIm
	521 β) 5'-W C G G G T W-3'	PyImImImHp- γ -PyPy- β -PyIm
	522 β) 5'-W C G G G A W-3'	PyImImImPy- γ -HpPy- β -PyIm
	523 β) 5'-W C G G C T W-3'	PyImIm- β -Hp- γ -PyIm- β -PyIm
	524 β) 5'-W C G G C A W-3'	PyImIm- β -Py- γ -HpIm- β -PyIm
	525 β) 5'-W C G C T T W-3'	PyImPyHpHp- γ -Py- β -ImPyIm
	525 β p) 5'-W C G C T T W-3'	PyImPy- β -Hp- γ -Py- β -ImPyIm
	526 β) 5'-W C G C T A W-3'	PyImPyHpPy- γ -Hp- β -ImPyIm
	526 β p) 5'-W C G C T A W-3'	PyImPy- β -Py- γ -Hp- β -ImPyIm
20	527 β) 5'-W C G C T G W-3'	PyIm- β -HpIm- γ -Py- β -ImPyIm
	528 β) 5'-W C G C T C W-3'	PyImPyHpPy- γ -Im- β -ImPyIm
	528 β p) 5'-W C G C T C W-3'	PyImPy- β -Py- γ -Im- β -ImPyIm
	529 β) 5'-W C G C A T W-3'	PyImPyPyHp- γ -Py- β -ImPyIm
	529 β p) 5'-W C G C A T W-3'	PyImPy- β -Hp- γ -Py- β -ImPyIm
25	530 β) 5'-W C G C A A W-3'	PyImPyPyPy- γ -Hp- β -ImPyIm
	530 β p) 5'-W C G C A A W-3'	PyImPy- β -Py- γ -Hp- β -ImPyIm
	531 β) 5'-W C G C A G W-3'	PyIm- β -PyIm- γ -Py- β -ImPyIm
	532 β) 5'-W C G C A C W-3'	PyImPyPyPy- γ -Im- β -ImPyIm
	532 β p) 5'-W C G C A C W-3'	PyImPy- β -Py- γ -Im- β -ImPyIm
30	533 β) 5'-W C G C G T W-3'	PyIm- β -ImHp- γ -Py- β -ImPyIm
	534 β) 5'-W C G C G A W-3'	PyIm- β -ImPy- γ -Hp- β -ImPyIm

TABLE 61 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	535 β) 5'-W C G C C T W-3'	PyImPyPyHp- γ -PyImIm- β -Im
5	536 β) 5'-W C G C C A W-3'	PyImPyPyPy- γ -HpImIm- β -Im
	G33 β) 5'-W C G G G G W-3'	PyImImImIm- γ -PyPy- β -PyIm
	G34 β) 5'-W C G G G C W-3'	PyImImImPy- γ -ImPy- β -PyIm
	G35 β) 5'-W C G G C G W-3'	PyImIm- β -Im- γ -PyIm- β -PyIm
	G36 β) 5'-W C G G C C W-3'	PyImIm- β -Py- γ -ImIm- β -PyIm
10	G37 β) 5'-W C G C G G W-3'	PyIm- β -ImIm- γ -Py- β -ImPyIm
	G38 β) 5'-W C G C G C W-3'	PyIm- β -ImPy- γ -Im- β -ImPyIm
	G39 β) 5'-W C G C C G W-3'	PyIm- β -PyIm- γ -PyImIm- β -Im
	G40 β) 5'-W C G C C C W-3'	PyImPyPyPy- γ -ImImIm- β -Im

15

TABLE 62: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	537 β) 5'-W C T T T T W-3'	PyHpHpHpHp- γ -PyPy- β -PyIm
5	537 β p) 5'-W C T T T T W-3'	PyHp- β -HpHp- γ -PyPy- β -PyIm
	538 β) 5'-W C T T T A W-3'	PyHpHpHpPy- γ -HpPy- β -PyIm
	538 β p) 5'-W C T T T A W-3'	PyHp- β -HpPy- γ -HpPy- β -PyIm
	539 β) 5'-W C T T T G W-3'	PyHp- β -HpIm- γ -PyPy- β -PyIm
	540 β) 5'-W C T T T C W-3'	PyHpHpHpPy- γ -ImPy- β -PyIm
10	540 β p) 5'-W C T T T C W-3'	PyHp- β -HpPy- γ -ImPy- β -PyIm
	541 β) 5'-W C T T A T W-3'	PyHpHpPyHp- γ -PyHp- β -PyIm
	541 β p) 5'-W C T T A T W-3'	PyHp- β -PyHp- γ -PyHp- β -PyIm
	542 β) 5'-W C T T A A W-3'	PyHpHpPyPy- γ -HpHp- β -PyIm
	542 β p) 5'-W C T T A A W-3'	PyHp- β -PyPy- γ -HpHp- β -PyIm
	543 β) 5'-W C T T A G W-3'	PyHp- β -PyIm- γ -PyHp- β -PyIm
	544 β) 5'-W C T T A C W-3'	PyHpHpPyPy- γ -ImHp- β -PyIm
	544 β p) 5'-W C T T A C W-3'	PyHp- β -PyPy- γ -ImHp- β -PyIm
	545 β) 5'-W C T T G T W-3'	PyHp- β -ImHp- γ -PyPy- β -PyIm
	546 β) 5'-W C T T G A W-3'	PyHp- β -ImPy- γ -HpPy- β -PyIm
20	547 β) 5'-W C T T G G W-3'	PyHp- β -ImIm- γ -PyPy- β -PyIm
	548 β) 5'-W C T T G C W-3'	PyHp- β -ImPy- γ -ImPy- β -PyIm
	549 β) 5'-W C T T C T W-3'	PyHpHpPyHp- γ -PyIm- β -PyIm
	549 β p) 5'-W C T T C T W-3'	PyHp- β -PyHp- γ -PyIm- β -PyIm
	550 β) 5'-W C T T C A W-3'	PyHpHpPyPy- γ -HpIm- β -PyIm
25	550 β p) 5'-W C T T C A W-3'	PyHp- β -PyPy- γ -HpIm- β -PyIm
	551 β) 5'-W C T T C G W-3'	PyHp- β -PyIm- γ -PyIm- β -PyIm
	552 β) 5'-W C T T C C W-3'	PyHpHpPyPy- γ -ImIm- β -PyIm
	552 β p) 5'-W C T T C C W-3'	PyHp- β -PyPy- γ -ImIm- β -PyIm
	553 β) 5'-W C T A T T W-3'	PyHpPyHpHp- γ -PyPy- β -PyIm
30	553 β p) 5'-W C T A T T W-3'	PyHp- β -HpHp- γ -PyPy- β -PyIm
	554 β) 5'-W C T A T A W-3'	PyHpPyHpPy- γ -HpPy- β -PyIm

TABLE 62 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	554 β p) 5'-W C T A T A W-3'	PyHp- β -HpPy- γ -HpPy- β -PyIm
5	555 β) 5'-W C T A T G W-3'	PyHp- β -HpIm- γ -PyPy- β -PyIm
	556 β) 5'-W C T A T C W-3'	PyHpPyHpPy- γ -ImPy- β -PyIm
	556 β p) 5'-W C T A T C W-3'	PyHp- β -HpPy- γ -ImPy- β -PyIm
	557 β) 5'-W C T A A T W-3'	PyHpPyPyHp- γ -PyHp- β -PyIm
	557 β p) 5'-W C T A A T W-3'	PyHp- β -PyHp- γ -PyHp- β -PyIm
10	558 β) 5'-W C T A A A W-3'	PyHpPyPyPy- γ -HpHp- β -PyIm
	558 β p) 5'-W C T A A A W-3'	PyHp- β -PyPy- γ -HpHp- β -PyIm
	559 β) 5'-W C T A A G W-3'	PyHp- β -PyIm- γ -PyHp- β -PyIm
	560 β) 5'-W C T A A C W-3'	PyHpPyPyPy- γ -ImHp- β -PyIm
	560 β p) 5'-W C T A A C W-3'	PyHp- β -PyPy- γ -ImHp- β -PyIm
15	561 β) 5'-W C T A G T W-3'	PyHp- β -ImHp- γ -PyPy- β -PyIm
	562 β) 5'-W C T A G A W-3'	PyHp- β -ImPy- γ -HpPy- β -PyIm
	563 β) 5'-W C T A G G W-3'	PyHp- β -ImIm- γ -PyPy- β -PyIm
	564 β) 5'-W C T A G C W-3'	PyHp- β -ImPy- γ -ImPy- β -PyIm
	565 β) 5'-W C T A C T W-3'	PyHpPyPyHp- γ -PyIm- β -PyIm
20	565 β p) 5'-W C T A C T W-3'	PyHp- β -PyHp- γ -PyIm- β -PyIm
	566 β) 5'-W C T A C A W-3'	PyHpPyPyPy- γ -HpIm- β -PyIm
	566 β p) 5'-W C T A C A W-3'	PyHp- β -PyPy- γ -HpIm- β -PyIm
	567 β) 5'-W C T A C G W-3'	PyHp- β -PyIm- γ -PyIm- β -PyIm
	568 β) 5'-W C T A C C W-3'	PyHpPyPyPy- γ -ImIm- β -PyIm
25	568 β p) 5'-W C T A C C W-3'	PyHp- β -PyPy- γ -ImIm- β -PyIm

TABLE 63: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTSNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	569 β) 5'-W C T G T T W-3'	Py- β -ImHpHp- γ -PyPy- β -PyIm
5	570 β) 5'-W C T G T A W-3'	Py- β -ImHpPy- γ -HpPy- β -PyIm
	571 β) 5'-W C T G T G W-3'	Py- β -ImHpIm- γ -PyPy- β -PyIm
	572 β) 5'-W C T G T C W-3'	Py- β -ImHpPy- γ -ImPy- β -PyIm
	573 β) 5'-W C T G A T W-3'	Py- β -ImPyHp- γ -PyHp- β -PyIm
	574 β) 5'-W C T G A A W-3'	Py- β -ImPyPy- γ -HpHp- β -PyIm
10	575 β) 5'-W C T G A G W-3'	Py- β -ImPyIm- γ -PyHp- β -PyIm
	576 β) 5'-W C T G A C W-3'	Py- β -ImPyPy- γ -ImHp- β -PyIm
	577 β) 5'-W C T G G T W-3'	Py- β -ImImHp- γ -PyPy- β -PyIm
	578 β) 5'-W C T G G A W-3'	Py- β -ImImPy- γ -HpPy- β -PyIm
	579 β) 5'-W C T G C T W-3'	Py- β -ImPyHp- γ -PyIm- β -PyIm
	580 β) 5'-W C T G C A W-3'	Py- β -ImPyPy- γ -HpIm- β -PyIm
	581 β) 5'-W C T G G G W-3'	Py- β -ImImIm- γ -PyPy- β -PyIm
	582 β) 5'-W C T G G C W-3'	Py- β -ImImPy- γ -ImPy- β -PyIm
	583 β) 5'-W C T G C G W-3'	Py- β -ImPyIm- γ -PyIm- β -PyIm
	584 β) 5'-W C T G C C W-3'	Py- β -ImPyPy- γ -ImIm- β -PyIm
20	585 β) 5'-W C T C T T W-3'	PyHpPyHpHp- γ -Py- β -ImPyIm
	585 β p) 5'-W C T C T T W-3'	PyHpPy- β -Hp- γ -Py- β -ImPyIm
	586 β) 5'-W C T C T A W-3'	PyHpPyHpPy- γ -Hp- β -ImPyIm
	586 β p) 5'-W C T C T A W-3'	PyHpPy- β -Py- γ -Hp- β -ImPyIm
	587 β) 5'-W C T C T G W-3'	PyHp- β -HpIm- γ -Py- β -ImPyIm
25	588 β) 5'-W C T C T C W-3'	PyHpPyHpPy- γ -Im- β -ImPyIm
	588 β p) 5'-W C T C T C W-3'	PyHpPy- β -Py- γ -Im- β -ImPyIm
	589 β) 5'-W C T C A T W-3'	PyHpPyPyHp- γ -Py- β -ImPyIm
	589 β p) 5'-W C T C A T W-3'	PyHpPy- β -Hp- γ -Py- β -ImPyIm
	590 β) 5'-W C T C A A W-3'	PyHpPyPyPy- γ -Hp- β -ImPyIm
30	590 β p) 5'-W C T C A A W-3'	PyHpPy- β -Py- γ -Hp- β -ImPyIm
	591 β) 5'-W C T C A G W-3'	PyHp- β -PyIm- γ -Py- β -ImPyIm

TABLE 63 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCTSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	592 β) 5'-W C T C A C W-3'	PyHpPyPyPy- γ -Im- β -ImPyIm
	592 β p) 5'-W C T C A C W-3'	PyHpPy- β -Py- γ -Im- β -ImPyIm
5	593 β) 5'-W C T C G T W-3'	PyHp- β -ImHp- γ -Py- β -ImPyIm
	594 β) 5'-W C T C G A W-3'	PyHp- β -ImPy- γ -Hp- β -ImPyIm
	595 β) 5'-W C T C C T W-3'	PyHpPyPyHp- γ -PyImIm- β -Im
	595 β p) 5'-W C T C C T W-3'	Py- β -PyPyHp- γ -PyImIm- β -Im
	596 β) 5'-W C T C C A W-3'	PyHpPyPyPy- γ -HpImIm- β -Im
10	596 β p) 5'-W C T C C A W-3'	Py- β -PyPyPy- γ -HpImIm- β -Im
	597 β) 5'-W C T C G G W-3'	PyHp- β -ImIm- γ -Py- β -ImPyIm
	598 β) 5'-W C T C G C W-3'	PyHp- β -ImPy- γ -Im- β -ImPyIm
	599 β) 5'-W C T C C G W-3'	PyHp- β -PyIm- γ -PyImIm- β -Im
	600 β) 5'-W C T C C C W-3'	PyHpPyPyPy- γ -ImImIm- β -Im
	600 β p) 5'-W C T C C C W-3'	Py- β -PyPyPy- γ -ImImIm- β -Im

TABLE 64: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCAWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	601 β) 5'-W C A T T T W-3'	PyPyHpHpHp- γ -PyPy- β -HpIm
5	601 β p) 5'-W C A T T T W-3'	PyPy- β -HpHp- γ -PyPy- β -HpIm
	602 β) 5'-W C A T T A W-3'	PyPyHpHpPy- γ -HpPy- β -HpIm
	602 β p) 5'-W C A T T A W-3'	PyPy- β -HpPy- γ -HpPy- β -HpIm
	603 β) 5'-W C A T T G W-3'	PyPy- β -HpIm- γ -PyPy- β -HpIm
	604 β) 5'-W C A T T C W-3'	PyPyHpHpPy- γ -ImPy- β -HpIm
10	604 β p) 5'-W C A T T C W-3'	PyPy- β -HpPy- γ -ImPy- β -HpIm
	605 β) 5'-W C A T A T W-3'	PyPyHpPyHp- γ -PyHp- β -HpIm
	605 β p) 5'-W C A T A T W-3'	PyPy- β -PyHp- γ -PyHp- β -HpIm
	606 β) 5'-W C A T A A W-3'	PyPyHpPyPy- γ -HpHp- β -HpIm
	606 β p) 5'-W C A T A A W-3'	PyPy- β -PyPy- γ -HpHp- β -HpIm
	607 β) 5'-W C A T A G W-3'	PyPy- β -PyIm- γ -PyHp- β -HpIm
	608 β) 5'-W C A T A C W-3'	PyPyHpPyPy- γ -ImHp- β -HpIm
	608 β p) 5'-W C A T A C W-3'	PyPy- β -PyPy- γ -ImHp- β -HpIm
	609 β) 5'-W C A T G T W-3'	PyPy- β -ImHp- γ -PyPy- β -HpIm
	610 β) 5'-W C A T G A W-3'	PyPy- β -ImPy- γ -HpPy- β -HpIm
20	611 β) 5'-W C A T G G W-3'	PyPy- β -ImIm- γ -PyPy- β -HpIm
	612 β) 5'-W C A T G C W-3'	PyPy- β -ImPy- γ -ImPy- β -HpIm
	613 β) 5'-W C A T C T W-3'	PyPyHpPyHp- γ -PyIm- β -HpIm
	613 β p) 5'-W C A T C T W-3'	PyPy- β -PyHp- γ -PyIm- β -HpIm
	614 β) 5'-W C A T C A W-3'	PyPyHpPyPy- γ -HpIm- β -HpIm
25	614 β p) 5'-W C A T C A W-3'	PyPy- β -PyPy- γ -HpIm- β -HpIm
	615 β) 5'-W C A T C G W-3'	PyPy- β -PyIm- γ -PyIm- β -HpIm
	616 β) 5'-W C A T C C W-3'	PyPyHpPyPy- γ -ImIm- β -HpIm
	616 β p) 5'-W C A T C C W-3'	PyPy- β -PyPy- γ -ImIm- β -HpIm
	617 β) 5'-W C A A T T W-3'	PyPyPyHpHp- γ -PyPy- β -HpIm
30	617 β p) 5'-W C A A T T W-3'	PyPy- β -HpHp- γ -PyPy- β -HpIm
	618 β) 5'-W C A A T A W-3'	PyPyPyHpPy- γ -HpPy- β -HpIm
	618 β p) 5'-W C A A T A W-3'	PyPy- β -HpPy- γ -HpPy- β -HpIm

TABLE 64 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCAWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	619 β) 5'-W C A A T G W-3'	PyPy- β -HpIm- γ -PyPy- β -HpIm
	620 β) 5'-W C A A T C W-3'	PyPyPyHpPy- γ -ImPy- β -HpIm
5	620 β p) 5'-W C A A T C W-3'	PyPy- β -HpPy- γ -ImPy- β -HpIm
	621 β) 5'-W C A A A T W-3'	PyPyPyPyHp- γ -PyHp- β -HpIm
	621 β p) 5'-W C A A A T W-3'	PyPy- β -PyHp- γ -PyHp- β -HpIm
	622 β) 5'-W C A A A A W-3'	PyPyPyPyPy- γ -HpHp- β -HpIm
	622 β p) 5'-W C A A A A W-3'	PyPy- β -PyPy- γ -HpHp- β -HpIm
10	623 β) 5'-W C A A A G W-3'	PyPy- β -PyIm- γ -PyHp- β -HpIm
	624 β) 5'-W C A A A C W-3'	PyPyPyPyPy- γ -ImHp- β -HpIm
	624 β p) 5'-W C A A A C W-3'	PyPy- β -PyPy- γ -ImHp- β -HpIm
	625 β) 5'-W C A A G T W-3'	PyPy- β -ImHp- γ -PyPy- β -HpIm
	626 β) 5'-W C A A G A W-3'	PyPy- β -ImPy- γ -HpPy- β -HpIm
	627 β) 5'-W C A A G G W-3'	PyPy- β -ImIm- γ -PyPy- β -HpIm
	628 β) 5'-W C A A G C W-3'	PyPy- β -ImPy- γ -ImPy- β -HpIm
	629 β) 5'-W C A A C T W-3'	PyPyPyPyHp- γ -PyIm- β -HpIm
	629 β p) 5'-W C A A C T W-3'	PyPy- β -PyHp- γ -PyIm- β -HpIm
	630 β) 5'-W C A A C A W-3'	PyPyPyPyPy- γ -HpIm- β -HpIm
	630 β p) 5'-W C A A C A W-3'	PyPy- β -PyPy- γ -HpIm- β -HpIm
	631 β) 5'-W C A A C G W-3'	PyPy- β -PyIm- γ -PyIm- β -HpIm
	632 β) 5'-W C A A C C W-3'	PyPyPyPyPy- γ -ImIm- β -HpIm
	632 β p) 5'-W C A A C C W-3'	PyPy- β -PyPy- γ -ImIm- β -HpIm

TABLE 65: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	633 β) 5'-W C A G T T W-3'	Py- β -ImHpHp- γ -PyPy- β -HpIm
5	634 β) 5'-W C A G T A W-3'	Py- β -ImHpPy- γ -HpPy- β -HpIm
	635 β) 5'-W C A G T G W-3'	Py- β -ImHpIm- γ -PyPy- β -HpIm
	636 β) 5'-W C A G T C W-3'	Py- β -ImHpPy- γ -ImPy- β -HpIm
	637 β) 5'-W C A G A T W-3'	Py- β -ImPyHp- γ -PyHp- β -HpIm
	638 β) 5'-W C A G A A W-3'	Py- β -ImPyPy- γ -HpHp- β -HpIm
10	639 β) 5'-W C A G A G W-3'	Py- β -ImPyIm- γ -PyHp- β -HpIm
	640 β) 5'-W C A G A C W-3'	Py- β -ImPyPy- γ -ImHp- β -HpIm
	641 β) 5'-W C A G G T W-3'	Py- β -ImImHp- γ -PyPy- β -HpIm
	642 β) 5'-W C A G G A W-3'	Py- β -ImImPy- γ -HpPy- β -HpIm
	643 β) 5'-W C A G C T W-3'	Py- β -ImPyHp- γ -PyIm- β -HpIm
	644 β) 5'-W C A G C A W-3'	Py- β -ImPyPy- γ -HpIm- β -HpIm
	645 β) 5'-W C A G G G W-3'	Py- β -ImImIm- γ -PyPy- β -HpIm
	646 β) 5'-W C A G G C W-3'	Py- β -ImImPy- γ -ImPy- β -HpIm
	647 β) 5'-W C A G C G W-3'	Py- β -ImPyIm- γ -PyIm- β -HpIm
	648 β) 5'-W C A G C C W-3'	Py- β -ImPyPy- γ -ImIm- β -HpIm
20	649 β) 5'-W C A C T T W-3'	PyPyPyHpHp- γ -Py- β -ImHpIm
	649 β p) 5'-W C A C T T W-3'	PyPyPy- β -Hp- γ -Py- β -ImHpIm
	650 β) 5'-W C A C T A W-3'	PyPyPyHpPy- γ -Hp- β -ImHpIm
	650 β p) 5'-W C A C T A W-3'	PyPyPy- β -Py- γ -Hp- β -ImHpIm
	651 β) 5'-W C A C T G W-3'	PyPy- β -HpIm- γ -Py- β -ImHpIm
25	652 β) 5'-W C A C T C W-3'	PyPyPyHpPy- γ -Im- β -ImHpIm
	652 β p) 5'-W C A C T C W-3'	PyPyPy- β -Py- γ -Im- β -ImHpIm
	653 β) 5'-W C A C A T W-3'	PyPyPyPyHp- γ -Py- β -ImHpIm
	653 β p) 5'-W C A C A T W-3'	PyPyPy- β -Hp- γ -Py- β -ImHpIm
	654 β) 5'-W C A C A A W-3'	PyPyPyPyPy- γ -Hp- β -ImHpIm
30	654 β p) 5'-W C A C A A W-3'	PyPyPy- β -Py- γ -Hp- β -ImHpIm
	655 β) 5'-W C A C A G W-3'	PyPy- β -PyIm- γ -Py- β -ImHpIm

TABLE 65 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	656 β) 5'-W C A C A C W-3'	PyPyPyPyPyPy- γ -Im- β -ImHpIm
	656 β p) 5'-W C A C A C W-3'	PyPyPy- β -Py- γ -Im- β -ImHpIm
5	657 β) 5'-W C A C G T W-3'	PyPy- β -ImHp- γ -Py- β -ImHpIm
	658 β p) 5'-W C A C G A W-3'	PyPy- β -ImPy- γ -Hp- β -ImHpIm
	659 β) 5'-W C A C C T W-3'	PyPyPyPyPyHp- γ -PyImIm- β -Im
	659 β p) 5'-W C A C C T W-3'	Py- β -PyPyHp- γ -PyImIm- β -Im
	660 β) 5'-W C A C C A W-3'	PyPyPyPyPyPy- γ -HpImIm- β -Im
10	660 β p) 5'-W C A C C A W-3'	Py- β -PyPyPyPy- γ -HpImIm- β -Im
	661 β) 5'-W C A C G G W-3'	PyPy- β -ImIm- γ -Py- β -ImHpIm
	662 β) 5'-W C A C G C W-3'	PyPy- β -ImPy- γ -Im- β -ImHpIm
	663 β) 5'-W C A C C G W-3'	PyPy- β -PyIm- γ -PyImIm- β -Im
	664 β) 5'-W C A C C C W-3'	PyPyPyPyPyPy- γ -ImImIm- β -Im
	664 β p) 5'-W C A C C C W-3'	Py- β -PyPyPyPy- γ -ImImIm- β -Im

TABLE 66: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	665 β) 5'-W C C T T T W-3'	PyPyHpHpHp- γ -PyPy- β -ImIm
5	665 β p) 5'-W C C T T T W-3'	PyPy- β -HpHp- γ -PyPy- β -ImIm
	666 β) 5'-W C C T T A W-3'	PyPyHpHpPy- γ -HpPy- β -ImIm
	666 β p) 5'-W C C T T A W-3'	PyPy- β -HpPy- γ -HpPy- β -ImIm
	667 β) 5'-W C C T T G W-3'	PyPy- β -HpIm- γ -PyPy- β -ImIm
	668 β) 5'-W C C T T C W-3'	PyPyHpHpPy- γ -ImPy- β -ImIm
10	668 β p) 5'-W C C T T C W-3'	PyPy- β -HpPy- γ -ImPy- β -ImIm
	669 β) 5'-W C C T A T W-3'	PyPyHpPyHp- γ -PyHp- β -ImIm
	669 β p) 5'-W C C T A T W-3'	PyPy- β -PyHp- γ -PyHp- β -ImIm
	670 β) 5'-W C C T A A W-3'	PyPyHpPyPy- γ -HpHp- β -ImIm
	670 β p) 5'-W C C T A A W-3'	PyPy- β -PyPy- γ -HpHp- β -ImIm
	671 β) 5'-W C C T A G W-3'	PyPy- β -PyIm- γ -PyHp- β -ImIm
	672 β) 5'-W C C T A C W-3'	PyPyHpPyPy- γ -ImHp- β -ImIm
	672 β p) 5'-W C C T A C W-3'	PyPy- β -PyPy- γ -ImHp- β -ImIm
	673 β) 5'-W C C T G T W-3'	PyPy- β -ImHp- γ -PyPy- β -ImIm
	674 β) 5'-W C C T G A W-3'	PyPy- β -ImPy- γ -HpPy- β -ImIm
20	675 β) 5'-W C C T G G W-3'	PyPy- β -ImIm- γ -PyPy- β -ImIm
	676 β) 5'-W C C T G C W-3'	PyPy- β -ImPy- γ -ImPy- β -ImIm
	677 β) 5'-W C C T C T W-3'	PyPyHpPyHp- γ -PyIm- β -ImIm
	677 β p) 5'-W C C T C T W-3'	PyPy- β -PyHp- γ -PyIm- β -ImIm
	678 β) 5'-W C C T C A W-3'	PyPyHpPyPy- γ -HpIm- β -ImIm
25	678 β p) 5'-W C C T C A W-3'	PyPy- β -PyPy- γ -HpIm- β -ImIm
	679 β) 5'-W C C T C G W-3'	PyPy- β -PyIm- γ -PyIm- β -ImIm
	680 β) 5'-W C C T C C W-3'	PyPyHpPyPy- γ -ImIm- β -ImIm
	680 β p) 5'-W C C T C C W-3'	PyPy- β -PyPy- γ -ImIm- β -ImIm
	681 β) 5'-W C C A T T W-3'	PyPyPyHpHp- γ -PyPy- β -ImIm
30	681 β p) 5'-W C C A T T W-3'	PyPy- β -HpHp- γ -PyPy- β -ImIm
	682 β) 5'-W C C A T A W-3'	PyPyPyHpPy- γ -HpPy- β -ImIm
	682 β p) 5'-W C C A T A W-3'	PyPy- β -HpPy- γ -HpPy- β -ImIm

TABLE 66: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	683 β) 5'-W C C A T G W-3'	PyPy- β -HpIm- γ -PyPy- β -ImIm
5	684 β) 5'-W C C A T C W-3'	PyPyPyHpPy- γ -ImPy- β -ImIm
	684 β p) 5'-W C C A T C W-3'	PyPy- β -HpPy- γ -ImPy- β -ImIm
	685 β) 5'-W C C A A T W-3'	PyPyPyPyHp- γ -PyHp- β -ImIm
	685 β p) 5'-W C C A A T W-3'	PyPy- β -PyHp- γ -PyHp- β -ImIm
	686 β) 5'-W C C A A A W-3'	PyPyPyPyPy- γ -HpHp- β -ImIm
10	686 β p) 5'-W C C A A A W-3'	PyPy- β -PyPy- γ -HpHp- β -ImIm
	687 β) 5'-W C C A A G W-3'	PyPy- β -PyIm- γ -PyHp- β -ImIm
	688 β) 5'-W C C A A C W-3'	PyPyPyPyPy- γ -ImHp- β -ImIm
	688 β p) 5'-W C C A A C W-3'	PyPy- β -PyPy- γ -ImHp- β -ImIm
	689 β) 5'-W C C A G T W-3'	PyPy- β -ImHp- γ -PyPy- β -ImIm
	690 β) 5'-W C C A G A W-3'	PyPy- β -ImPy- γ -HpPy- β -ImIm
	691 β) 5'-W C C A G G W-3'	PyPy- β -ImIm- γ -PyPy- β -ImIm
	692 β) 5'-W C C A G C W-3'	PyPy- β -ImPy- γ -ImPy- β -ImIm
	693 β) 5'-W C C A C T W-3'	PyPyPyPyHp- γ -PyIm- β -ImIm
	693 β p) 5'-W C C A C T W-3'	PyPy- β -PyHp- γ -PyIm- β -ImIm
20	694 β) 5'-W C C A C A W-3'	PyPyPyPyPy- γ -HpIm- β -ImIm
	694 β p) 5'-W C C A C A W-3'	PyPy- β -PyPy- γ -HpIm- β -ImIm
	695 β) 5'-W C C A C G W-3'	PyPy- β -PyIm- γ -PyIm- β -ImIm
	696 β) 5'-W C C A C C W-3'	PyPyPyPyPy- γ -ImIm- β -ImIm
	696 β p) 5'-W C C A C C W-3'	PyPy- β -PyPy- γ -ImIm- β -ImIm

TABLE 67: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	697 β) 5'-W C C G T T W-3'	Py- β -ImHpHp- γ -PyPy- β -ImIm
5	698 β) 5'-W C C G T A W-3'	Py- β -ImHpPy- γ -HpPy- β -ImIm
	699 β) 5'-W C C G T G W-3'	Py- β -ImHpIm- γ -PyPy- β -ImIm
	700 β) 5'-W C C G T C W-3'	Py- β -ImHpPy- γ -ImPy- β -ImIm
	701 β) 5'-W C C G A T W-3'	Py- β -ImPyHp- γ -PyHp- β -ImIm
	702 β) 5'-W C C G A A W-3'	Py- β -ImPyPy- γ -HpHp- β -ImIm
10	703 β) 5'-W C C G A G W-3'	Py- β -ImPyIm- γ -PyHp- β -ImIm
	704 β) 5'-W C C G A C W-3'	Py- β -ImPyPy- γ -ImHp- β -ImIm
	705 β) 5'-W C C G G T W-3'	Py- β -ImImHp- γ -PyPy- β -ImIm
	706 β) 5'-W C C G G A W-3'	Py- β -ImImPy- γ -HpPy- β -ImIm
	707 β) 5'-W C C G C T W-3'	Py- β -ImPyHp- γ -PyIm- β -ImIm
	708 β) 5'-W C C G C A W-3'	Py- β -ImPyPy- γ -HpIm- β -ImIm
	709 β) 5'-W C C C T T W-3'	PyPyPyHpHp- γ -Py- β -ImImIm
	709 β p) 5'-W C C C T T W-3'	PyPyPy- β -Hp- γ -Py- β -ImImIm
	710 β) 5'-W C C C T A W-3'	PyPyPyHpPy- γ -Hp- β -ImImIm
	710 β p) 5'-W C C C T A W-3'	PyPyPy- β -Py- γ -Hp- β -ImImIm
20	711 β) 5'-W C C C T G W-3'	PyPy- β -HpIm- γ -Py- β -ImImIm
	712 β) 5'-W C C C T C W-3'	PyPyPyHpPy- γ -Im- β -ImImIm
	712 β p) 5'-W C C C T C W-3'	PyPyPy- β -Py- γ -Im- β -ImImIm
	713 β) 5'-W C C C A T W-3'	PyPyPyPyHp- γ -Py- β -ImImIm
	713 β p) 5'-W C C C A T W-3'	PyPyPy- β -Hp- γ -Py- β -ImImIm
25	714 β) 5'-W C C C A A W-3'	PyPyPyPyPy- γ -Hp- β -ImImIm
	714 β p) 5'-W C C C A A W-3'	PyPyPy- β -Py- γ -Hp- β -ImImIm
	715 β) 5'-W C C C A G W-3'	PyPy- β -PyIm- γ -Py- β -ImImIm
	716 β) 5'-W C C C A C W-3'	PyPyPyPyPy- γ -Im- β -ImImIm
	716 β p) 5'-W C C C A C W-3'	PyPyPy- β -Py- γ -Im- β -ImImIm
30	717 β) 5'-W C C C G T W-3'	PyPy- β -ImHp- γ -Py- β -ImImIm
	718 β) 5'-W C C C G A W-3'	PyPy- β -ImPy- γ -Hp- β -ImImIm

TABLE 67 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WCCSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	G41 β) 5'-W C C G G G W-3'	Py- β -ImImIm- γ -PyPy- β -ImIm
	G42 β) 5'-W C C G G C W-3'	Py- β -ImImPy- γ -ImPy- β -ImIm
5	G43 β) 5'-W C C G C G W-3'	Py- β -ImPyIm- γ -PyIm- β -ImIm
	G44 β) 5'-W C C G C C W-3'	Py- β -ImPyPy- γ -ImIm- β -ImIm
	G45 β) 5'-W C C C G G W-3'	PyPy- β -ImIm- γ -Py- β -ImImIm
	G46 β) 5'-W C C C G C W-3'	PyPy- β -ImPy- γ -Im- β -ImImIm
	G47 β) 5'-W C C C C G W-3'	PyPy- β -PyIm- γ -PyImImImIm

TABLE 68: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAGWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	723 β) 5'-W A G T T G W-3'	PyIm- β -HpIm- γ -PyPyPyPyHp
5	723 β p) 5'-W A G T T G W-3'	PyIm- β -HpIm- γ -PyPy- β -PyHp
	727 β) 5'-W A G T A G W-3'	PyIm- β -PyIm- γ -PyHpPyPyHp
	727 β p) 5'-W A G T A G W-3'	PyIm- β -PyIm- γ -PyHp- β -PyHp
	729 β) 5'-W A G T G T W-3'	PyIm- β -ImHp- γ -PyPyPyPyHp
	729 β p) 5'-W A G T G T W-3'	PyIm- β -ImHp- γ -PyPy- β -PyHp
10	730 β) 5'-W A G T G A W-3'	PyIm- β -ImPy- γ -HpPyPyPyHp
	730 β p) 5'-W A G T G A W-3'	PyIm- β -ImPy- γ -HpPy- β -PyHp
	731 β) 5'-W A G T G G W-3'	PyIm- β -ImIm- γ -PyPyPyPyHp
	731 β p) 5'-W A G T G G W-3'	PyIm- β -ImIm- γ -PyPy- β -PyHp
	732 β) 5'-W A G T G C W-3'	PyIm- β -ImPy- γ -ImPyPyPyHp
	732 β p) 5'-W A G T G C W-3'	PyIm- β -ImPy- γ -ImPy- β -PyHp
	735 β) 5'-W A G T C G W-3'	PyIm- β -PyIm- γ -PyImPyPyHp
	735 β p) 5'-W A G T C G W-3'	PyIm- β -PyIm- γ -PyIm- β -PyHp
	739 β) 5'-W A G A T G W-3'	PyIm- β -HpIm- γ -PyPyHpPyHp
	739 β p) 5'-W A G A T G W-3'	PyIm- β -HpIm- γ -PyPy- β -PyHp
	743 β) 5'-W A G A A G W-3'	PyIm- β -PyIm- γ -PyHpHpPyHp
	743 β p) 5'-W A G A A G W-3'	PyIm- β -PyIm- γ -PyHp- β -PyHp
	745 β) 5'-W A G A G T W-3'	PyIm- β -ImHp- γ -PyPyHpPyHp
	745 β p) 5'-W A G A G T W-3'	PyIm- β -ImHp- γ -PyPy- β -PyHp
	746 β) 5'-W A G A G A W-3'	PyIm- β -ImPy- γ -HpPyHpPyHp
25	746 β p) 5'-W A G A G A W-3'	PyIm- β -ImPy- γ -HpPy- β -PyHp
	747 β) 5'-W A G A G G W-3'	PyIm- β -ImIm- γ -PyPyHpPyHp
	747 β p) 5'-W A G A G G W-3'	PyIm- β -ImIm- γ -PyPy- β -PyHp
	748 β) 5'-W A G A G C W-3'	PyIm- β -ImPy- γ -ImPyHpPyHp
	748 β p) 5'-W A G A G C W-3'	PyIm- β -ImPy- γ -ImPy- β -PyHp
30	751 β) 5'-W A G A C G W-3'	PyIm- β -PyIm- γ -PyImHpPyHp
	751 β p) 5'-W A G A C G W-3'	PyIm- β -PyIm- γ -PyIm- β -PyHp

TABLE 69: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	753 β) 5'-W A G G T T W-3'	PyImIm- β -Hp- γ -PyPyPyPyHp
5	753 β p) 5'-W A G G T T W-3'	PyImIm- β -Hp- γ -Py- β -PyPyHp
	754 β) 5'-W A G G T A W-3'	PyImIm- β -Py- γ -HpPyPyPyHp
	754 β p) 5'-W A G G T A W-3'	PyImIm- β -Py- γ -Hp- β -PyPyHp
	755 β) 5'-W A G G T G W-3'	PyImIm- β -Im- γ -PyPyPyPyHp
	755 β p) 5'-W A G G T G W-3'	PyImIm- β -Im- γ -Py- β -PyPyHp
10	756 β) 5'-W A G G T C W-3'	PyImIm- β -Py- γ -ImPyPyPyHp
	756 β p) 5'-W A G G T C W-3'	PyImIm- β -Py- γ -Im- β -PyPyHp
	757 β) 5'-W A G G A T W-3'	PyImIm- β -Hp- γ -PyHpPyPyHp
	757 β p) 5'-W A G G A T W-3'	PyImIm- β -Hp- γ -Py- β -PyPyHp
	758 β) 5'-W A G G A A W-3'	PyImIm- β -Py- γ -HpHpPyPyHp
	758 β p) 5'-W A G G A A W-3'	PyImIm- β -Py- γ -Hp- β -PyPyHp
	759 β) 5'-W A G G A G W-3'	PyImIm- β -Im- γ -PyHpPyPyHp
	759 β p) 5'-W A G G A G W-3'	PyImIm- β -Im- γ -Py- β -PyPyHp
	760 β) 5'-W A G G A C W-3'	PyImIm- β -Py- γ -ImHpPyPyHp
	760 β p) 5'-W A G G A C W-3'	PyImIm- β -Py- γ -Im- β -PyPyHp
20	763 β) 5'-W A G G C T W-3'	PyImIm- β -Hp- γ -PyImPyPyHp
	764 β) 5'-W A G G C A W-3'	PyImIm- β -Py- γ -HpImPyPyHp
	765 β) 5'-W A G C T T W-3'	PyImPyHpHp- γ -Py- β -ImPyHp
	765 β p) 5'-W A G C T T W-3'	PyImPy- β -Hp- γ -Py- β -ImPyHp
	766 β) 5'-W A G C T A W-3'	PyImPyHpPy- γ -Hp- β -ImPyHp
25	766 β p) 5'-W A G C T A W-3'	PyImPy- β -Py- γ -Hp- β -ImPyHp
	767 β) 5'-W A G C T G W-3'	PyIm- β -HpIm- γ -Py- β -ImPyHp
	768 β) 5'-W A G C T C W-3'	PyImPyHpPy- γ -Im- β -ImPyHp
	768 β p) 5'-W A G C T C W-3'	PyImPy- β -Py- γ -Im- β -ImPyHp
	769 β) 5'-W A G C A T W-3'	PyImPyPyHp- γ -Py- β -ImPyHp
30	769 β p) 5'-W A G C A T W-3'	PyImPy- β -Hp- γ -Py- β -ImPyHp
	770 β) 5'-W A G C A A W-3'	PyImPyPyPy- γ -Hp- β -ImPyHp

TABLE 69 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	770 β p) 5'-W A G C A A W-3'	PyImPy- β -Py- γ -Hp- β -ImPyHp
5	771 β) 5'-W A G C A G W-3'	PyIm- β -PyIm- γ -Py- β -ImPyHp
	772 β) 5'-W A G C A C W-3'	PyImPyPyPy- γ -Im- β -ImPyHp
	772 β p) 5'-W A G C A C W-3'	PyImPy- β -Py- γ -Im- β -ImPyHp
	773 β) 5'-W A G C G T W-3'	PyIm- β -ImHp- γ -Py- β -ImPyHp
	774 β) 5'-W A G C G A W-3'	PyIm- β -ImPy- γ -Hp- β -ImPyHp
10	775 β) 5'-W A G C C T W-3'	PyImPyPyHp- γ -PyImIm- β -Hp
	776 β) 5'-W A G C C A W-3'	PyImPyPyPy- γ -HpImIm- β -Hp
	779 β) 5'-W A G G C G W-3'	PyImIm- β -Im- γ -PyImPyPyHp
	780 β) 5'-W A G G C C W-3'	PyImIm- β -Py- γ -ImImPyPyHp
	781 β) 5'-W A G C G G W-3'	PyIm- β -ImIm- γ -Py- β -ImPyHp
	782 β) 5'-W A G C G C W-3'	PyIm- β -ImPy- γ -Im- β -ImPyHp
	783 β) 5'-W A G C C G W-3'	PyIm- β -PyIm- γ -PyImIm- β -Hp
	784 β) 5'-W A G C C C W-3'	PyImPyPyPy- γ -ImImIm- β -Hp

TABLE 70: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WATWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	787 β) 5'-W A T T T G W-3'	PyHp- β -HpIm- γ -PyPyPyPyHp
5	787 β p) 5'-W A T T T G W-3'	PyHp- β -HpIm- γ -PyPy- β -PyHp
	791 β) 5'-W A T T A G W-3'	PyHp- β -PyIm- γ -PyHpPyPyHp
	791 β p) 5'-W A T T A G W-3'	PyHp- β -PyIm- γ -PyHp- β -PyHp
	793 β) 5'-W A T T G T W-3'	PyHp- β -ImHp- γ -PyPyPyPyHp
	793 β p) 5'-W A T T G T W-3'	PyHp- β -ImHp- γ -PyPy- β -PyHp
10	794 β) 5'-W A T T G A W-3'	PyHp- β -ImPy- γ -HpPyPyPyHp
	794 β p) 5'-W A T T G A W-3'	PyHp- β -ImPy- γ -HpPy- β -PyHp
	795 β) 5'-W A T T G G W-3'	PyHp- β -ImIm- γ -PyPyPyPyHp
	795 β p) 5'-W A T T G G W-3'	PyHp- β -ImPy- γ -ImPyPyPyHp
	796 β p) 5'-W A T T G C W-3'	PyHp- β -ImPy- γ -ImPy- β -PyHp
15	799 β) 5'-W A T T C G W-3'	PyHp- β -PyIm- γ -PyImPyPyHp
	799 β p) 5'-W A T T C G W-3'	PyHp- β -PyIm- γ -PyIm- β -PyHp
	803 β) 5'-W A T A T G W-3'	PyHp- β -HpIm- γ -PyPyHpPyHp
	803 β p) 5'-W A T A T G W-3'	PyHp- β -HpIm- γ -PyPy- β -PyHp
	807 β) 5'-W A T A A G W-3'	PyHp- β -PyIm- γ -PyHpHpPyHp
20	807 β p) 5'-W A T A A G W-3'	PyHp- β -PyIm- γ -PyHp- β -PyHp
	809 β) 5'-W A T A G T W-3'	PyHp- β -ImHp- γ -PyPyHpPyHp
	809 β p) 5'-W A T A G T W-3'	PyHp- β -ImHp- γ -PyPy- β -PyHp
	810 β) 5'-W A T A G A W-3'	PyHp- β -ImPy- γ -HpPyHpPyHp
	810 β p) 5'-W A T A G A W-3'	PyHp- β -ImPy- γ -HpPy- β -PyHp
25	811 β) 5'-W A T A G G W-3'	PyHp- β -ImIm- γ -PyPyHpPyHp
	811 β p) 5'-W A T A G G W-3'	PyHp- β -ImIm- γ -PyPy- β -PyHp
	812 β) 5'-W A T A G C W-3'	PyHp- β -ImPy- γ -ImPyHpPyHp
	812 β p) 5'-W A T A G C W-3'	PyHp- β -ImPy- γ -ImPy- β -PyHp
	815 β) 5'-W A T A C G W-3'	PyHp- β -PyIm- γ -PyImHpPyHp
30	815 β p) 5'-W A T A C G W-3'	PyHp- β -PyIm- γ -PyIm- β -PyHp

TABLE 71: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WATSNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	817 β) 5'-W A T G T T W-3'	Py- β -ImHpHp- γ -PyPyPyPyHp
5	817 β p) 5'-W A T G T T W-3'	Py- β -ImHpHp- γ -PyPyPy- β -Hp
	818 β) 5'-W A T G T A W-3'	Py- β -ImHpPy- γ -HpPyPyPyHp
	818 β p) 5'-W A T G T A W-3'	Py- β -ImHpPy- γ -HpPyPy- β -Hp
	819 β) 5'-W A T G T G W-3'	Py- β -ImHpIm- γ -PyPyPyPyHp
	819 β p) 5'-W A T G T G W-3'	Py- β -ImHpIm- γ -PyPyPy- β -Hp
10	820 β) 5'-W A T G T C W-3'	Py- β -ImHpPy- γ -ImPyPyPyHp
	820 β p) 5'-W A T G T C W-3'	Py- β -ImHpPy- γ -ImPyPy- β -Hp
	821 β) 5'-W A T G A T W-3'	Py- β -ImPyHp- γ -PyHpPyPyHp
	821 β p) 5'-W A T G A T W-3'	Py- β -ImPyHp- γ -PyHpPy- β -Hp
	822 β) 5'-W A T G A A W-3'	Py- β -ImPyPy- γ -HpHpPyPyHp
	822 β p) 5'-W A T G A A W-3'	Py- β -ImPyPy- γ -HpHpPy- β -Hp
	823 β) 5'-W A T G A G W-3'	Py- β -ImPyIm- γ -PyHpPyPyHp
	823 β p) 5'-W A T G A G W-3'	Py- β -ImPyIm- γ -PyHpPy- β -Hp
	824 β) 5'-W A T G A C W-3'	Py- β -ImPyPy- γ -ImHpPyPyHp
	824 β p) 5'-W A T G A C W-3'	Py- β -ImPyPy- γ -ImHpPy- β -Hp
20	825 β) 5'-W A T G G T W-3'	Py- β -ImImHp- γ -PyPyPyPyHp
	825 β p) 5'-W A T G G T W-3'	Py- β -ImImHp- γ -PyPyPy- β -Hp
	826 β) 5'-W A T G G A W-3'	Py- β -ImImPy- γ -HpPyPyPyHp
	826 β p) 5'-W A T G G A W-3'	Py- β -ImImPy- γ -HpPyPy- β -Hp
	827 β) 5'-W A T G C T W-3'	Py- β -ImPyHp- γ -PyImPyPyHp
25	827 β p) 5'-W A T G C T W-3'	Py- β -ImPyHp- γ -PyImPy- β -Hp
	828 β) 5'-W A T G C A W-3'	Py- β -ImPyPy- γ -HpImPyPyHp
	828 β p) 5'-W A T G C A W-3'	Py- β -ImPyPy- γ -HpImPy- β -Hp
	829 β) 5'-W A T G G G W-3'	Py- β -ImImIm- γ -PyPyPyPyHp
	829 β p) 5'-W A T G G G W-3'	Py- β -ImImIm- γ -PyPyPy- β -Hp
30	830 β) 5'-W A T G G C W-3'	Py- β -ImImPy- γ -ImPyPyPyHp
	830 β p) 5'-W A T G G C W-3'	Py- β -ImImPy- γ -ImPyPy- β -Hp
	831 β) 5'-W A T G C G W-3'	Py- β -ImPyIm- γ -PyImPyPyHp
	831 β p) 5'-W A T G C G W-3'	Py- β -ImPyIm- γ -PyImPy- β -Hp

TABLE 71: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WATSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	832 β) 5'-W A T G C C W-3'	Py- β -ImPyPy- γ -ImImPyPyHp
5	832 β p) 5'-W A T G C C W-3'	Py- β -ImPyPy- γ -ImImPy- β -Hp
	833 β) 5'-W A T C T T W-3'	PyHpPyHpHp- γ -Py- β -ImPyHp
	833 β p) 5'-W A T C T T W-3'	PyHpPy- β -Hp- γ -Py- β -ImPyHp
	834 β) 5'-W A T C T A W-3'	PyHpPyHpPy- γ -Hp- β -ImPyHp
	834 β p) 5'-W A T C T A W-3'	PyHpPy- β -Py- γ -Hp- β -ImPyHp
10	835 β) 5'-W A T C T G W-3'	PyHp- β -HpIm- γ -Py- β -ImPyHp
	836 β) 5'-W A T C T C W-3'	PyHpPyHpPy- γ -Im- β -ImPyHp
	836 β p) 5'-W A T C T C W-3'	PyHpPy- β -Py- γ -Im- β -ImPyHp
	837 β) 5'-W A T C A T W-3'	PyHpPyPyHp- γ -Py- β -ImPyHp
	837 β p) 5'-W A T C A T W-3'	PyHpPy- β -Hp- γ -Py- β -ImPyHp
	838 β) 5'-W A T C A A W-3'	PyHpPyPyPy- γ -Hp- β -ImPyHp
	838 β p) 5'-W A T C A A W-3'	PyHpPy- β -Py- γ -Hp- β -ImPyHp
	839 β) 5'-W A T C A G W-3'	PyHp- β -PyIm- γ -Py- β -ImPyHp
	840 β) 5'-W A T C A C W-3'	PyHpPyPyPy- γ -Im- β -ImPyHp
	840 β p) 5'-W A T C A C W-3'	PyHpPy- β -Py- γ -Im- β -ImPyHp
20	841 β) 5'-W A T C G T W-3'	PyHp- β -ImHp- γ -Py- β -ImPyHp
	842 β) 5'-W A T C G A W-3'	PyHp- β -ImPy- γ -Hp- β -ImPyHp
	843 β) 5'-W A T C C T W-3'	PyHpPyPyHp- γ -PyImIm- β -Hp
	843 β p) 5'-W A T C C T W-3'	Py- β -PyPyHp- γ -PyImIm- β -Hp
	844 β) 5'-W A T C C A W-3'	PyHpPyPyPy- γ -HpImIm- β -Hp
25	844 β p) 5'-W A T C C A W-3'	Py- β -PyPyPy- γ -HpImIm- β -Hp
	845 β) 5'-W A T C G G W-3'	PyHp- β -ImIm- γ -Py- β -ImPyHp
	846 β) 5'-W A T C G C W-3'	PyHp- β -ImPy- γ -Im- β -ImPyHp
	847 β) 5'-W A T C C G W-3'	PyHp- β -PyIm- γ -PyImIm- β -Hp
	848 β) 5'-W A T C C C W-3'	PyHpPyPyPy- γ -ImImIm- β -Hp
30	848 β p) 5'-W A T C C C W-3'	Py- β -PyPyPy- γ -ImImIm- β -Hp

TABLE 72: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAAWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	851 β) 5'-W A A T T G W-3'	PyPy- β -HpIm- γ -PyPyPyHpHp
5	851 β p) 5'-W A A T T G W-3'	PyPy- β -HpIm- γ -PyPy- β -HpHp
	855 β) 5'-W A A T A G W-3'	PyPy- β -PyIm- γ -PyHpPyHpHp
	855 β p) 5'-W A A T A G W-3'	PyPy- β -PyIm- γ -PyHp- β -HpHp
	857 β) 5'-W A A T G T W-3'	PyPy- β -ImHp- γ -PyPyPyHpHp
	857 β p) 5'-W A A T G T W-3'	PyPy- β -ImHp- γ -PyPy- β -HpHp
10	858 β) 5'-W A A T G A W-3'	PyPy- β -ImPy- γ -HpPyPyHpHp
	858 β p) 5'-W A A T G A W-3'	PyPy- β -ImPy- γ -HpPy- β -HpHp
	859 β) 5'-W A A T G G W-3'	PyPy- β -ImIm- γ -PyPyPyHpHp
	859 β p) 5'-W A A T G G W-3'	PyPy- β -ImIm- γ -PyPy- β -HpHp
	860 β) 5'-W A A T G C W-3'	PyPy- β -ImPy- γ -ImPyPyHpHp
	860 β p) 5'-W A A T G C W-3'	PyPy- β -ImPy- γ -ImPy- β -HpHp
	863 β) 5'-W A A T C G W-3'	PyPy- β -PyIm- γ -PyImPyHpHp
	863 β p) 5'-W A A T C G W-3'	PyPy- β -PyIm- γ -PyIm- β -HpHp
	867 β) 5'-W A A A T G W-3'	PyPy- β -HpIm- γ -PyPyHpHpHp
	867 β p) 5'-W A A A T G W-3'	PyPy- β -HpIm- γ -PyPy- β -HpHp
20	871 β) 5'-W A A A A G W-3'	PyPy- β -PyIm- γ -PyHpHpHpHp
	871 β p) 5'-W A A A A G W-3'	PyPy- β -PyIm- γ -PyHp- β -HpHp
	873 β) 5'-W A A A G T W-3'	PyPy- β -ImHp- γ -PyPyHpHpHp
	873 β p) 5'-W A A A G T W-3'	PyPy- β -ImHp- γ -PyPy- β -HpHp
	874 β) 5'-W A A A G A W-3'	PyPy- β -ImPy- γ -HpPyHpHpHp
25	874 β p) 5'-W A A A G A W-3'	PyPy- β -ImPy- γ -HpPy- β -HpHp
	875 β) 5'-W A A A G G W-3'	PyPy- β -ImIm- γ -PyPyHpHpHp
	875 β p) 5'-W A A A G G W-3'	PyPy- β -ImIm- γ -PyPy- β -HpHp
	876 β) 5'-W A A A G C W-3'	PyPy- β -ImPy- γ -ImPyHpHpHp
	876 β p) 5'-W A A A G C W-3'	PyPy- β -ImPy- γ -ImPy- β -HpHp
30	879 β) 5'-W A A A C G W-3'	PyPy- β -PyIm- γ -PyImHpHpHp
	879 β p) 5'-W A A A C G W-3'	PyPy- β -PyIm- γ -PyIm- β -HpHp

TABLE 73: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	881 β) 5'-W A A G T T W-3'	Py- β -ImHpHp- γ -PyPyPyHpHp
5	881 β p) 5'-W A A G T T W-3'	Py- β -ImHpHp- γ -PyPyPy- β -Hp
	882 β) 5'-W A A G T A W-3'	Py- β -ImHpPy- γ -HpPyPyHpHp
	882 β p) 5'-W A A G T A W-3'	Py- β -ImHpPy- γ -HpPyPy- β -Hp
	883 β) 5'-W A A G T G W-3'	Py- β -ImHpIm- γ -PyPyPyHpHp
	883 β p) 5'-W A A G T G W-3'	Py- β -ImHpIm- γ -PyPyPy- β -Hp
10	884 β) 5'-W A A G T C W-3'	Py- β -ImHpPy- γ -ImPyPyHpHp
	884 β p) 5'-W A A G T C W-3'	Py- β -ImHpPy- γ -ImPyPy- β -Hp
	885 β) 5'-W A A G A T W-3'	Py- β -ImPyHp- γ -PyHpPyHpHp
	885 β p) 5'-W A A G A T W-3'	Py- β -ImPyHp- γ -PyHpPy- β -Hp
	886 β) 5'-W A A G A A W-3'	Py- β -ImPyPy- γ -HpHpPyHpHp
	886 β p) 5'-W A A G A A W-3'	Py- β -ImPyPy- γ -HpHpPy- β -Hp
	887 β) 5'-W A A G A G W-3'	Py- β -ImPyIm- γ -PyHpPyHpHp
	887 β p) 5'-W A A G A G W-3'	Py- β -ImPyIm- γ -PyHpPy- β -Hp
	888 β) 5'-W A A G A C W-3'	Py- β -ImPyPy- γ -ImHpPyHpHp
	888 β p) 5'-W A A G A C W-3'	Py- β -ImPyPy- γ -ImHpPy- β -Hp
20	889 β) 5'-W A A G G T W-3'	Py- β -ImImHp- γ -PyPyPyHpHp
	889 β p) 5'-W A A G G T W-3'	Py- β -ImImHp- γ -PyPyPy- β -Hp
	890 β) 5'-W A A G G A W-3'	Py- β -ImImPy- γ -HpPyPyHpHp
	890 β p) 5'-W A A G G A W-3'	Py- β -ImImPy- γ -HpPyPy- β -Hp
	891 β) 5'-W A A G C T W-3'	Py- β -ImPyHp- γ -PyImPyHpHp
25	891 β p) 5'-W A A G C T W-3'	Py- β -ImPyHp- γ -PyImPy- β -Hp
	892 β) 5'-W A A G C A W-3'	Py- β -ImPyPy- γ -HpImPyHpHp
	892 β p) 5'-W A A G C A W-3'	Py- β -ImPyPy- γ -HpImPy- β -Hp
	893 β) 5'-W A A G G G W-3'	Py- β -ImImIm- γ -PyPyPyHpHp
	893 β p) 5'-W A A G G G W-3'	Py- β -ImImIm- γ -PyPyPy- β -Hp
30	894 β) 5'-W A A G G C W-3'	Py- β -ImImPy- γ -ImPyPyHpHp
	894 β p) 5'-W A A G G C W-3'	Py- β -ImImPy- γ -ImPyPy- β -Hp
	895 β) 5'-W A A G C G W-3'	Py- β -ImPyIm- γ -PyImPyHpHp
	895 β p) 5'-W A A G C G W-3'	Py- β -ImPyIm- γ -PyImPy- β -Hp

TABLE 73 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WAASNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	896 β) 5'-W A A G C C W-3'	Py- β -ImPyPy- γ -ImImPyHpHp
	896 β p) 5'-W A A G C C W-3'	Py- β -ImPyPy- γ -ImImPy- β -Hp
5	897 β) 5'-W A A C T T W-3'	PyPyPyHpHp- γ -Py- β -ImHpHp
	897 β p) 5'-W A A C T T W-3'	PyPyPy- β -Hp- γ -Py- β -ImHpHp
	898 β) 5'-W A A C T A W-3'	PyPyPyHpPy- γ -Hp- β -ImHpHp
	898 β p) 5'-W A A C T A W-3'	PyPyPy- β -Py- γ -Hp- β -ImHpHp
	899 β) 5'-W A A C T G W-3'	PyPy- β -HpIm- γ -Py- β -ImHpHp
10	900 β) 5'-W A A C T C W-3'	PyPyPyHpPy- γ -Im- β -ImHpHp
	900 β p) 5'-W A A C T C W-3'	PyPyPy- β -Py- γ -Im- β -ImHpHp
	901 β) 5'-W A A C A T W-3'	PyPyPyPyHp- γ -Py- β -ImHpHp
	901 β p) 5'-W A A C A T W-3'	PyPyPy- β -Hp- γ -Py- β -ImHpHp
	902 β) 5'-W A A C A A W-3'	PyPyPyPyPy- γ -Hp- β -ImHpHp
	902 β p) 5'-W A A C A A W-3'	PyPyPy- β -Py- γ -Hp- β -ImHpHp
	903 β) 5'-W A A C A G W-3'	PyPy- β -PyIm- γ -Py- β -ImHpHp
	904 β) 5'-W A A C A C W-3'	PyPyPyPyPy- γ -Im- β -ImHpHp
	904 β p) 5'-W A A C A C W-3'	PyPyPy- β -Py- γ -Im- β -ImHpHp
	905 β) 5'-W A A C G T W-3'	PyPy- β -ImHp- γ -Py- β -ImHpHp
20	906 β) 5'-W A A C G A W-3'	PyPy- β -ImPy- γ -Hp- β -ImHpHp
	907 β) 5'-W A A C C T W-3'	PyPyPyPyHp- γ -PyImIm- β -Hp
	907 β p) 5'-W A A C C T W-3'	Py- β -PyPyHp- γ -PyImIm- β -Hp
	908 β) 5'-W A A C C A W-3'	PyPyPyPyPy- γ -HpImIm- β -Hp
	908 β p) 5'-W A A C C A W-3'	Py- β -PyPyPy- γ -HpImIm- β -Hp
25	909 β) 5'-W A A C G G W-3'	PyPy- β -ImIm- γ -Py- β -ImHpHp
	910 β) 5'-W A A C G C W-3'	PyPy- β -ImPy- γ -Im- β -ImHpHp
	911 β) 5'-W A A C C G W-3'	PyPy- β -PyIm- γ -PyImIm- β -Hp
	912 β) 5'-W A A C C C W-3'	PyPyPyPyPy- γ -ImImIm- β -Hp
	912 β p) 5'-W A A C C C W-3'	Py- β -PyPyPy- γ -ImImIm- β -Hp

TABLE 74: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	913 β) 5'-W A C T T T W-3'	PyPyHpHpHp- γ -PyPy- β -ImHp
5	913 β p) 5'-W A C T T T W-3'	PyPy- β -HpHp- γ -PyPy- β -ImHp
	914 β) 5'-W A C T T A W-3'	PyPyHpHpPy- γ -HpPy- β -ImHp
	914 β p) 5'-W A C T T A W-3'	PyPy- β -HpPy- γ -HpPy- β -ImHp
	915 β) 5'-W A C T T G W-3'	PyPy- β -HpIm- γ -PyPy- β -ImHp
	916 β) 5'-W A C T T C W-3'	PyPyHpHpPy- γ -ImPy- β -ImHp
10	916 β p) 5'-W A C T T C W-3'	PyPy- β -HpPy- γ -ImPy- β -ImHp
	917 β) 5'-W A C T A T W-3'	PyPyHpPyHp- γ -PyHp- β -ImHp
	917 β p) 5'-W A C T A T W-3'	PyPyHpPyHp- γ -PyHp- β -ImHp
	918 β) 5'-W A C T A A W-3'	PyPyHpPyPy- γ -HpHp- β -ImHp
	918 β p) 5'-W A C T A A W-3'	PyPy- β -PyPy- γ -HpHp- β -ImHp
	919 β) 5'-W A C T A G W-3'	PyPy- β -PyIm- γ -PyHp- β -ImHp
	920 β) 5'-W A C T A C W-3'	PyPyHpPyPy- γ -ImHp- β -ImHp
	920 β p) 5'-W A C T A C W-3'	PyPy- β -PyPy- γ -ImHp- β -ImHp
	921 β) 5'-W A C T G T W-3'	PyPy- β -ImHp- γ -PyPy- β -ImHp
	922 β) 5'-W A C T G A W-3'	PyPy- β -ImPy- γ -HpPy- β -ImHp
20	923 β) 5'-W A C T G G W-3'	PyPy- β -ImIm- γ -PyPy- β -ImHp
	924 β) 5'-W A C T G C W-3'	PyPy- β -ImPy- γ -ImPy- β -ImHp
	925 β) 5'-W A C T C T W-3'	PyPyHpPyHp- γ -PyIm- β -ImHp
	925 β p) 5'-W A C T C T W-3'	PyPy- β -PyHp- γ -PyIm- β -ImHp
	926 β) 5'-W A C T C A W-3'	PyPyHpPyPy- γ -HpIm- β -ImHp
25	926 β p) 5'-W A C T C A W-3'	PyPy- β -PyPy- γ -HpIm- β -ImHp
	927 β) 5'-W A C T C G W-3'	PyPy- β -PyIm- γ -PyIm- β -ImHp
	928 β) 5'-W A C T C C W-3'	PyPyHpPyPy- γ -ImIm- β -ImHp
	928 β p) 5'-W A C T C C W-3'	PyPy- β -PyPy- γ -ImIm- β -ImHp
	929 β) 5'-W A C A T T W-3'	PyPyPyHpHp- γ -PyPy- β -ImHp
30	929 β p) 5'-W A C A T T W-3'	PyPy- β -HpHp- γ -PyPy- β -ImHp
	930 β) 5'-W A C A T A W-3'	PyPyPyHpPy- γ -HpPy- β -ImHp
	930 β p) 5'-W A C A T A W-3'	PyPy- β -HpPy- γ -HpPy- β -ImHp
	931 β) 5'-W A C A T G W-3'	PyPy- β -HpIm- γ -PyPy- β -ImHp

TABLE 74 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	932 β) 5'-W A C A T C W-3'	PyPyPyHpPy- γ -ImPy- β -ImHp
5	932 β p) 5'-W A C A T C W-3'	PyPy- β -HpPy- γ -ImPy- β -ImHp
	933 β) 5'-W A C A A T W-3'	PyPyPyPyHpPy- γ -PyHp- β -ImHp
	933 β p) 5'-W A C A A T W-3'	PyPy- β -PyHp- γ -PyHp- β -ImHp
	934 β) 5'-W A C A A A W-3'	PyPyPyPyPyPy- γ -HpHp- β -ImHp
	934 β p) 5'-W A C A A A W-3'	PyPy- β -PyPy- γ -HpHp- β -ImHp
10	935 β) 5'-W A C A A G W-3'	PyPy- β -PyIm- γ -PyHp- β -ImHp
	936 β) 5'-W A C A A C W-3'	PyPyPyPyPyPy- γ -ImHp- β -ImHp
	936 β p) 5'-W A C A A C W-3'	PyPy- β -PyPy- γ -ImHp- β -ImHp
	937 β) 5'-W A C A G T W-3'	PyPy- β -ImHp- γ -PyPy- β -ImHp
	938 β) 5'-W A C A G A W-3'	PyPy- β -ImPy- γ -HpPy- β -ImHp
15	939 β) 5'-W A C A G G W-3'	PyPy- β -ImIm- γ -PyPy- β -ImHp
	940 β) 5'-W A C A G C W-3'	PyPy- β -ImPy- γ -ImPy- β -ImHp
	941 β) 5'-W A C A C T W-3'	PyPyPyPyHpPy- γ -PyIm- β -ImHp
	941 β p) 5'-W A C A C T W-3'	PyPy- β -PyHp- γ -PyIm- β -ImHp
	942 β) 5'-W A C A C A W-3'	PyPyPyPyPyPy- γ -HpIm- β -ImHp
20	942 β p) 5'-W A C A C A W-3'	PyPy- β -PyPy- γ -HpIm- β -ImHp
	943 β) 5'-W A C A C G W-3'	PyPy- β -PyIm- γ -PyIm- β -ImHp
	944 β) 5'-W A C A C C W-3'	PyPyPyPyPyPy- γ -ImIm- β -ImHp
	944 β p) 5'-W A C A C C W-3'	PyPy- β -PyPy- γ -ImIm- β -ImHp

TABLE 75: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACSNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
5	945 β) 5'-W A C G T T W-3'	Py- β -ImHpHp- γ -PyPy- β -ImHp
	946 β) 5'-W A C G T A W-3'	Py- β -ImHpPy- γ -HpPy- β -ImHp
	947 β) 5'-W A C G T G W-3'	Py- β -ImHpIm- γ -PyPy- β -ImHp
	948 β) 5'-W A C G T C W-3'	Py- β -ImHpPy- γ -ImPy- β -ImHp
	949 β) 5'-W A C G A T W-3'	Py- β -ImPyHp- γ -PyHp- β -ImHp
10	950 β) 5'-W A C G A A W-3'	Py- β -ImPyPy- γ -HpHp- β -ImHp
	951 β) 5'-W A C G A G W-3'	Py- β -ImPyIm- γ -PyHp- β -ImHp
	952 β) 5'-W A C G A C W-3'	Py- β -ImPyPy- γ -ImHp- β -ImHp
	953 β) 5'-W A C G G T W-3'	Py- β -ImImHp- γ -PyPy- β -ImHp
	954 β) 5'-W A C G G A W-3'	Py- β -ImImPy- γ -HpPy- β -ImHp
15	955 β) 5'-W A C G C T W-3'	Py- β -ImPyHp- γ -PyIm- β -ImHp
	956 β) 5'-W A C G C A W-3'	Py- β -ImPyPy- γ -HpIm- β -ImHp
	957 β) 5'-W A C C T T W-3'	PyPyPyHpHp- γ -Py- β -ImImHp
	957 β_p) 5'-W A C C T T W-3'	PyPyPy- β -Hp- γ -Py- β -ImImHp
	958 β) 5'-W A C C T A W-3'	PyPyPyHpPy- γ -Hp- β -ImImHp
20	958 β_p) 5'-W A C C T A W-3'	PyPyPy- β -Py- γ -Hp- β -ImImHp
	959 β) 5'-W A C C T G W-3'	PyPy- β -HpIm- γ -Py- β -ImImHp
	960 β) 5'-W A C C T C W-3'	PyPyPyHpPy- γ -Im- β -ImImHp
	960 β_p) 5'-W A C C T C W-3'	PyPyPy- β -Py- γ -Im- β -ImImHp
	961 β) 5'-W A C C A T W-3'	PyPyPyPyHp- γ -Py- β -ImImHp
25	961 β_p) 5'-W A C C A T W-3'	PyPyPy- β -Hp- γ -Py- β -ImImHp
	962 β) 5'-W A C C A A W-3'	PyPyPyPyPy- γ -Hp- β -ImImHp
	962 β_p) 5'-W A C C A A W-3'	PyPyPy- β -Py- γ -Hp- β -ImImHp
	963 β) 5'-W A C C A G W-3'	PyPy- β -PyIm- γ -Py- β -ImImHp
	964 β) 5'-W A C C A C W-3'	PyPyPyPyPy- γ -Im- β -ImImHp
30	964 β_p) 5'-W A C C A C W-3'	PyPyPy- β -Py- γ -Im- β -ImImHp
	965 β) 5'-W A C C G T W-3'	PyPy- β -ImHp- γ -Py- β -ImImHp
	966 β) 5'-W A C C G A W-3'	PyPy- β -ImPy- γ -Hp- β -ImImHp
	969 β) 5'-W A C G G G W-3'	Py- β -ImImIm- γ -PyPy- β -ImHp
	970 β) 5'-W A C G G C W-3'	Py- β -ImImPy- γ -ImPy- β -ImHp

TABLE 75 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WACSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	971 β) 5' -W A C G C G W-3'	Py- β -ImPyIm- γ -PyIm- β -ImHp
	972 β) 5' -W A C G C C W-3'	Py- β -ImPyPy- γ -ImIm- β -ImHp
5	973 β) 5' -W A C C G G W-3'	PyPy- β -ImIm- γ -Py- β -ImImHp
	974 β) 5' -W A C C G C W-3'	PyPy- β -ImPy- γ -Im- β -ImImHp
	975 β) 5' -W A C C C G W-3'	PyPy- β -PyIm- γ -PyImImImHp

TABLE 76: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTGWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	979 β) 5'-W T G T T G W-3'	HpIm- β -HpIm- γ -PyPyPyPyPy
5	979 β p) 5'-W T G T T G W-3'	HpIm- β -HpIm- γ -PyPy- β -PyPy
	983 β) 5'-W T G T A G W-3'	HpIm- β -PyIm- γ -PyHpPyPyPy
	983 β p) 5'-W T G T A G W-3'	HpIm- β -PyIm- γ -PyHp- β -PyPy
	985 β) 5'-W T G T G T W-3'	HpIm- β -ImHp- γ -PyPyPyPyPy
	985 β p) 5'-W T G T G T W-3'	HpIm- β -ImHp- γ -PyPy- β -PyPy
10	986 β) 5'-W T G T G A W-3'	HpIm- β -ImPy- γ -HpPyPyPyPy
	986 β p) 5'-W T G T G A W-3'	HpIm- β -ImPy- γ -HpPy- β -PyPy
	987 β) 5'-W T G T G G W-3'	HpIm- β -ImIm- γ -PyPyPyPyPy
	987 β p) 5'-W T G T G G W-3'	HpIm- β -ImIm- γ -PyPy- β -PyPy
	988 β) 5'-W T G T G C W-3'	HpIm- β -ImPy- γ -ImPyPyPyPy
	988 β p) 5'-W T G T G C W-3'	HpIm- β -ImPy- γ -ImPy- β -PyPy
	991 β) 5'-W T G T C G W-3'	HpIm- β -PyIm- γ -PyImPyPyPy
	991 β p) 5'-W T G T C G W-3'	HpIm- β -PyIm- γ -PyIm- β -PyPy
	995 β) 5'-W T G A T G W-3'	HpIm- β -HpIm- γ -PyPyHpPyPy
	995 β p) 5'-W T G A T G W-3'	HpIm- β -HpIm- γ -PyPy- β -PyPy
20	999 β) 5'-W T G A A G W-3'	HpIm- β -PyIm- γ -PyHpHpPyPy
	999 β p) 5'-W T G A A G W-3'	HpIm- β -PyIm- γ -PyHp- β -PyPy
	1001 β) 5'-W T G A G T W-3'	HpIm- β -ImHp- γ -PyPyHpPyPy
	1001 β p) 5'-W T G A G T W-3'	HpIm- β -ImHp- γ -PyPy- β -PyPy
	1002 β) 5'-W T G A G A W-3'	HpIm- β -ImPy- γ -HpPyHpPyPy
25	1002 β p) 5'-W T G A G A W-3'	HpIm- β -ImPy- γ -HpPy- β -PyPy
	1003 β) 5'-W T G A G G W-3'	HpIm- β -ImIm- γ -PyPyHpPyPy
	1003 β p) 5'-W T G A G G W-3'	HpIm- β -ImIm- γ -PyPy- β -PyPy
	1004 β) 5'-W T G A G C W-3'	HpIm- β -ImPy- γ -ImPyHpPyPy
	1004 β p) 5'-W T G A G C W-3'	HpIm- β -ImPy- γ -ImPy- β -PyPy
30	1007 β) 5'-W T G A C G W-3'	HpIm- β -PyIm- γ -PyImHpPyPy
	1007 β p) 5'-W T G A C G W-3'	HpIm- β -PyIm- γ -PyIm- β -PyPy

TABLE 77: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	1009 β) 5'-W T G G T T W-3'	HpImIm- β -Hp- γ -PyPyPyPyPy
5	1009 β p) 5'-W T G G T T W-3'	HpImIm- β -Hp- γ -Py- β -PyPyPy
	1010 β) 5'-W T G G T A W-3'	HpImIm- β -Py- γ -HpPyPyPyPy
	1010 β p) 5'-W T G G T A W-3'	HpImIm- β -Py- γ -Hp- β -PyPyPy
	1011 β) 5'-W T G G T G W-3'	HpImIm- β -Im- γ -PyPyPyPyPy
	1011 β p) 5'-W T G G T G W-3'	HpImIm- β -Im- γ -Py- β -PyPyPy
10	1012 β) 5'-W T G G T C W-3'	HpImIm- β -Py- γ -ImPyPyPyPy
	1012 β p) 5'-W T G G T C W-3'	HpImIm- β -Py- γ -Im- β -PyPyPy
	1013 β) 5'-W T G G A T W-3'	HpImIm- β -Hp- γ -PyHpPyPyPy
	1013 β p) 5'-W T G G A T W-3'	HpImIm- β -Hp- γ -Py- β -PyPyPy
	1014 β) 5'-W T G G A A W-3'	HpImIm- β -Py- γ -HpHpPyPyPy
	1014 β p) 5'-W T G G A A W-3'	HpImIm- β -Py- γ -Hp- β -PyPyPy
	1015 β) 5'-W T G G A G W-3'	HpImIm- β -Im- γ -PyHpPyPyPy
	1015 β p) 5'-W T G G A G W-3'	HpImIm- β -Im- γ -Py- β -PyPyPy
	1016 β) 5'-W T G G A C W-3'	HpImIm- β -Py- γ -ImHpPyPyPy
	1016 β p) 5'-W T G G A C W-3'	HpImIm- β -Py- γ -Im- β -PyPyPy
	1019 β) 5'-W T G G C T W-3'	HpImIm- β -Hp- γ -PyImPyPyPy
	1020 β) 5'-W T G G C A W-3'	HpImIm- β -Py- γ -HpImPyPyPy
	1021 β) 5'-W T G C T T W-3'	HpImPyHpHp- γ -Py- β -ImPyPy
	1021 β p) 5'-W T G C T T W-3'	HpImPy- β -Hp- γ -Py- β -ImPyPy
	1022 β) 5'-W T G C T A W-3'	HpImPyHpPy- γ -Hp- β -ImPyPy
25	1022 β p) 5'-W T G C T A W-3'	HpImPy- β -Py- γ -Hp- β -ImPyPy
	1023 β) 5'-W T G C T G W-3'	HpIm- β -HpIm- γ -Py- β -ImPyPy
	1024 β) 5'-W T G C T C W-3'	HpImPyHpPy- γ -Im- β -ImPyPy
	1024 β p) 5'-W T G C T C W-3'	HpImPy- β -Py- γ -Im- β -ImPyPy
	1025 β) 5'-W T G C A T W-3'	HpImPyPyHp- γ -Py- β -ImPyPy
30	1025 β p) 5'-W T G C A T W-3'	HpImPy- β -Hp- γ -Py- β -ImPyPy
	1026 β) 5'-W T G C A A W-3'	HpImPyPyPy- γ -Hp- β -ImPyPy
	1026 β p) 5'-W T G C A A W-3'	HpImPy- β -Py- γ -Hp- β -ImPyPy
	1027 β) 5'-W T G C A G W-3'	HpIm- β -PyIm- γ -Py- β -ImPyPy

TABLE 77 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTGSNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	1028 β) 5'-W T G C A C W-3'	HpImPyPyPy- γ -Im- β -ImPyP
5	1028 β p) 5'-W T G C A C W-3'	HpImPy- β -Py- γ -Im- β -ImPyPy
	1029 β) 5'-W T G C G T W-3'	HpIm- β -ImHp- γ -Py- β -ImPyPy
	1030 β) 5'-W T G C G A W-3'	HpIm- β -ImPy- γ -Hp- β -ImPyPy
	1031 β) 5'-W T G C C T W-3'	HpImPyPyHp- γ -PyImIm- β -Py
	1031 β p) 5'-W T G C C T W-3'	HpImPy- β -Hp- γ -PyImIm- β -Py
10	1032 β) 5'-W T G C C A W-3'	HpImPyPyPy- γ -HpImIm- β -Py
	1032 β p) 5'-W T G C C A W-3'	HpImPy- β -Py- γ -HpImIm- β -Py
	1035 β) 5'-W T G G C G W-3'	HpImIm- β -Im- γ -PyImPyPyPy
	1036 β) 5'-W T G G C C W-3'	HpImIm- β -Py- γ -ImImPyPyPy
	1037 β) 5'-W T G C G G W-3'	HpIm- β -ImIm- γ -Py- β -ImPyPy
	1038 β) 5'-W T G C G C W-3'	HpIm- β -ImPy- γ -Im- β -ImPyPy
	1039 β) 5'-W T G C C G W-3'	HpIm- β -PyIm- γ -PyImIm- β -Py
	1040 β) 5'-W T G C C C W-3'	HpImPyPyPy- γ -ImImIm- β -Py

TABLE 78: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTTWNNW-3' with β substitutions.

	DNA sequence	aromatic amino acid sequence
	1043 β) 5'-W T T T T G W-3'	HpHp- β -HpIm- γ -PyPyPyPyPy
5	1043 β p) 5'-W T T T T G W-3'	HpHp- β -HpIm- γ -PyPy- β -PyPy
	1047 β) 5'-W T T T A G W-3'	HpHp- β -PyIm- γ -PyHpPyPyPy
	1047 β p) 5'-W T T T A G W-3'	HpHp- β -PyIm- γ -PyHp- β -PyPy
	1049 β) 5'-W T T T G T W-3'	HpHp- β -ImHp- γ -PyPyPyPyPy
	1049 β p) 5'-W T T T G T W-3'	HpHp- β -ImHp- γ -PyPy- β -PyPy
10	1050 β) 5'-W T T T G A W-3'	HpHp- β -ImPy- γ -HpPyPyPyPy
	1050 β p) 5'-W T T T G A W-3'	HpHp- β -ImPy- γ -HpPy- β -PyPy
	1051 β) 5'-W T T T G G W-3'	HpHp- β -ImIm- γ -PyPyPyPyPy
	1051 β p) 5'-W T T T G G W-3'	HpHp- β -ImIm- γ -PyPy- β -PyPy
	1052 β) 5'-W T T T G C W-3'	HpHp- β -ImPy- γ -ImPyPyPyPy
	1052 β p) 5'-W T T T G C W-3'	HpHp- β -ImPy- γ -ImPy- β -PyPy
	1055 β) 5'-W T T T C G W-3'	HpHp- β -PyIm- γ -PyImPyPyPy
	1055 β p) 5'-W T T T C G W-3'	HpHp- β -PyIm- γ -PyIm- β -PyPy
	1059 β) 5'-W T T A T G W-3'	HpHp- β -HpIm- γ -PyPyHpPyPy
	1059 β p) 5'-W T T A T G W-3'	HpHp- β -HpIm- γ -PyPy- β -PyPy
20	1063 β) 5'-W T T A A G W-3'	HpHp- β -PyIm- γ -PyHpHpPyPy
	1063 β p) 5'-W T T A A G W-3'	HpHp- β -PyIm- γ -PyHp- β -PyPy
	1065 β) 5'-W T T A G T W-3'	HpHp- β -ImHp- γ -PyPyHpPyPy
	1065 β p) 5'-W T T A G T W-3'	HpHp- β -ImHp- γ -PyPy- β -PyPy
	1066 β) 5'-W T T A G A W-3'	HpHp- β -ImPy- γ -HpPyHpPyPy
25	1066 β p) 5'-W T T A G A W-3'	HpHp- β -ImPy- γ -HpPy- β -PyPy
	1067 β) 5'-W T T A G G W-3'	HpHp- β -ImIm- γ -PyPyHpPyPy
	1067 β p) 5'-W T T A G G W-3'	HpHp- β -ImIm- γ -PyPy- β -PyPy
	1068 β) 5'-W T T A G C W-3'	HpHp- β -ImPy- γ -ImPyHpPyPy
	1068 β p) 5'-W T T A G C W-3'	HpHp- β -ImPy- γ -ImPy- β -PyPy
30	1071 β) 5'-W T T A C G W-3'	HpHp- β -PyIm- γ -PyImHpPyPy
	1071 β p) 5'-W T T A C G W-3'	HpHp- β -PyIm- γ -PyIm- β -PyPy

TABLE 79: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTTSNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1073 β) 5'-W T T G T T W-3'	Hp- β -ImHpHp- γ -PyPyPyPyPy
5	1073 β p) 5'-W T T G T T W-3'	Hp- β -ImHpHp- γ -PyPyPy- β -Py
	1074 β) 5'-W T T G T A W-3'	Hp- β -ImHpPy- γ -HpPyPyPyPy
	1074 β p) 5'-W T T G T A W-3'	Hp- β -ImHpPy- γ -HpPyPy- β -Py
	1075 β) 5'-W T T G T G W-3'	Hp- β -ImHpIm- γ -PyPyPyPyPy
	1075 β p) 5'-W T T G T G W-3'	Hp- β -ImHpIm- γ -PyPyPy- β -Py
10	1076 β) 5'-W T T G T C W-3'	Hp- β -ImHpPy- γ -ImPyPyPyPy
	1076 β p) 5'-W T T G T C W-3'	Hp- β -ImHpPy- γ -ImPyPy- β -Py
	1077 β) 5'-W T T G A T W-3'	Hp- β -ImPyHp- γ -PyHpPyPyPy
	1077 β p) 5'-W T T G A T W-3'	Hp- β -ImPyHp- γ -PyHpPy- β -Py
	1078 β) 5'-W T T G A A W-3'	Hp- β -ImPyPy- γ -HpHpPyPyPy
	1078 β p) 5'-W T T G A A W-3'	Hp- β -ImPyPy- γ -HpHpPy- β -Py
	1079 β) 5'-W T T G A G W-3'	Hp- β -ImPyIm- γ -PyHpPyPyPy
	1079 β p) 5'-W T T G A G W-3'	Hp- β -ImPyIm- γ -PyHpPy- β -Py
	1080 β) 5'-W T T G A C W-3'	Hp- β -ImPyPy- γ -ImHpPyPyPy
	1080 β p) 5'-W T T G A C W-3'	Hp- β -ImPyPy- γ -ImHpPy- β -Py
20	1081 β) 5'-W T T G G T W-3'	Hp- β -ImImHp- γ -PyPyPyPyPy
	1081 β p) 5'-W T T G G T W-3'	Hp- β -ImImHp- γ -PyPyPy- β -Py
	1082 β) 5'-W T T G G A W-3'	Hp- β -ImImPy- γ -HpPyPyPyPy
	1082 β p) 5'-W T T G G A W-3'	Hp- β -ImImPy- γ -HpPyPy- β -Py
	1083 β) 5'-W T T G C T W-3'	Hp- β -ImPyHp- γ -PyImPyPyPy
25	1083 β p) 5'-W T T G C T W-3'	Hp- β -ImPyHp- γ -PyImPy- β -Py
	1084 β) 5'-W T T G C A W-3'	Hp- β -ImPyPy- γ -HpImPyPyPy
	1084 β p) 5'-W T T G C A W-3'	Hp- β -ImPyPy- γ -HpImPy- β -Py
	1085 β) 5'-W T T G G G W-3'	Hp- β -ImImIm- γ -PyPyPyPyPy
	1085 β p) 5'-W T T G G G W-3'	Hp- β -ImImIm- γ -PyPyPy- β -Py
30	1086 β) 5'-W T T G G C W-3'	Hp- β -ImImPy- γ -ImPyPyPyPy
	1086 β p) 5'-W T T G G C W-3'	Hp- β -ImImPy- γ -ImPyPy- β -Py
	1087 β) 5'-W T T G C G W-3'	Hp- β -ImPyIm- γ -PyImPyPyPy
	1087 β p) 5'-W T T G C G W-3'	Hp- β -ImPyIm- γ -PyImPy- β -Py

TABLE 79 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTTSNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1088 β) 5'-W T T G C C W-3'	Hp- β -ImPyPy- γ -ImImPyPyPy
5	1088 β p) 5'-W T T G C C W-3'	Hp- β -ImPyPy- γ -ImImPy- β -Py
	1089 β) 5'-W T T C T T W-3'	HpHpPyHpHp- γ -Py- β -ImPyPy
	1089 β p) 5'-W T T C T T W-3'	HpHpPy- β -Hp- γ -Py- β -ImPyPy
	1090 β) 5'-W T T C T A W-3'	HpHpPyHpPy- γ -Hp- β -ImPyPy
	1090 β p) 5'-W T T C T A W-3'	HpHpPy- β -Py- γ -Hp- β -ImPyPy
10	1091 β) 5'-W T T C T G W-3'	HpHp- β -HpIm- γ -Py- β -ImPyPy
	1092 β) 5'-W T T C T C W-3'	HpHpPyHpPy- γ -Im- β -ImPyPy
	1092 β p) 5'-W T T C T C W-3'	HpHpPy- β -Py- γ -Im- β -ImPyPy
	1093 β) 5'-W T T C A T W-3'	HpHpPyPyHp- γ -Py- β -ImPyPy
	1093 β p) 5'-W T T C A T W-3'	HpHpPy- β -Hp- γ -Py- β -ImPyPy
	1094 β) 5'-W T T C A A W-3'	HpHpPyPyPy- γ -Hp- β -ImPyPy
	1094 β p) 5'-W T T C A A W-3'	HpHpPy- β -Py- γ -Hp- β -ImPyPy
	1095 β) 5'-W T T C A G W-3'	HpHp- β -PyIm- γ -Py- β -ImPyPy
	1096 β) 5'-W T T C A C W-3'	HpHpPyPyPy- γ -Im- β -ImPyPy
	1096 β p) 5'-W T T C A C W-3'	HpHpPy- β -Py- γ -Im- β -ImPyPy
	1097 β) 5'-W T T C G T W-3'	HpHp- β -ImHp- γ -Py- β -ImPyPy
	1098 β) 5'-W T T C G A W-3'	HpHp- β -ImPy- γ -Hp- β -ImPyPy
	1099 β) 5'-W T T C C T W-3'	HpHpPyPyHp- γ -PyImIm- β -Py
	1099 β p) 5'-W T T C C T W-3'	Hp- β -PyPyHp- γ -PyImIm- β -Py
	1100 β) 5'-W T T C C A W-3'	HpHpPyPyPy- γ -HpImIm- β -Py
25	1100 β p) 5'-W T T C C A W-3'	Hp- β -PyPyPy- γ -HpImIm- β -Py
	1101 β) 5'-W T T C G G W-3'	HpHp- β -ImIm- γ -Py- β -ImPyPy
	1102 β) 5'-W T T C G C W-3'	HpHp- β -ImPy- γ -Im- β -ImPyPy
	1103 β) 5'-W T T C C G W-3'	HpHp- β -PyIm- γ -PyImIm- β -Py

TABLE 80: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTAWNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1107 β) 5'-W T A T T G W-3'	HpPy- β -HpIm- γ -PyPyPyHpPy
5	1107 β p) 5'-W T A T T G W-3'	HpPy- β -HpIm- γ -PyPy- β -HpPy
	1111 β) 5'-W T A T A G W-3'	HpPy- β -PyIm- γ -PyHpPyHpPy
	1111 β p) 5'-W T A T A G W-3'	HpPy- β -PyIm- γ -PyHp- β -HpPy
	1113 β) 5'-W T A T G T W-3'	HpPy- β -ImHp- γ -PyPyPyHpPy
	1113 β p) 5'-W T A T G T W-3'	HpPy- β -ImHp- γ -PyPy- β -HpPy
10	1114 β) 5'-W T A T G A W-3'	HpPy- β -ImPy- γ -HpPyPyHpPy
	1114 β p) 5'-W T A T G A W-3'	HpPy- β -ImPy- γ -HpPy- β -HpPy
	1115 β) 5'-W T A T G G W-3'	HpPy- β -ImIm- γ -PyPyPyHpPy
	1115 β p) 5'-W T A T G G W-3'	HpPy- β -ImIm- γ -PyPy- β -HpPy
	1116 β) 5'-W T A T G C W-3'	HpPy- β -ImPy- γ -ImPyPyHpPy
15	1116 β p) 5'-W T A T G C W-3'	HpPy- β -ImPy- γ -ImPy- β -HpPy
	1119 β) 5'-W T A T C G W-3'	HpPy- β -PyIm- γ -PyImPyHpPy
	1119 β p) 5'-W T A T C G W-3'	HpPy- β -PyIm- γ -PyIm- β -HpPy
	1123 β) 5'-W T A A T G W-3'	HpPy- β -HpIm- γ -PyPyHpHpPy
	1123 β p) 5'-W T A A T G W-3'	HpPy- β -HpIm- γ -PyPy- β -HpPy
20	1127 β) 5'-W T A A A G W-3'	HpPy- β -PyIm- γ -PyHpHpHpPy
	1127 β p) 5'-W T A A A G W-3'	HpPy- β -PyIm- γ -PyHp- β -HpPy
	1129 β) 5'-W T A A G T W-3'	HpPy- β -ImHp- γ -PyPyHpHpPy
	1129 β p) 5'-W T A A G T W-3'	HpPy- β -ImHp- γ -PyPy- β -HpPy
	1130 β) 5'-W T A A G A W-3'	HpPy- β -ImPy- γ -HpPyHpHpPy
25	1130 β p) 5'-W T A A G A W-3'	HpPy- β -ImPy- γ -HpPy- β -HpPy
	1131 β) 5'-W T A A G G W-3'	HpPy- β -ImIm- γ -PyPyHpHpPy
	1131 β p) 5'-W T A A G G W-3'	HpPy- β -ImIm- γ -PyPy- β -HpPy
	1132 β) 5'-W T A A G C W-3'	HpPy- β -ImPy- γ -ImPyHpHpPy
	1132 β p) 5'-W T A A G C W-3'	HpPy- β -ImPy- γ -ImPy- β -HpPy
30	1135 β) 5'-W T A A C G W-3'	HpPy- β -PyIm- γ -PyImHpHpPy
	1135 β p) 5'-W T A A C G W-3'	HpPy- β -PyIm- γ -PyIm- β -HpPy

TABLE 81: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTASNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1137 β) 5'-W T A G T T W-3'	Hp- β -ImHpHp- γ -PyPyPyHpPy
5	1137 β p) 5'-W T A G T T W-3'	Hp- β -ImHpHp- γ -PyPyPy- β -Py
	1138 β) 5'-W T A G T A W-3'	Hp- β -ImHpPy- γ -HpPyPyHpPy
	1138 β p) 5'-W T A G T A W-3'	Hp- β -ImHpPy- γ -HpPyPy- β -Py
	1139 β) 5'-W T A G T G W-3'	Hp- β -ImHpIm- γ -PyPyPyHpPy
	1139 β p) 5'-W T A G T G W-3'	Hp- β -ImHpIm- γ -PyPyPy- β -Py
10	1140 β) 5'-W T A G T C W-3'	Hp- β -ImHpPy- γ -ImPyPyHpPy
	1140 β p) 5'-W T A G T C W-3'	Hp- β -ImHpPy- γ -ImPyPy- β -Py
	1141 β) 5'-W T A G A T W-3'	Hp- β -ImPyHp- γ -PyHpPyHpPy
	1141 β p) 5'-W T A G A T W-3'	Hp- β -ImPyHp- γ -PyHpPy- β -Py
	1142 β) 5'-W T A G A A W-3'	Hp- β -ImPyPy- γ -HpHpPyHpPy
	1142 β p) 5'-W T A G A A W-3'	Hp- β -ImPyPy- γ -HpHpPy- β -Py
	1143 β) 5'-W T A G A G W-3'	Hp- β -ImPyIm- γ -PyHpPyHpPy
	1143 β p) 5'-W T A G A G W-3'	Hp- β -ImPyIm- γ -PyHpPy- β -Py
	1144 β) 5'-W T A G A C W-3'	Hp- β -ImPyPy- γ -ImHpPyHpPy
	1144 β p) 5'-W T A G A C W-3'	Hp- β -ImPyPy- γ -ImHpPy- β -Py
20	1145 β) 5'-W T A G G T W-3'	Hp- β -ImImHp- γ -PyPyPyHpPy
	1145 β p) 5'-W T A G G T W-3'	Hp- β -ImImHp- γ -PyPyPy- β -Py
	1146 β) 5'-W T A G G A W-3'	Hp- β -ImImPy- γ -HpPyPyHpPy
	1146 β p) 5'-W T A G G A W-3'	Hp- β -ImImPy- γ -HpPyPy- β -Py
	1147 β) 5'-W T A G C T W-3'	Hp- β -ImPyHp- γ -PyImPyHpPy
25	1147 β p) 5'-W T A G C T W-3'	Hp- β -ImPyHp- γ -PyImPy- β -Py
	1148 β) 5'-W T A G C A W-3'	Hp- β -ImPyPy- γ -HpImPyHpPy
	1148 β p) 5'-W T A G C A W-3'	Hp- β -ImPyPy- γ -HpImPy- β -Py
	1149 β) 5'-W T A G G G W-3'	Hp- β -ImImIm- γ -PyPyPyHpPy
	1149 β p) 5'-W T A G G G W-3'	Hp- β -ImImIm- γ -PyPyPy- β -Py
30	1150 β) 5'-W T A G G C W-3'	Hp- β -ImImPy- γ -ImPyPyHpPy
	1150 β p) 5'-W T A G G C W-3'	Hp- β -ImImPy- γ -ImPyPy- β -Py
	1151 β) 5'-W T A G C G W-3'	Hp- β -ImPyIm- γ -PyImPyHpPy
	1151 β p) 5'-W T A G C G W-3'	Hp- β -ImPyIm- γ -PyImPy- β -Py

TABLE 81 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTASNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1152 β) 5'-W T A G C C W-3'	Hp- β -ImPyPy- γ -ImImPyHpPy
5	1152 β p) 5'-W T A G C C W-3'	Hp- β -ImPyPy- γ -ImImPy- β -Py
	1153 β) 5'-W T A C T T W-3'	HpPyPyHpHp- γ -Py- β -ImHpPy
	1153 β p) 5'-W T A C T T W-3'	HpPyPy- β -Hp- γ -Py- β -ImHpPy
	1154 β) 5'-W T A C T A W-3'	HpPyPyHpPy- γ -Hp- β -ImHpPy
	1154 β p) 5'-W T A C T A W-3'	HpPyPy- β -Py- γ -Hp- β -ImHpPy
10	1155 β) 5'-W T A C T G W-3'	HpPy- β -HpIm- γ -Py- β -ImHpPy
	1156 β) 5'-W T A C T C W-3'	HpPyPyHpPy- γ -Im- β -ImHpPy
	1156 β p) 5'-W T A C T C W-3'	HpPyPy- β -Py- γ -Im- β -ImHpPy
	1157 β) 5'-W T A C A T W-3'	HpPyPyPyHp- γ -Py- β -ImHpPy
	1157 β p) 5'-W T A C A T W-3'	HpPyPy- β -Hp- γ -Py- β -ImHpPy
	1158 β) 5'-W T A C A A W-3'	HpPyPyPyPy- γ -Hp- β -ImHpPy
	1158 β p) 5'-W T A C A A W-3'	HpPyPy- β -Py- γ -Hp- β -ImHpPy
	1159 β) 5'-W T A C A G W-3'	HpPy- β -PyIm- γ -Py- β -ImHpPy
	1160 β) 5'-W T A C A C W-3'	HpPyPyPyPy- γ -Im- β -ImHpPy
	1160 β p) 5'-W T A C A C W-3'	HpPyPy- β -Py- γ -Im- β -ImHpPy
20	1161 β) 5'-W T A C G T W-3'	HpPy- β -ImHp- γ -Py- β -ImHpPy
	1162 β) 5'-W T A C G A W-3'	HpPy- β -ImPy- γ -Hp- β -ImHpPy
	1163 β) 5'-W T A C C T W-3'	HpPyPyPyHp- γ -PyImIm- β -Py
	1163 β p) 5'-W T A C C T W-3'	Hp- β -PyPyHp- γ -PyImIm- β -Py
	1164 β) 5'-W T A C C A W-3'	HpPyPyPyPy- γ -HpImIm- β -Py
25	1164 β p) 5'-W T A C C A W-3'	Hp- β -PyPyPy- γ -HpImIm- β -Py
	1165 β) 5'-W T A C G G W-3'	HpPy- β -ImIm- γ -Py- β -ImHpPy
	1166 β) 5'-W T A C G C W-3'	HpPy- β -ImPy- γ -Im- β -ImHpPy
	1167 β) 5'-W T A C C G W-3'	HpPy- β -PyIm- γ -PyImIm- β -Py

TABLE 82: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCWNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1170 β) 5'-W T C T T A W-3'	HpPyHpHpPy- γ -HpPy- β -ImPy
5	1170 β p) 5'-W T C T T A W-3'	HpPy- β -HpPy- γ -HpPy- β -ImPy
	1171 β) 5'-W T C T T G W-3'	HpPy- β -HpIm- γ -PyPy- β -ImPy
	1172 β) 5'-W T C T T C W-3'	HpPyHpHpPy- γ -ImPy- β -ImPy
	1172 β p) 5'-W T C T T C W-3'	HpPy- β -HpPy- γ -ImPy- β -ImPy
	1173 β) 5'-W T C T A T W-3'	HpPyHpPyHp- γ -PyHp- β -ImPy
10	1173 β p) 5'-W T C T A T W-3'	HpPy- β -PyHp- γ -PyHp- β -ImPy
	1174 β) 5'-W T C T A A W-3'	HpPyHpPyPy- γ -HpHp- β -ImPy
	1174 β p) 5'-W T C T A A W-3'	HpPy- β -PyPy- γ -HpHp- β -ImPy
	1175 β) 5'-W T C T A G W-3'	HpPy- β -PyIm- γ -PyHp- β -ImPy
	1176 β) 5'-W T C T A C W-3'	HpPyHpPyPy- γ -ImHp- β -ImPy
	1176 β p) 5'-W T C T A C W-3'	HpPy- β -PyPy- γ -ImHp- β -ImPy
	1177 β) 5'-W T C T G T W-3'	HpPy- β -ImHp- γ -PyPy- β -ImPy
	1178 β) 5'-W T C T G A W-3'	HpPy- β -ImPy- γ -HpPy- β -ImPy
	1179 β) 5'-W T C T G G W-3'	HpPy- β -ImIm- γ -PyPy- β -ImPy
	1180 β) 5'-W T C T G C W-3'	HpPy- β -ImPy- γ -ImPy- β -ImPy
20	1181 β) 5'-W T C T C T W-3'	HpPyHpPyHp- γ -PyIm- β -ImPy
	1181 β p) 5'-W T C T C T W-3'	HpPy- β -PyHp- γ -PyIm- β -ImPy
	1182 β) 5'-W T C T C A W-3'	HpPyHpPyPy- γ -HpIm- β -ImPy
	1182 β p) 5'-W T C T C A W-3'	HpPy- β -PyPy- γ -HpIm- β -ImPy
	1183 β) 5'-W T C T C G W-3'	HpPy- β -PyIm- γ -PyIm- β -ImPy
25	1184 β) 5'-W T C T C C W-3'	HpPyHpPyPy- γ -ImIm- β -ImPy
	1184 β p) 5'-W T C T C C W-3'	HpPy- β -PyPy- γ -ImIm- β -ImPy
	1185 β) 5'-W T C A T T W-3'	HpPyPyHpHp- γ -PyPy- β -ImPy
	1185 β p) 5'-W T C A T T W-3'	HpPy- β -HpHp- γ -PyPy- β -ImPy
	1186 β) 5'-W T C A T A W-3'	HpPyPyHpPy- γ -HpPy- β -ImPy
30	1186 β p) 5'-W T C A T A W-3'	HpPy- β -HpPy- γ -HpPy- β -ImPy
	1187 β) 5'-W T C A T G W-3'	HpPy- β -HpIm- γ -PyPy- β -ImPy

TABLE 82 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCWNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1188 β) 5'-W T C A T C W-3'	HpPyPyHpPy- γ -ImPy- β -ImPy
5	1188 β p) 5'-W T C A T C W-3'	HpPy- β -HpPy- γ -ImPy- β -ImPy
	1189 β) 5'-W T C A A T W-3'	HpPyPyPyHpPy- γ -PyHp- β -ImPy
	1189 β p) 5'-W T C A A T W-3'	HpPy- β -PyHp- γ -PyHp- β -ImPy
	1190 β) 5'-W T C A A A W-3'	HpPyPyPyPy- γ -HpHp- β -ImPy
	1190 β p) 5'-W T C A A A W-3'	HpPy- β -PyPy- γ -HpHp- β -ImPy
10	1191 β) 5'-W T C A A G W-3'	HpPy- β -PyIm- γ -PyHp- β -ImPy
	1192 β) 5'-W T C A A C W-3'	HpPyPyPyPy- γ -ImHp- β -ImPy
	1192 β p) 5'-W T C A A C W-3'	HpPy- β -PyPy- γ -ImHp- β -ImPy
	1193 β) 5'-W T C A G T W-3'	HpPy- β -ImHp- γ -PyPy- β -ImPy
	1194 β) 5'-W T C A G A W-3'	HpPy- β -ImPy- γ -HpPy- β -ImPy
	1195 β) 5'-W T C A G G W-3'	HpPy- β -ImIm- γ -PyPy- β -ImPy
	1196 β) 5'-W T C A G C W-3'	HpPy- β -ImPy- γ -ImPy- β -ImPy
	1197 β) 5'-W T C A C T W-3'	HpPyPyPyHpPy- γ -PyIm- β -ImPy
	1197 β p) 5'-W T C A C T W-3'	HpPy- β -PyHp- γ -PyIm- β -ImPy
	1198 β) 5'-W T C A C A W-3'	HpPyPyPyPy- γ -HpIm- β -ImPy
20	1198 β p) 5'-W T C A C A W-3'	HpPy- β -PyPy- γ -HpIm- β -ImPy
	1199 β) 5'-W T C A C G W-3'	HpPy- β -PyIm- γ -PyIm- β -ImPy
	1200 β) 5'-W T C A C C W-3'	HpPyPyPyPy- γ -ImIm- β -ImPy
	1200 β p) 5'-W T C A C C W-3'	HpPy- β -PyPy- γ -ImIm- β -ImPy

TABLE 83: 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCSNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1201 β) 5'-W T C G T T W-3'	Hp- β -ImHpHp- γ -PyPy- β -ImPy
5	1202 β) 5'-W T C G T A W-3'	Hp- β -ImHpPy- γ -HpPy- β -ImPy
	1203 β) 5'-W T C G T G W-3'	Hp- β -ImHpIm- γ -PyPy- β -ImPy
	1204 β) 5'-W T C G T C W-3'	Hp- β -ImHpPy- γ -ImPy- β -ImPy
	1205 β) 5'-W T C G A T W-3'	Hp- β -ImPyHp- γ -PyHp- β -ImPy
	1206 β) 5'-W T C G A A W-3'	Hp- β -ImPyPy- γ -HpHp- β -ImPy
10	1207 β) 5'-W T C G A G W-3'	Hp- β -ImPyIm- γ -PyHp- β -ImPy
	1208 β) 5'-W T C G A C W-3'	Hp- β -ImPyPy- γ -ImHp- β -ImPy
	1209 β) 5'-W T C G G T W-3'	Hp- β -ImImHp- γ -PyPy- β -ImPy
	1210 β) 5'-W T C G G A W-3'	Hp- β -ImImPy- γ -HpPy- β -ImPy
	1211 β) 5'-W T C G C T W-3'	Hp- β -ImPyHp- γ -PyIm- β -ImPy
	1212 β) 5'-W T C G C A W-3'	Hp- β -ImPyPy- γ -HpIm- β -ImPy
	1213 β) 5'-W T C C T T W-3'	HpPyPyHpHp- γ -Py- β -ImImPy
	1213 β_p) 5'-W T C C T T W-3'	HpPyPy- β -Hp- γ -Py- β -ImImPy
	1214 β) 5'-W T C C T A W-3'	HpPyPyHpPy- γ -Hp- β -ImImPy
	1214 β_p) 5'-W T C C T A W-3'	HpPyPy- β -Py- γ -Hp- β -ImImPy
20	1215 β) 5'-W T C C T G W-3'	HpPy- β -HpIm- γ -Py- β -ImImPy
	1216 β) 5'-W T C C T C W-3'	HpPyPyHpPy- γ -Im- β -ImImPy
	1216 β_p) 5'-W T C C T C W-3'	HpPyPy- β -Py- γ -Im- β -ImImPy
	1217 β) 5'-W T C C A T W-3'	HpPyPyPyHp- γ -Py- β -ImImPy
	1217 β_p) 5'-W T C C A T W-3'	HpPyPy- β -Hp- γ -Py- β -ImImPy
25	1218 β) 5'-W T C C A A W-3'	HpPyPyPyPy- γ -Hp- β -ImImPy
	1218 β_p) 5'-W T C C A A W-3'	HpPyP- β -Py- γ -Hp- β -ImImPy
	1219 β) 5'-W T C C A G W-3'	HpPy- β -PyIm- γ -Py- β -ImImPy
	1220 β) 5'-W T C C A C W-3'	HpPyPyPyPy- γ -Im- β -ImImPy
	1220 β_p) 5'-W T C C A C W-3'	HpPyPy- β -Py- γ -Im- β -ImImPy
30	1221 β) 5'-W T C C G T W-3'	HpPy- β -ImHp- γ -Py- β -ImImPy
	1222 β) 5'-W T C C G A W-3'	HpPy- β -ImPy- γ -Hp- β -ImImPy
	1225 β) 5'-W T C G G G W-3'	Hp- β -ImImIm- γ -PyPy- β -ImPy

TABLE 83 (cont): 10-ring Hairpin Polyamides for recognition of 7-bp 5'-WTCSNNW-3' with β substitutions

	DNA sequence	aromatic amino acid sequence
	1226 β) 5'-W T C G G C W-3'	Hp- β -ImImPy- γ -ImPy- β -ImPy
5	1227 β) 5'-W T C G C G W-3'	Hp- β -ImPyIm- γ -PyIm- β -ImPy
	1228 β) 5'-W T C G C C W-3'	Hp- β -ImPyPy- γ -ImIm- β -ImPy
	1229 β) 5'-W T C C G G W-3'	HpPy- β -ImIm- γ -Py- β -ImImPy
	1230 β) 5'-W T C C G C W-3'	HpPy- β -ImPy- γ -Im- β -ImImPy
10	1231 β) 5'-W T C C C G W-3'	HpPy- β -PyIm- γ -PyImImImPy

If the process described above of designing a preferred polyamide molecule comprising four or five carboxamide binding pairs does not produce a selective polyamide that binds to the target identified DNA sequence with subnanomolar affinity and with a selectivity over mismatch sequences of greater than a factor of ten, a polyamide molecule

X₁X₂X₃X₄X₅X₆- γ -X₇X₈X₉X₁₀X₁₁X₁₂ having six carboxamide binding pairs can be designed that is selective for an eight base pair identified target 5'-WNNNNNNW-3' sequence. The design and synthesis of six binding pair polyamides is essentially the same as that of the four and five binding pair polyamides described above.

The polyamide design process for six carboxamide binding pair polyamides is shown schematically in Figure 10 A and the upper half of 10B. The method for choosing the residues that can be replaced by a β -alanine residue is shown schematically in the lower half of Figure 10 B and in Figure 11. The 1024 possible 12-ring hairpins which target the 1024 5'-GNNNNN-3' core sequences are listed in Tables 84-115. Each DNA sequence entry can be correlated to its corresponding polyamide recognition sequence using the process outlined in this figure. The 1024 possible 12-ring hairpins which target the 1024 5'-CNNNNN-3' core sequences are listed in Tables 116-147. Each DNA sequence entry can be correlated to its corresponding polyamide recognition sequence using the process outlined in this figure.

Figure 11 shows a process for replacement of aromatic amino acid residues with aliphatic β -alanine 'spring' residues in order to enhance the DNA binding properties of 12-ring hairpin polyamides. Selective placement of an aliphatic β -alanine (β) residue paired side-by-side with either a pyrrole (Py) or imidazole (Im) aromatic amino acid or another β -alanine residue is found

to compensate for sequence composition effects for recognition of the minor groove of DNA by hairpin pyrrole-imidazole polyamides. If an all-ring polyamide has been found to have an affinity which is not subnanomolar, or a specificity versus mismatch sequences which is less than 10-fold it may be caused by DNA sequence-composition effects which can be tuned out by replacement of an aromatic amino acid with an aliphatic β -alanine spring. Rules have been determined to help determine the exact placement of the β -spring residues. For example, within the 12-ring template, it is only beneficial to place β -alanine within positions X₂, X₃, X₄, X₅, X₈, X₉, and X₁₀ X₁₁. No more than two β -alanine residues may be placed within a single hairpin structure. No more than a single β -residue may be placed within each individual polyamide subunit. Tables 148-1079 list derivatives of sequences (1233-2224) labeled (1223 β -2224 β) which contain two β -alanine residues assigned according to the process outlined in Figure 11A & B.

TABLE 84: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1233) 5'-W G G G T T T W-3'	ImImImHpHpHp-γ-PyPyPyPyPyPy
5	1234) 5'-W G G G T T A W-3'	ImImImHpHpPy-γ-HpPyPyPyPyPy
	1235) 5'-W G G G T T G W-3'	ImImImHpHpIm-γ-PyPyPyPyPyPy
	1236) 5'-W G G G T T C W-3'	ImImImHpHpPy-γ-ImPyPyPyPyPy
	1237) 5'-W G G G T A T W-3'	ImImImHpPyHp-γ-PyHpPyPyPyPy
	1238) 5'-W G G G T A A W-3'	ImImImHpPyPy-γ-HpHpPyPyPyPy
10	1239) 5'-W G G G T A G W-3'	ImImImHpPyIm-γ-PyHpPyPyPyPy
	1240) 5'-W G G G T A C W-3'	ImImImHpPyPy-γ-ImHpPyPyPyPy
	1241) 5'-W G G G T G T W-3'	ImImImHpImHp-γ-PyPyPyPyPyPy
	1242) 5'-W G G G T G A W-3'	ImImImHpImPy-γ-HpPyPyPyPyPy
	1243) 5'-W G G G T G G W-3'	ImImImHpImIm-γ-PyPyPyPyPyPy
	1244) 5'-W G G G T G C W-3'	ImImImHpImPy-γ-ImPyPyPyPyPy
	1245) 5'-W G G G T C T W-3'	ImImImHpPyHp-γ-PyImPyPyPyPy
	1246) 5'-W G G G T C A W-3'	ImImImHpPyPy-γ-HpImPyPyPyPy
	1247) 5'-W G G G T C G W-3'	ImImImHpPyIm-γ-PyImPyPyPyPy
	1248) 5'-W G G G T C C W-3'	ImImImHpPyPy-γ-ImImPyPyPyPy
	1249) 5'-W G G G A T T W-3'	ImImImPyHpHp-γ-PyPyHpPyPyPy
	1250) 5'-W G G G A T A W-3'	ImImImPyHpPy-γ-HpPyHpPyPyPy
	1251) 5'-W G G G A T G W-3'	ImImImPyHpIm-γ-PyPyHpPyPyPy
	1252) 5'-W G G G A T C W-3'	ImImImPyHpPy-γ-ImPyHpPyPyPy
	1253) 5'-W G G G A A T W-3'	ImImImPyPyHp-γ-PyHpHpPyPyPy
25	1254) 5'-W G G G A A A W-3'	ImImImPyPyPy-γ-HpHpHpPyPyPy
	1255) 5'-W G G G A A G W-3'	ImImImPyPyIm-γ-PyHpHpPyPyPy
	1256) 5'-W G G G A A C W-3'	ImImImPyPyPy-γ-ImHpHpPyPyPy
	1257) 5'-W G G G A G T W-3'	ImImImPyImHp-γ-PyPyHpPyPyPy
	1258) 5'-W G G G A G A W-3'	ImImImPyImPy-γ-HpPyHpPyPyPy
30	1259) 5'-W G G G A G G W-3'	ImImImPyImIm-γ-PyPyHpPyPyPy
	1260) 5'-W G G G A G C W-3'	ImImImPyImPy-γ-ImPyHpPyPyPy
	1261) 5'-W G G G A C T W-3'	ImImImPyPyHp-γ-PyImHpPyPyPy
	1262) 5'-W G G G A C A W-3'	ImImImPyPyPy-γ-HpImHpPyPyPy
	1263) 5'-W G G G A C G W-3'	ImImImPyPyIm-γ-PyImHpPyPyPy
35	1264) 5'-W G G G A C C W-3'	ImImImPyPyPy-γ-ImImHpPyPyPy

TABLE 85: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1265) 5'-W G G G G T T W-3'	ImImImImHpHp-γ-PyPyPyPyPyPy
	1266) 5'-W G G G G T A W-3'	ImImImImHpPy-γ-HpPyPyPyPyPy
5	1267) 5'-W G G G G T G W-3'	ImImImImHpIm-γ-PyPyPyPyPyPy
	1268) 5'-W G G G G T C W-3'	ImImImImHpPy-γ-ImPyPyPyPyPy
	1269) 5'-W G G G G A T W-3'	ImImImImPyHp-γ-PyHpPyPyPyPy
	1270) 5'-W G G G G A A W-3'	ImImImImPyPy-γ-HpHpPyPyPyPy
	1271) 5'-W G G G G A G W-3'	ImImImImPyIm-γ-PyHpPyPyPyPy
10	1272) 5'-W G G G G A C W-3'	ImImImImPyPy-γ-ImHpPyPyPyPy
	1273) 5'-W G G G G G T W-3'	ImImImImImHp-γ-PyPyPyPyPyPy
	1274) 5'-W G G G G G A W-3'	ImImImImImPy-γ-HpPyPyPyPyPy
	1275) 5'-W G G G G C T W-3'	ImImImImPyHp-γ-PyImPyPyPyPy
	1276) 5'-W G G G G C A W-3'	ImImImImPyPy-γ-HpImPyPyPyPy
	1277) 5'-W G G G C T T W-3'	ImImImPyHpHp-γ-PyPyImPyPyPy
	1278) 5'-W G G G C T A W-3'	ImImImPyHpPy-γ-HpPyImPyPyPy
	1279) 5'-W G G G C T G W-3'	ImImImPyHpIm-γ-PyPyImPyPyPy
	1280) 5'-W G G G C T C W-3'	ImImImPyHpPy-γ-ImPyImPyPyPy
	1281) 5'-W G G G C A T W-3'	ImImImPyPyHp-γ-PyHpImPyPyPy
20	1282) 5'-W G G G C A A W-3'	ImImImPyPyPy-γ-HpHpImPyPyPy
	1283) 5'-W G G G C A G W-3'	ImImImPyPyIm-γ-PyHpImPyPyPy
	1284) 5'-W G G G C A C W-3'	ImImImPyPyPy-γ-ImHpImPyPyPy
	1285) 5'-W G G G C G T W-3'	ImImImPyImHp-γ-PyPyImPyPyPy
	1286) 5'-W G G G C G A W-3'	ImImImPyImPy-γ-HpPyImPyPyPy
25	1287) 5'-W G G G C C T W-3'	ImImImPyPyHp-γ-PyImImPyPyPy
	1288) 5'-W G G G C C A W-3'	ImImImPyPyPy-γ-HpImImPyPyPy
	G49) 5'-W G G G G G G W-3'	ImImImImImIm-γ-PyPyPyPyPyPy
	G50) 5'-W G G G G G C W-3'	ImImImImImPy-γ-ImPyPyPyPyPy
	G51) 5'-W G G G G C G W-3'	ImImImImPyIm-γ-PyImPyPyPyPy
30	G52) 5'-W G G G G C C W-3'	ImImImImPyPy-γ-ImImPyPyPyPy
	G53) 5'-W G G G C G G W-3'	ImImImPyImIm-γ-PyPyImPyPyPy
	G54) 5'-W G G G C G C W-3'	ImImImPyImPy-γ-ImPyImPyPyPy
	G55) 5'-W G G G C C G W-3'	ImImImPyPyIm-γ-PyImImPyPyPy
	G56) 5'-W G G G C C C W-3'	ImImImPyPyPy-γ-ImImImPyPyPy

TABLE 86: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1289) 5'-W G G T T T T W-3'	ImImHpHpHpHp-γ-PyPyPyPyPyPy
5	1290) 5'-W G G T T T A W-3'	ImImHpHpHpPy-γ-HpPyPyPyPyPy
	1291) 5'-W G G T T T G W-3'	ImImHpHpHpIm-γ-PyPyPyPyPyPy
	1292) 5'-W G G T T T C W-3'	ImImHpHpHpPy-γ-ImPyPyPyPyPy
	1293) 5'-W G G T T A T W-3'	ImImHpHpPyHp-γ-PyHpPyPyPyPy
	1294) 5'-W G G T T A A W-3'	ImImHpHpPyPy-γ-HpHpPyPyPyPy
10	1295) 5'-W G G T T A G W-3'	ImImHpHpPyIm-γ-PyHpPyPyPyPy
	1296) 5'-W G G T T A C W-3'	ImImHpHpPyPy-γ-ImHpPyPyPyPy
	1297) 5'-W G G T T G T W-3'	ImImHpHpImHp-γ-PyPyPyPyPyPy
	1298) 5'-W G G T T G A W-3'	ImImHpHpImPy-γ-HpPyPyPyPyPy
	1299) 5'-W G G T T G G W-3'	ImImHpHpImIm-γ-PyPyPyPyPyPy
15	1300) 5'-W G G T T G C W-3'	ImImHpHpImPy-γ-ImPyPyPyPyPy
	1301) 5'-W G G T T C T W-3'	ImImHpHpPyHp-γ-PyImPyPyPyPy
	1302) 5'-W G G T T C A W-3'	ImImHpHpPyPy-γ-HpImPyPyPyPy
	1303) 5'-W G G T T C G W-3'	ImImHpHpPyIm-γ-PyImPyPyPyPy
	1304) 5'-W G G T T C C W-3'	ImImHpHpPyPy-γ-ImImPyPyPyPy
20	1305) 5'-W G G T A T T W-3'	ImImHpPyHpHp-γ-PyPyHpPyPyPy
	1306) 5'-W G G T A T A W-3'	ImImHpPyHpPy-γ-HpPyHpPyPyPy
	1307) 5'-W G G T A T G W-3'	ImImHpPyHpIm-γ-PyPyHpPyPyPy
	1308) 5'-W G G T A T C W-3'	ImImHpPyHpPy-γ-ImPyHpPyPyPy
	1309) 5'-W G G T A A T W-3'	ImImHpPyPyHp-γ-PyHpHpPyPyPy
25	1310) 5'-W G G T A A A W-3'	ImImHpPyPyPy-γ-HpHpHpPyPyPy
	1311) 5'-W G G T A A G W-3'	ImImHpPyPyIm-γ-PyHpHpPyPyPy
	1312) 5'-W G G T A A C W-3'	ImImHpPyPyPy-γ-ImHpHpPyPyPy
	1313) 5'-W G G T A G T W-3'	ImImHpPyImHp-γ-PyPyHpPyPyPy
	1314) 5'-W G G T A G A W-3'	ImImHpPyImPy-γ-HpPyHpPyPyPy
30	1315) 5'-W G G T A G G W-3'	ImImHpPyImIm-γ-PyPyHpPyPyPy
	1316) 5'-W G G T A G C W-3'	ImImHpPyImPy-γ-ImPyHpPyPyPy
	1317) 5'-W G G T A C T W-3'	ImImHpPyPyHp-γ-PyImHpPyPyPy
	1318) 5'-W G G T A C A W-3'	ImImHpPyPyPy-γ-HpImHpPyPyPy
	1319) 5'-W G G T A C G W-3'	ImImHpPyPyIm-γ-PyImHpPyPyPy
35	1320) 5'-W G G T A C C W-3'	ImImHpPyPyPy-γ-ImImHpPyPyPy

TABLE 87: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1321) 5'-W G G T G T T W-3'	ImImHpImHpHp- γ -PyPyPyPyPyPy
5	1322) 5'-W G G T G T A W-3'	ImImHpImHpPy- γ -HpPyPyPyPyPy
	1323) 5'-W G G T G T G W-3'	ImImHpImHpIm- γ -PyPyPyPyPyPy
	1324) 5'-W G G T G T C W-3'	ImImHpImHpPy- γ -ImPyPyPyPyPy
	1325) 5'-W G G T G A T W-3'	ImImHpImPyHp- γ -PyHpPyPyPyPy
	1326) 5'-W G G T G A A W-3'	ImImHpImPyPy- γ -HpHpPyPyPyPy
10	1327) 5'-W G G T G A G W-3'	ImImHpImPyIm- γ -PyHpPyPyPyPy
	1328) 5'-W G G T G A C W-3'	ImImHpImPyPy- γ -ImHpPyPyPyPy
	1329) 5'-W G G T G G T W-3'	ImImHpImImHp- γ -PyPyPyPyPyPy
	1330) 5'-W G G T G G A W-3'	ImImHpImImPy- γ -HpPyPyPyPyPy
	1331) 5'-W G G T G C T W-3'	ImImHpImPyHp- γ -PyImPyPyPyPy
	1332) 5'-W G G T G C A W-3'	ImImHpImPyPy- γ -HpImPyPyPyPy
	1333) 5'-W G G T G G G W-3'	ImImHpImImIm- γ -PyPyPyPyPyPy
	1334) 5'-W G G T G G C W-3'	ImImHpImImPy- γ -ImPyPyPyPyPy
	1335) 5'-W G G T G C G W-3'	ImImHpImPyIm- γ -PyImPyPyPyPy
	1336) 5'-W G G T G C C W-3'	ImImHpImPyPy- γ -ImImPyPyPyPy
20	1337) 5'-W G G T C T T W-3'	ImImHpPyHpHp- γ -PyPyImPyPyPy
	1338) 5'-W G G T C T A W-3'	ImImHpPyHpPy- γ -HpPyImPyPyPy
	1339) 5'-W G G T C T G W-3'	ImImHpPyHpIm- γ -PyPyImPyPyPy
	1340) 5'-W G G T C T C W-3'	ImImHpPyHpPy- γ -ImPyImPyPyPy
	1341) 5'-W G G T C A T W-3'	ImImHpPyPyHp- γ -PyHpImPyPyPy
25	1342) 5'-W G G T C A A W-3'	ImImHpPyPyPy- γ -HpHpImPyPyPy
	1343) 5'-W G G T C A G W-3'	ImImHpPyPyIm- γ -PyHpImPyPyPy
	1344) 5'-W G G T C A C W-3'	ImImHpPyPyPy- γ -ImHpImPyPyPy
	1345) 5'-W G G T C G T W-3'	ImImHpPyImHp- γ -PyPyImPyPyPy
	1346) 5'-W G G T C G A W-3'	ImImHpPyImPy- γ -HpPyImPyPyPy
30	1347) 5'-W G G T C C T W-3'	ImImHpPyPyHp- γ -PyImImPyPyPy
	1348) 5'-W G G T C C A W-3'	ImImHpPyPyPy- γ -HpImImPyPyPy
	1349) 5'-W G G T C G G W-3'	ImImHpPyImIm- γ -PyPyImPyPyPy
	1350) 5'-W G G T C G C W-3'	ImImHpPyImPy- γ -ImPyImPyPyPy
	1351) 5'-W G G T C C G W-3'	ImImHpPyPyIm- γ -PyImImPyPyPy
35	1352) 5'-W G G T C C C W-3'	ImImHpPyPyPy- γ -ImImImPyPyPy

TABLE 88: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1353) 5'-W G G A T T T W-3'	ImImPyHpHpHp- γ -PyPyPyHpPyPy
5	1354) 5'-W G G A T T A W-3'	ImImPyHpHpPy- γ -HpPyPyHpPyPy
	1355) 5'-W G G A T T G W-3'	ImImPyHpHpIm- γ -PyPyPyHpPyPy
	1356) 5'-W G G A T T C W-3'	ImImPyHpHpPy- γ -ImPyPyHpPyPy
	1357) 5'-W G G A T A T W-3'	ImImPyHpPyHp- γ -PyHpPyHpPyPy
	1358) 5'-W G G A T A A W-3'	ImImPyHpPyPy- γ -HpHpPyHpPyPy
10	1359) 5'-W G G A T A G W-3'	ImImPyHpPyIm- γ -PyHpPyHpPyPy
	1360) 5'-W G G A T A C W-3'	ImImPyHpPyPy- γ -ImHpPyHpPyPy
	1361) 5'-W G G A T G T W-3'	ImImPyHpImHp- γ -PyPyPyHpPyPy
	1362) 5'-W G G A T G A W-3'	ImImPyHpImPy- γ -HpPyPyHpPyPy
	1363) 5'-W G G A T G G W-3'	ImImPyHpImIm- γ -PyPyPyHpPyPy
	1364) 5'-W G G A T G C W-3'	ImImPyHpImPy- γ -ImPyPyHpPyPy
	1365) 5'-W G G A T C T W-3'	ImImPyHpPyHp- γ -PyImPyHpPyPy
	1366) 5'-W G G A T C A W-3'	ImImPyHpPyPy- γ -HpImPyHpPyPy
	1367) 5'-W G G A T C G W-3'	ImImPyHpPyIm- γ -PyImPyHpPyPy
	1368) 5'-W G G A T C C W-3'	ImImPyHpPyPy- γ -ImImPyHpPyPy
20	1369) 5'-W G G A A T T W-3'	ImImPyPyHpHp- γ -PyPyHpHpPyPy
	1370) 5'-W G G A A T A W-3'	ImImPyPyHpPy- γ -HpPyHpHpPyPy
	1371) 5'-W G G A A T G W-3'	ImImPyPyHpIm- γ -PyPyHpHpPyPy
	1372) 5'-W G G A A T C W-3'	ImImPyPyHpPy- γ -ImPyHpHpPyPy
	1373) 5'-W G G A A A T W-3'	ImImPyPyPyHp- γ -PyHpHpHpPyPy
25	1374) 5'-W G G A A A A W-3'	ImImPyPyPyPy- γ -HpHpHpHpPyPy
	1375) 5'-W G G A A A G W-3'	ImImPyPyPyIm- γ -PyHpHpHpPyPy
	1376) 5'-W G G A A A C W-3'	ImImPyPyPyPy- γ -ImHpHpHpPyPy
	1377) 5'-W G G A A G T W-3'	ImImPyPyImHp- γ -PyPyHpHpPyPy
	1378) 5'-W G G A A G A W-3'	ImImPyPyImPy- γ -HpPyHpHpPyPy
30	1379) 5'-W G G A A G G W-3'	ImImPyPyImIm- γ -PyPyHpHpPyPy
	1380) 5'-W G G A A G C W-3'	ImImPyPyImPy- γ -ImPyHpHpPyPy
	1381) 5'-W G G A A C T W-3'	ImImPyPyPyHp- γ -PyImHpHpPyPy
	1382) 5'-W G G A A C A W-3'	ImImPyPyPyPy- γ -HpImHpHpPyPy
	1383) 5'-W G G A A C G W-3'	ImImPyPyPyIm- γ -PyImHpHpPyPy
35	1384) 5'-W G G A A C C W-3'	ImImPyPyPyPy- γ -ImImHpHpPyPy

TABLE 89: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1385) 5'-W G G A G T T W-3'	ImImPyImHpHp- γ -PyPyPyHpPyPy
5	1386) 5'-W G G A G T A W-3'	ImImPyImHpPy- γ -HpPyPyHpPyPy
	1387) 5'-W G G A G T G W-3'	ImImPyImHpIm- γ -PyPyPyHpPyPy
	1388) 5'-W G G A G T C W-3'	ImImPyImHpPy- γ -ImPyPyHpPyPy
	1389) 5'-W G G A G A T W-3'	ImImPyImPyHp- γ -PyHpPyHpPyPy
	1390) 5'-W G G A G A A W-3'	ImImPyImPyPy- γ -HpHpPyHpPyPy
10	1391) 5'-W G G A G A G W-3'	ImImPyImPyIm- γ -PyHpPyHpPyPy
	1392) 5'-W G G A G A C W-3'	ImImPyImPyPy- γ -ImHpPyHpPyPy
	1393) 5'-W G G A G G T W-3'	ImImPyImImHp- γ -PyPyPyHpPyPy
	1394) 5'-W G G A G G A W-3'	ImImPyImImPy- γ -HpPyPyHpPyPy
	1395) 5'-W G G A G C T W-3'	ImImPyImPyHp- γ -PyImPyHpPyPy
	1396) 5'-W G G A G C A W-3'	ImImPyImPyPy- γ -HpImPyHpPyPy
	1397) 5'-W G G A G G G W-3'	ImImPyImImIm- γ -PyPyPyHpPyPy
	1398) 5'-W G G A G G C W-3'	ImImPyImImPy- γ -ImPyPyHpPyPy
	1399) 5'-W G G A G C G W-3'	ImImPyImPyIm- γ -PyImPyHpPyPy
	1400) 5'-W G G A G C C W-3'	ImImPyImPyPy- γ -ImImPyHpPyPy
	1401) 5'-W G G A C T T W-3'	ImImPyPyHpHp- γ -PyPyImHpPyPy
	1402) 5'-W G G A C T A W-3'	ImImPyPyHpPy- γ -HpPyImHpPyPy
	1403) 5'-W G G A C T G W-3'	ImImPyPyHpIm- γ -PyPyImHpPyPy
	1404) 5'-W G G A C T C W-3'	ImImPyPyHpPy- γ -ImPyImHpPyPy
	1405) 5'-W G G A C A T W-3'	ImImPyPyPyHp- γ -PyHpImHpPyPy
25	1406) 5'-W G G A C A A W-3'	ImImPyPyPyPy- γ -HpHpImHpPyPy
	1407) 5'-W G G A C A G W-3'	ImImPyPyPyIm- γ -PyHpImHpPyPy
	1408) 5'-W G G A C A C W-3'	ImImPyPyPyPy- γ -ImHpImHpPyPy
	1409) 5'-W G G A C G T W-3'	ImImPyPyImHp- γ -PyPyImHpPyPy
	1410) 5'-W G G A C G A W-3'	ImImPyPyImPy- γ -HpPyImHpPyPy
30	1411) 5'-W G G A C C T W-3'	ImImPyPyPyHp- γ -PyImImHpPyPy
	1412) 5'-W G G A C C A W-3'	ImImPyPyPyPy- γ -HpImImHpPyPy
	1413) 5'-W G G A C G G W-3'	ImImPyPyImIm- γ -PyPyImHpPyPy
	1414) 5'-W G G A C G C W-3'	ImImPyPyImPy- γ -ImPyImHpPyPy
	1415) 5'-W G G A C C G W-3'	ImImPyPyPyIm- γ -PyImImHpPyPy
35	1416) 5'-W G G A C C C W-3'	ImImPyPyPyPy- γ -ImImImHpPyPy

TABLE 90: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1417) 5'-W G G C T T T W-3'	ImImPyHpHpHp- γ -PyPyPyImPyPy
5	1418) 5'-W G G C T T A W-3'	ImImPyHpHpPy- γ -HpPyPyImPyPy
	1419) 5'-W G G C T T G W-3'	ImImPyHpHpIm- γ -PyPyPyImPyPy
	1420) 5'-W G G C T T C W-3'	ImImPyHpHpPy- γ -ImPyPyImPyPy
	1421) 5'-W G G C T A T W-3'	ImImPyHpPyHp- γ -PyHpPyImPyPy
	1422) 5'-W G G C T A A W-3'	ImImPyHpPyPy- γ -HpHpPyImPyPy
10	1423) 5'-W G G C T A G W-3'	ImImPyHpPyIm- γ -PyHpPyImPyPy
	1424) 5'-W G G C T A C W-3'	ImImPyHpPyPy- γ -ImHpPyImPyPy
	1425) 5'-W G G C T G T W-3'	ImImPyHpImHp- γ -PyPyPyImPyPy
	1426) 5'-W G G C T G A W-3'	ImImPyHpImPy- γ -HpPyPyImPyPy
	1427) 5'-W G G C T G G W-3'	ImImPyHpImIm- γ -PyPyPyImPyPy
15	1428) 5'-W G G C T G C W-3'	ImImPyHpImPy- γ -ImPyPyImPyPy
	1429) 5'-W G G C T C T W-3'	ImImPyHpPyHp- γ -PyImPyImPyPy
	1430) 5'-W G G C T C A W-3'	ImImPyHpPyPy- γ -HpImPyImPyPy
	1431) 5'-W G G C T C G W-3'	ImImPyHpPyIm- γ -PyImPyImPyPy
	1432) 5'-W G G C T C C W-3'	ImImPyHpPyPy- γ -ImImPyImPyPy
20	1433) 5'-W G G C A T T W-3'	ImImPyPyHpHp- γ -PyPyHpImPyPy
	1434) 5'-W G G C A T A W-3'	ImImPyPyHpPy- γ -HpPyHpImPyPy
	1435) 5'-W G G C A T G W-3'	ImImPyPyHpIm- γ -PyPyHpImPyPy
	1436) 5'-W G G C A T C W-3'	ImImPyPyHpPy- γ -ImPyHpImPyPy
	1437) 5'-W G G C A A T W-3'	ImImPyPyPyHp- γ -PyHpHpImPyPy
25	1438) 5'-W G G C A A A W-3'	ImImPyPyPyPy- γ -HpHpHpImPyPy
	1439) 5'-W G G C A A G W-3'	ImImPyPyPyIm- γ -PyHpHpImPyPy
	1440) 5'-W G G C A A C W-3'	ImImPyPyPyPy- γ -ImHpHpImPyPy
	1441) 5'-W G G C A G T W-3'	ImImPyPyImHp- γ -PyPyHpImPyPy
	1442) 5'-W G G C A G A W-3'	ImImPyPyImPy- γ -HpPyHpImPyPy
30	1443) 5'-W G G C A G G W-3'	ImImPyPyImIm- γ -PyPyHpImPyPy
	1444) 5'-W G G C A G C W-3'	ImImPyPyImPy- γ -ImPyHpImPyPy
	1445) 5'-W G G C A C T W-3'	ImImPyPyPyHp- γ -PyImHpImPyPy
	1446) 5'-W G G C A C A W-3'	ImImPyPyPyPy- γ -HpImHpImPyPy
	1447) 5'-W G G C A C G W-3'	ImImPyPyPyIm- γ -PyImHpImPyPy
35	1448) 5'-W G G C A C C W-3'	ImImPyPyPyPy- γ -ImImHpImPyPy

TABLE 91: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGGCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1449) 5'-W G G C G T T W-3'	ImImPyImHpHp-γ-PyPyPyImPyPy
5	1450) 5'-W G G C G T A W-3'	ImImPyImHpPy-γ-HpPyPyImPyPy
	1451) 5'-W G G C G T G W-3'	ImImPyImHpIm-γ-PyPyPyImPyPy
	1452) 5'-W G G C G T C W-3'	ImImPyImHpPy-γ-ImPyPyImPyPy
	1453) 5'-W G G C G A T W-3'	ImImPyImPyHp-γ-PyHpPyImPyPy
	1454) 5'-W G G C G A A W-3'	ImImPyImPyPy-γ-HpHpPyImPyPy
10	1455) 5'-W G G C G A G W-3'	ImImPyImPyIm-γ-PyHpPyImPyPy
	1456) 5'-W G G C G A C W-3'	ImImPyImPyPy-γ-ImHpPyImPyPy
	1457) 5'-W G G C G G T W-3'	ImImPyImImHp-γ-PyPyPyImPyPy
	1458) 5'-W G G C G G A W-3'	ImImPyImImPy-γ-HpPyPyImPyPy
	1459) 5'-W G G C G C T W-3'	ImImPyImPyHp-γ-PyImPyImPyPy
15	1460) 5'-W G G C G C A W-3'	ImImPyImPyPy-γ-HpImPyImPyPy
	1461) 5'-W G G C C T T W-3'	ImImPyPyHpHp-γ-PyPyImImPyPy
	1462) 5'-W G G C C T A W-3'	ImImPyPyHpPy-γ-HpPyImImPyPy
	1463) 5'-W G G C C T G W-3'	ImImPyPyHpIm-γ-PyPyImImPyPy
	1464) 5'-W G G C C T C W-3'	ImImPyPyHpPy-γ-ImPyImImPyPy
20	1465) 5'-W G G C C A T W-3'	ImImPyPyPyHp-γ-PyHpImImPyPy
	1466) 5'-W G G C C A A W-3'	ImImPyPyPyPy-γ-HpHpImImPyPy
	1467) 5'-W G G C C A G W-3'	ImImPyPyPyIm-γ-PyHpImImPyPy
	1468) 5'-W G G C C A C W-3'	ImImPyPyPyPy-γ-ImHpImImPyPy
	1469) 5'-W G G C C G T W-3'	ImImPyPyImHp-γ-PyPyImImPyPy
25	1470) 5'-W G G C C G A W-3'	ImImPyPyImPy-γ-HpPyImImPyPy
	1471) 5'-W G G C C C T W-3'	ImImPyPyPyHp-γ-PyImImImPyPy
	1472) 5'-W G G C C C A W-3'	ImImPyPyPyPy-γ-HpImImImPyPy
	G57) 5'-W G G C G G G W-3'	ImImPyImImIm-γ-PyPyPyImPyPy
	G58) 5'-W G G C G G C W-3'	ImImPyImImPy-γ-ImPyPyImPyPy
30	G59) 5'-W G G C G C G W-3'	ImImPyImPyIm-γ-PyImPyImPyPy
	G60) 5'-W G G C G C C W-3'	ImImPyImPyPy-γ-ImImPyImPyPy
	G61) 5'-W G G C C G G W-3'	ImImPyPyImIm-γ-PyPyImImPyPy
	G62) 5'-W G G C C G C W-3'	ImImPyPyImPy-γ-ImPyImImPyPy
	G63) 5'-W G G C C C G W-3'	ImImPyPyPyIm-γ-PyImImImPyPy
35	G64) 5'-W G G C C C C W-3'	ImImPyPyPyPy-γ-ImImImImPyPy

TABLE 92: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGGWNW-3'

	DNA sequence	aromatic amino acid sequence
	1473) 5'-W G C G T T T W-3'	ImPyImHpHpHp-γ-PyPyPyPyImPy
5	1474) 5'-W G C G T T A W-3'	ImPyImHpHpPy-γ-HpPyPyPyImPy
	1475) 5'-W G C G T T G W-3'	ImPyImHpHpIm-γ-PyPyPyPyImPy
	1476) 5'-W G C G T T C W-3'	ImPyImHpHpPy-γ-ImPyPyPyImPy
	1477) 5'-W G C G T A T W-3'	ImPyImHpPyHp-γ-PyHpPyPyImPy
	1478) 5'-W G C G T A A W-3'	ImPyImHpPyPy-γ-HpHpPyPyImPy
10	1479) 5'-W G C G T A G W-3'	ImPyImHpPyIm-γ-PyHpPyPyImPy
	1480) 5'-W G C G T A C W-3'	ImPyImHpPyPy-γ-ImHpPyPyImPy
	1481) 5'-W G C G T G T W-3'	ImPyImHpImHp-γ-PyPyPyPyImPy
	1482) 5'-W G C G T G A W-3'	ImPyImHpImPy-γ-HpPyPyPyImPy
	1483) 5'-W G C G T G G W-3'	ImPyImHpImIm-γ-PyPyPyPyImPy
	1484) 5'-W G C G T G C W-3'	ImPyImHpImPy-γ-ImPyPyPyImPy
	1485) 5'-W G C G T C T W-3'	ImPyImHpPyHp-γ-PyImPyPyImPy
	1486) 5'-W G C G T C A W-3'	ImPyImHpPyPy-γ-HpImPyPyImPy
	1487) 5'-W G C G T C G W-3'	ImPyImHpPyIm-γ-PyImPyPyImPy
	1488) 5'-W G C G T C C W-3'	ImPyImHpPyPy-γ-ImImPyPyImPy
20	1489) 5'-W G C G A T T W-3'	ImPyImPyHpHp-γ-PyPyHpPyImPy
	1490) 5'-W G C G A T A W-3'	ImPyImPyHpPy-γ-HpPyHpPyImPy
	1491) 5'-W G C G A T G W-3'	ImPyImPyHpIm-γ-PyPyHpPyImPy
	1492) 5'-W G C G A T C W-3'	ImPyImPyHpPy-γ-ImPyHpPyImPy
	1493) 5'-W G C G A A T W-3'	ImPyImPyPyHp-γ-PyHpHpPyImPy
25	1494) 5'-W G C G A A A W-3'	ImPyImPyPyPy-γ-HpHpHpPyImPy
	1495) 5'-W G C G A A G W-3'	ImPyImPyPyIm-γ-PyHpHpPyImPy
	1496) 5'-W G C G A A C W-3'	ImPyImPyPyPy-γ-ImHpHpPyImPy
	1497) 5'-W G C G A G T W-3'	ImPyImPyImHp-γ-PyPyHpPyImPy
	1498) 5'-W G C G A G A W-3'	ImPyImPyImPy-γ-HpPyHpPyImPy
30	1499) 5'-W G C G A G G W-3'	ImPyImPyImIm-γ-PyPyHpPyImPy
	1490) 5'-W G C G A G C W-3'	ImPyImPyImPy-γ-ImPyHpPyImPy
	1501) 5'-W G C G A C T W-3'	ImPyImPyPyHp-γ-PyImHpPyImPy
	1502) 5'-W G C G A C A W-3'	ImPyImPyPyPy-γ-HpImHpPyImPy
	1503) 5'-W G C G A C G W-3'	ImPyImPyPyIm-γ-PyImHpPyImPy
35	1504) 5'-W G C G A C C W-3'	ImPyImPyPyPy-γ-ImImHpPyImPy

TABLE 93: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1505) 5'-W G C G G T T W-3'	ImPyImImHpHp- γ -PyPyPyPyImPy
5	1506) 5'-W G C G G T A W-3'	ImPyImImHpPy- γ -HpPyPyPyImPy
	1507) 5'-W G C G G T G W-3'	ImPyImImHpIm- γ -PyPyPyPyImPy
	1508) 5'-W G C G G T C W-3'	ImPyImImHpPy- γ -ImPyPyPyImPy
	1509) 5'-W G C G G A T W-3'	ImPyImImPyHp- γ -PyHpPyPyImPy
	1510) 5'-W G C G G A A W-3'	ImPyImImPyPy- γ -HpHpPyPyImPy
10	1511) 5'-W G C G G A G W-3'	ImPyImImPyIm- γ -PyHpPyPyImPy
	1512) 5'-W G C G G A C W-3'	ImPyImImPyPy- γ -ImHpPyPyImPy
	1513) 5'-W G C G G G T W-3'	ImPyImImImHp- γ -PyPyPyPyImPy
	1514) 5'-W G C G G G A W-3'	ImPyImImImPy- γ -HpPyPyPyImPy
	1515) 5'-W G C G G C T W-3'	ImPyImImPyHp- γ -PyImPyPyImPy
	1516) 5'-W G C G G C A W-3'	ImPyImImPyPy- γ -HpImPyPyImPy
	1517) 5'-W G C G C T T W-3'	ImPyImPyHpHp- γ -PyPyImPyImPy
	1518) 5'-W G C G C T A W-3'	ImPyImPyHpPy- γ -HpPyImPyImPy
	1519) 5'-W G C G C T G W-3'	ImPyImPyHpIm- γ -PyPyImPyImPy
	1520) 5'-W G C G C T C W-3'	ImPyImPyHpPy- γ -ImPyImPyImPy
20	1521) 5'-W G C G C A T W-3'	ImPyImPyPyHp- γ -PyHpImPyImPy
	1522) 5'-W G C G C A A W-3'	ImPyImPyPyPy- γ -HpHpImPyImPy
	1523) 5'-W G C G C A G W-3'	ImPyImPyPyIm- γ -PyHpImPyImPy
	1524) 5'-W G C G C A C W-3'	ImPyImPyPyPy- γ -ImHpImPyImPy
	1525) 5'-W G C G C G T W-3'	ImPyImPyImHp- γ -PyPyImPyImPy
25	1526) 5'-W G C G C G A W-3'	ImPyImPyImPy- γ -HpPyImPyImPy
	1527) 5'-W G C G C C T W-3'	ImPyImPyPyHp- γ -PyImImPyImPy
	1528) 5'-W G C G C C A W-3'	ImPyImPyPyPy- γ -HpImImPyImPy
	G65) 5'-W G C G G G G W-3'	ImPyImImImIm- γ -PyPyPyPyImPy
	G66) 5'-W G C G G G C W-3'	ImPyImImImPy- γ -ImPyPyPyImPy
30	G67) 5'-W G C G G C G W-3'	ImPyImImPyIm- γ -PyImPyPyImPy
	G68) 5'-W G C G G C C W-3'	ImPyImImPyPy- γ -ImImPyPyImPy
	G69) 5'-W G C G C G G W-3'	ImPyImPyImIm- γ -PyPyImPyImPy
	G70) 5'-W G C G C G C W-3'	ImPyImPyImPy- γ -ImPyImPyImPy
	G71) 5'-W G C G C C G W-3'	ImPyImPyPyIm- γ -PyImImPyImPy
35	G72) 5'-W G C G C C C W-3'	ImPyImPyPyPy- γ -ImImImPyImPy

TABLE 94: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1529) 5'-W G C T T T T W-3'	ImPyHpHpHpHp- γ -PyPyPyPyImPy
5	1530) 5'-W G C T T T A W-3'	ImPyHpHpHpPy- γ -HpPyPyPyImPy
	1531) 5'-W G C T T T G W-3'	ImPyHpHpHpIm- γ -PyPyPyPyImPy
	1532) 5'-W G C T T T C W-3'	ImPyHpHpHpPy- γ -ImPyPyPyImPy
	1533) 5'-W G C T T A T W-3'	ImPyHpHpPyHp- γ -PyHpPyPyImPy
	1534) 5'-W G C T T A A W-3'	ImPyHpHpPyPy- γ -HpHpPyPyImPy
10	1535) 5'-W G C T T A G W-3'	ImPyHpHpPyIm- γ -PyHpPyPyImPy
	1536) 5'-W G C T T A C W-3'	ImPyHpHpPyPy- γ -ImHpPyPyImPy
	1537) 5'-W G C T T G T W-3'	ImPyHpHpImHp- γ -PyPyPyPyImPy
	1538) 5'-W G C T T G A W-3'	ImPyHpHpImPy- γ -HpPyPyPyImPy
	1539) 5'-W G C T T G G W-3'	ImPyHpHpImIm- γ -PyPyPyPyImPy
15	1540) 5'-W G C T T G C W-3'	ImPyHpHpImPy- γ -ImPyPyPyImPy
	1541) 5'-W G C T T C T W-3'	ImPyHpHpPyHp- γ -PyImPyPyImPy
	1542) 5'-W G C T T C A W-3'	ImPyHpHpPyPy- γ -HpImPyPyImPy
	1543) 5'-W G C T T C G W-3'	ImPyHpHpPyIm- γ -PyImPyPyImPy
	1544) 5'-W G C T T C C W-3'	ImPyHpHpPyPy- γ -ImImPyPyImPy
20	1545) 5'-W G C T A T T W-3'	ImPyHpPyHpHp- γ -PyPyHpPyImPy
	1546) 5'-W G C T A T A W-3'	ImPyHpPyHpPy- γ -HpPyHpPyImPy
	1547) 5'-W G C T A T G W-3'	ImPyHpPyHpIm- γ -PyPyHpPyImPy
	1548) 5'-W G C T A T C W-3'	ImPyHpPyHpPy- γ -ImPyHpPyImPy
	1549) 5'-W G C T A A T W-3'	ImPyHpPyPyHp- γ -PyHpHpPyImPy
25	1550) 5'-W G C T A A A W-3'	ImPyHpPyPyPy- γ -HpHpHpPyImPy
	1551) 5'-W G C T A A G W-3'	ImPyHpPyPyIm- γ -PyHpHpPyImPy
	1552) 5'-W G C T A A C W-3'	ImPyHpPyPyPy- γ -ImHpHpPyImPy
	1553) 5'-W G C T A G T W-3'	ImPyHpPyImHp- γ -PyPyHpPyImPy
	1554) 5'-W G C T A G A W-3'	ImPyHpPyImPy- γ -HpPyHpPyImPy
30	1555) 5'-W G C T A G G W-3'	ImPyHpPyImIm- γ -PyPyHpPyImPy
	1556) 5'-W G C T A G C W-3'	ImPyHpPyImPy- γ -ImPyHpPyImPy
	1557) 5'-W G C T A C T W-3'	ImPyHpPyPyHp- γ -PyImHpPyImPy
	1558) 5'-W G C T A C A W-3'	ImPyHpPyPyPy- γ -HpImHpPyImPy
	1559) 5'-W G C T A C G W-3'	ImPyHpPyPyIm- γ -PyImHpPyImPy
35	1560) 5'-W G C T A C C W-3'	ImPyHpPyPyPy- γ -ImImHpPyImPy

TABLE 95: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCTSNW-3'

	DNA sequence	aromatic amino acid sequence
	1561) 5'-W G C T G T T W-3'	ImPyHpImHpHp- γ -PyPyPyPyImPy
5	1562) 5'-W G C T G T A W-3'	ImPyHpImHpPy- γ -HpPyPyPyImPy
	1563) 5'-W G C T G T G W-3'	ImPyHpImHpIm- γ -PyPyPyPyImPy
	1564) 5'-W G C T G T C W-3'	ImPyHpImHpPy- γ -ImPyPyPyImPy
	1565) 5'-W G C T G A T W-3'	ImPyHpImPyHp- γ -PyHpPyPyImPy
	1566) 5'-W G C T G A A W-3'	ImPyHpImPyPy- γ -HpHpPyPyImPy
10	1567) 5'-W G C T G A G W-3'	ImPyHpImPyIm- γ -PyHpPyPyImPy
	1568) 5'-W G C T G A C W-3'	ImPyHpImPyPy- γ -ImHpPyPyImPy
	1569) 5'-W G C T G G T W-3'	ImPyHpImImHp- γ -PyPyPyPyImPy
	1570) 5'-W G C T G G A W-3'	ImPyHpImImPy- γ -HpPyPyPyImPy
	1571) 5'-W G C T G C T W-3'	ImPyHpImPyHp- γ -PyImPyPyImPy
	1572) 5'-W G C T G C A W-3'	ImPyHpImPyPy- γ -HpImPyPyImPy
	1573) 5'-W G C T G G G W-3'	ImPyHpImImIm- γ -PyPyPyPyImPy
	1574) 5'-W G C T G G C W-3'	ImPyHpImImPy- γ -ImPyPyPyImPy
	1575) 5'-W G C T G C G W-3'	ImPyHpImPyIm- γ -PyImPyPyImPy
	1576) 5'-W G C T G C C W-3'	ImPyHpImPyPy- γ -ImImPyPyImPy
	1577) 5'-W G C T C T T W-3'	ImPyHpPyHpHp- γ -PyPyImPyImPy
	1578) 5'-W G C T C T A W-3'	ImPyHpPyHpPy- γ -HpPyImPyImPy
	1579) 5'-W G C T C T G W-3'	ImPyHpPyHpIm- γ -PyPyImPyImPy
	1580) 5'-W G C T C T C W-3'	ImPyHpPyHpPy- γ -ImPyImPyImPy
	1581) 5'-W G C T C A T W-3'	ImPyHpPyPyHp- γ -PyHpImPyImPy
25	1582) 5'-W G C T C A A W-3'	ImPyHpPyPyPy- γ -HpHpImPyImPy
	1583) 5'-W G C T C A G W-3'	ImPyHpPyPyIm- γ -PyHpImPyImPy
	1584) 5'-W G C T C A C W-3'	ImPyHpPyPyPy- γ -ImHpImPyImPy
	1585) 5'-W G C T C G T W-3'	ImPyHpPyImHp- γ -PyPyImPyImPy
	1586) 5'-W G C T C G A W-3'	ImPyHpPyImPy- γ -HpPyImPyImPy
30	1587) 5'-W G C T C C T W-3'	ImPyHpPyPyHp- γ -PyImImPyImPy
	1588) 5'-W G C T C C A W-3'	ImPyHpPyPyPy- γ -HpImImPyImPy
	1589) 5'-W G C T C G G W-3'	ImPyHpPyImIm- γ -PyPyImPyImPy
	1590) 5'-W G C T C G C W-3'	ImPyHpPyImPy- γ -ImPyImPyImPy
	1591) 5'-W G C T C C G W-3'	ImPyHpPyPyIm- γ -PyImImPyImPy
35	1592) 5'-W G C T C C C W-3'	ImPyHpPyPyPy- γ -ImImImPyImPy

TABLE 96: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1593) 5'-W G C A T T T W-3'	ImPyPyHpHpHp-γ-PyPyPyHpImPy
5	1594) 5'-W G C A T T A W-3'	ImPyPyHpHpPy-γ-HpPyPyHpImPy
	1595) 5'-W G C A T T G W-3'	ImPyPyHpHpIm-γ-PyPyPyHpImPy
	1596) 5'-W G C A T T C W-3'	ImPyPyHpHpPy-γ-ImPyPyHpImPy
	1597) 5'-W G C A T A T W-3'	ImPyPyHpPyHp-γ-PyHpPyHpImPy
	1598) 5'-W G C A T A A W-3'	ImPyPyHpPyPy-γ-HpHpPyHpImPy
10	1599) 5'-W G C A T A G W-3'	ImPyPyHpPyIm-γ-PyHpPyHpImPy
	1600) 5'-W G C A T A C W-3'	ImPyPyHpPyPy-γ-ImHpPyHpImPy
	1601) 5'-W G C A T G T W-3'	ImPyPyHpImHp-γ-PyPyPyHpImPy
	1602) 5'-W G C A T G A W-3'	ImPyPyHpImPy-γ-HpPyPyHpImPy
	1603) 5'-W G C A T G G W-3'	ImPyPyHpImIm-γ-PyPyPyHpImPy
	1604) 5'-W G C A T G C W-3'	ImPyPyHpImPy-γ-ImPyPyHpImPy
	1605) 5'-W G C A T C T W-3'	ImPyPyHpPyHp-γ-PyImPyHpImPy
	1606) 5'-W G C A T C A W-3'	ImPyPyHpPyPy-γ-HpImPyHpImPy
	1607) 5'-W G C A T C G W-3'	ImPyPyHpPyIm-γ-PyImPyHpImPy
	1608) 5'-W G C A T C C W-3'	ImPyPyHpPyPy-γ-ImImPyHpImPy
20	1609) 5'-W G C A A T T W-3'	ImPyPyPyHpHp-γ-PyPyHpHpImPy
	1610) 5'-W G C A A T A W-3'	ImPyPyPyHpPy-γ-HpPyHpHpImPy
	1611) 5'-W G C A A T G W-3'	ImPyPyPyHpIm-γ-PyPyHpHpImPy
	1612) 5'-W G C A A T C W-3'	ImPyPyPyHpPy-γ-ImPyHpHpImPy
	1613) 5'-W G C A A A T W-3'	ImPyPyPyPyHp-γ-PyHpHpHpImPy
25	1614) 5'-W G C A A A A W-3'	ImPyPyPyPyPy-γ-HpHpHpHpImPy
	1615) 5'-W G C A A A G W-3'	ImPyPyPyPyIm-γ-PyHpHpHpImPy
	1616) 5'-W G C A A A C W-3'	ImPyPyPyPyPy-γ-ImHpHpHpImPy
	1617) 5'-W G C A A G T W-3'	ImPyPyPyImHp-γ-PyPyHpHpImPy
	1618) 5'-W G C A A G A W-3'	ImPyPyPyImPy-γ-HpPyHpHpImPy
30	1619) 5'-W G C A A G G W-3'	ImPyPyPyImIm-γ-PyPyHpHpImPy
	1620) 5'-W G C A A G C W-3'	ImPyPyPyImPy-γ-ImPyHpHpImPy
	1621) 5'-W G C A A C T W-3'	ImPyPyPyPyHp-γ-PyImHpHpImPy
	1622) 5'-W G C A A C A W-3'	ImPyPyPyPyPy-γ-HpImHpHpImPy
	1623) 5'-W G C A A C G W-3'	ImPyPyPyPyIm-γ-PyImHpHpImPy
35	1624) 5'-W G C A A C C W-3'	ImPyPyPyPyPy-γ-ImImHpHpImPy

TABLE 97: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1625) 5'-W G C A G T T W-3'	ImPyPyImHpHp-γ-PyPyPyHpImPy
5	1626) 5'-W G C A G T A W-3'	ImPyPyImHpPy-γ-HpPyPyHpImPy
	1627) 5'-W G C A G T G W-3'	ImPyPyImHpIm-γ-PyPyPyHpImPy
	1628) 5'-W G C A G T C W-3'	ImPyPyImHpPy-γ-ImPyPyHpImPy
	1629) 5'-W G C A G A T W-3'	ImPyPyImPyHp-γ-PyHpPyHpImPy
	1630) 5'-W G C A G A A W-3'	ImPyPyImPyPy-γ-HpHpPyHpImPy
10	1631) 5'-W G C A G A G W-3'	ImPyPyImPyIm-γ-PyHpPyHpImPy
	1632) 5'-W G C A G A C W-3'	ImPyPyImPyPy-γ-ImHpPyHpImPy
	1633) 5'-W G C A G G T W-3'	ImPyPyImImHp-γ-PyPyPyHpImPy
	1634) 5'-W G C A G G A W-3'	ImPyPyImImPy-γ-HpPyPyHpImPy
	1635) 5'-W G C A G C T W-3'	ImPyPyImPyHp-γ-PyImPyHpImPy
	1636) 5'-W G C A G C A W-3'	ImPyPyImPyPy-γ-HpImPyHpImPy
	1637) 5'-W G C A G G G W-3'	ImPyPyImImIm-γ-PyPyPyHpImPy
	1638) 5'-W G C A G G C W-3'	ImPyPyImImPy-γ-ImPyPyHpImPy
	1639) 5'-W G C A G C G W-3'	ImPyPyImPyIm-γ-PyImPyHpImPy
	1640) 5'-W G C A G C C W-3'	ImPyPyImPyPy-γ-ImImPyHpImPy
20	1641) 5'-W G C A C T T W-3'	ImPyPyPyHpHp-γ-PyPyImHpImPy
	1642) 5'-W G C A C T A W-3'	ImPyPyPyHpPy-γ-HpPyImHpImPy
	1643) 5'-W G C A C T G W-3'	ImPyPyPyHpIm-γ-PyPyImHpImPy
	1644) 5'-W G C A C T C W-3'	ImPyPyPyHpPy-γ-ImPyImHpImPy
	1645) 5'-W G C A C A T W-3'	ImPyPyPyPyHp-γ-PyHpImHpImPy
25	1646) 5'-W G C A C A A W-3'	ImPyPyPyPyPy-γ-HpHpImHpImPy
	1647) 5'-W G C A C A G W-3'	ImPyPyPyPyIm-γ-PyHpImHpImPy
	1648) 5'-W G C A C A C W-3'	ImPyPyPyPyPy-γ-ImHpImHpImPy
	1649) 5'-W G C A C G T W-3'	ImPyPyPyImHp-γ-PyPyImHpImPy
	1650) 5'-W G C A C G A W-3'	ImPyPyPyImPy-γ-HpPyImHpImPy
30	1651) 5'-W G C A C C T W-3'	ImPyPyPyPyHp-γ-PyImImHpImPy
	1652) 5'-W G C A C C A W-3'	ImPyPyPyPyPy-γ-HpImImHpImPy
	1653) 5'-W G C A C G G W-3'	ImPyPyPyImIm-γ-PyPyImHpImPy
	1654) 5'-W G C A C G C W-3'	ImPyPyPyImPy-γ-ImPyImHpImPy
	1655) 5'-W G C A C C G W-3'	ImPyPyPyPyIm-γ-PyImImHpImPy
35	1656) 5'-W G C A C C C W-3'	ImPyPyPyPyPy-γ-ImImImHpImPy

TABLE 98: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1657) 5'-W G C C T T T W-3'	ImPyPyHpHpHp-γ-PyPyPyImImPy
5	1658) 5'-W G C C T T A W-3'	ImPyPyHpHpPy-γ-HpPyPyImImPy
	1659) 5'-W G C C T T G W-3'	ImPyPyHpHpIm-γ-PyPyPyImImPy
	1660) 5'-W G C C T T C W-3'	ImPyPyHpHpPy-γ-ImPyPyImImPy
	1661) 5'-W G C C T A T W-3'	ImPyPyHpPyHp-γ-PyHpPyImImPy
	1662) 5'-W G C C T A A W-3'	ImPyPyHpPyPy-γ-HpHpPyImImPy
10	1663) 5'-W G C C T A G W-3'	ImPyPyHpPyIm-γ-PyHpPyImImPy
	1664) 5'-W G C C T A C W-3'	ImPyPyHpPyPy-γ-ImHpPyImImPy
	1665) 5'-W G C C T G T W-3'	ImPyPyHpImHp-γ-PyPyPyImImPy
	1666) 5'-W G C C T G A W-3'	ImPyPyHpImPy-γ-HpPyPyImImPy
	1667) 5'-W G C C T G G W-3'	ImPyPyHpImIm-γ-PyPyPyImImPy
	1668) 5'-W G C C T G C W-3'	ImPyPyHpImPy-γ-ImPyPyImImPy
	1669) 5'-W G C C T C T W-3'	ImPyPyHpPyHp-γ-PyImPyImImPy
	1670) 5'-W G C C T C A W-3'	ImPyPyHpPyPy-γ-HpImPyImImPy
	1671) 5'-W G C C T C G W-3'	ImPyPyHpPyIm-γ-PyImPyImImPy
	1672) 5'-W G C C T C C W-3'	ImPyPyHpPyPy-γ-ImImPyImImPy
20	1673) 5'-W G C C A T T W-3'	ImPyPyPyHpHp-γ-PyPyHpImImPy
	1674) 5'-W G C C A T A W-3'	ImPyPyPyHpPy-γ-HpPyHpImImPy
	1675) 5'-W G C C A T G W-3'	ImPyPyPyHpIm-γ-PyPyHpImImPy
	1676) 5'-W G C C A T C W-3'	ImPyPyPyHpPy-γ-ImPyHpImImPy
	1677) 5'-W G C C A A T W-3'	ImPyPyPyPyHp-γ-PyHpHpImImPy
25	1678) 5'-W G C C A A A W-3'	ImPyPyPyPyPy-γ-HpHpHpImImPy
	1679) 5'-W G C C A A G W-3'	ImPyPyPyPyIm-γ-PyHpHpImImPy
	1680) 5'-W G C C A A C W-3'	ImPyPyPyPyPy-γ-ImHpHpImImPy
	1681) 5'-W G C C A G T W-3'	ImPyPyPyImHp-γ-PyPyHpImImPy
	1682) 5'-W G C C A G A W-3'	ImPyPyPyImPy-γ-HpPyHpImImPy
30	1683) 5'-W G C C A G G W-3'	ImPyPyPyImIm-γ-PyPyHpImImPy
	1684) 5'-W G C C A G C W-3'	ImPyPyPyImPy-γ-ImPyHpImImPy
	1685) 5'-W G C C A C T W-3'	ImPyPyPyPyHp-γ-PyImHpImImPy
	1686) 5'-W G C C A C A W-3'	ImPyPyPyPyPy-γ-HpImHpImImPy
	1687) 5'-W G C C A C G W-3'	ImPyPyPyPyIm-γ-PyImHpImImPy
35	1688) 5'-W G C C A C C W-3'	ImPyPyPyPyPy-γ-ImImHpImImPy

TABLE 99: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGCCSNW-3'

	DNA sequence	aromatic amino acid sequence
	1689) 5'-W G C C G T T W-3'	ImPyPyImHpHp- γ -PyPyPyImImPy
5	1690) 5'-W G C C G T A W-3'	ImPyPyImHpPy- γ -HpPyPyImImPy
	1691) 5'-W G C C G T G W-3'	ImPyPyImHpIm- γ -PyPyPyImImPy
	1692) 5'-W G C C G T C W-3'	ImPyPyImHpPy- γ -ImPyPyImImPy
	1693) 5'-W G C C G A T W-3'	ImPyPyImPyHp- γ -PyHpPyImImPy
	1694) 5'-W G C C G A A W-3'	ImPyPyImPyPy- γ -HpHpPyImImPy
10	1695) 5'-W G C C G A G W-3'	ImPyPyImPyIm- γ -PyHpPyImImPy
	1696) 5'-W G C C G A C W-3'	ImPyPyImPyPy- γ -ImHpPyImImPy
	1697) 5'-W G C C G G T W-3'	ImPyPyImImHp- γ -PyPyPyImImPy
	1698) 5'-W G C C G G A W-3'	ImPyPyImImPy- γ -HpPyPyImImPy
	1699) 5'-W G C C G C T W-3'	ImPyPyImPyHp- γ -PyImPyImImPy
	1700) 5'-W G C C G C A W-3'	ImPyPyImPyPy- γ -HpImPyImImPy
	1701) 5'-W G C C C T T W-3'	ImPyPyPyHpHp- γ -PyPyImImImPy
	1702) 5'-W G C C C T A W-3'	ImPyPyPyHpPy- γ -HpPyImImImPy
	1703) 5'-W G C C C T G W-3'	ImPyPyPyHpIm- γ -PyPyImImImPy
	1704) 5'-W G C C C T C W-3'	ImPyPyPyHpPy- γ -ImPyImImImPy
20	1705) 5'-W G C C C A T W-3'	ImPyPyPyPyHp- γ -PyHpImImImPy
	1706) 5'-W G C C C A A W-3'	ImPyPyPyPyPy- γ -HpHpImImImPy
	1707) 5'-W G C C C A G W-3'	ImPyPyPyPyIm- γ -PyHpImImImPy
	1708) 5'-W G C C C A C W-3'	ImPyPyPyPyPy- γ -ImHpImImImPy
	1709) 5'-W G C C C G T W-3'	ImPyPyPyImHp- γ -PyPyImImImPy
25	1710) 5'-W G C C C G A W-3'	ImPyPyPyImPy- γ -HpPyImImImPy
	1711) 5'-W G C C C C T W-3'	ImPyPyPyPyHp- γ -PyImImImImPy
	1712) 5'-W G C C C C A W-3'	ImPyPyPyPyPy- γ -HpImImImImPy
	G73) 5'-W G C C G G G W-3'	ImPyPyImImIm- γ -PyPyPyImImPy
	G74) 5'-W G C C G G C W-3'	ImPyPyImImPy- γ -ImPyPyImImPy
30	G75) 5'-W G C C G C G W-3'	ImPyPyImPyIm- γ -PyImPyImImPy
	G76) 5'-W G C C G C C W-3'	ImPyPyImPyPy- γ -ImImPyImImPy
	G77) 5'-W G C C C G G W-3'	ImPyPyPyImIm- γ -PyPyImImImPy
	G78) 5'-W G C C C G C W-3'	ImPyPyPyImPy- γ -ImPyImImImPy
	G79) 5'-W G C C C C G W-3'	ImPyPyPyPyIm- γ -PyImImImImPy
35	G80) 5'-W G C C C C C W-3'	ImPyPyPyPyPy- γ -ImImImImImPy

TABLE 100: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGAGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1713) 5'-W G A G T T T W-3'	ImPyImHpHpHp-γ-PyPyPyPyHpPy
5	1714) 5'-W G A G T T A W-3'	ImPyImHpHpPy-γ-HpPyPyPyHpPy
	1715) 5'-W G A G T T G W-3'	ImPyImHpHpIm-γ-PyPyPyPyHpPy
	1716) 5'-W G A G T T C W-3'	ImPyImHpHpPy-γ-ImPyPyPyHpPy
	1717) 5'-W G A G T A T W-3'	ImPyImHpPyHp-γ-PyHpPyPyHpPy
	1718) 5'-W G A G T A A W-3'	ImPyImHpPyPy-γ-HpHpPyPyHpPy
10	1719) 5'-W G A G T A G W-3'	ImPyImHpPyIm-γ-PyHpPyPyHpPy
	1720) 5'-W G A G T A C W-3'	ImPyImHpPyPy-γ-ImHpPyPyHpPy
	1721) 5'-W G A G T G T W-3'	ImPyImHpImHp-γ-PyPyPyPyHpPy
	1722) 5'-W G A G T G A W-3'	ImPyImHpImPy-γ-HpPyPyPyHpPy
	1723) 5'-W G A G T G G W-3'	ImPyImHpImIm-γ-PyPyPyPyHpPy
	1724) 5'-W G A G T G C W-3'	ImPyImHpImPy-γ-ImPyPyPyHpPy
	1725) 5'-W G A G T C T W-3'	ImPyImHpPyHp-γ-PyImPyPyHpPy
	1726) 5'-W G A G T C A W-3'	ImPyImHpPyPy-γ-HpImPyPyHpPy
	1727) 5'-W G A G T C G W-3'	ImPyImHpPyIm-γ-PyImPyPyHpPy
	1728) 5'-W G A G T C C W-3'	ImPyImHpPyPy-γ-ImImPyPyHpPy
20	1729) 5'-W G A G A T T W-3'	ImPyImPyHpHp-γ-PyPyHpPyHpPy
	1730) 5'-W G A G A T A W-3'	ImPyImPyHpPy-γ-HpPyHpPyHpPy
	1731) 5'-W G A G A T G W-3'	ImPyImPyHpIm-γ-PyPyHpPyHpPy
	1732) 5'-W G A G A T C W-3'	ImPyImPyHpPy-γ-ImPyHpPyHpPy
	1733) 5'-W G A G A A T W-3'	ImPyImPyPyHp-γ-PyHpHpPyHpPy
25	1734) 5'-W G A G A A A W-3'	ImPyImPyPyPy-γ-HpHpHpPyHpPy
	1735) 5'-W G A G A A G W-3'	ImPyImPyPyIm-γ-PyHpHpPyHpPy
	1736) 5'-W G A G A A C W-3'	ImPyImPyPyPy-γ-ImHpHpPyHpPy
	1737) 5'-W G A G A G T W-3'	ImPyImPyImHp-γ-PyPyHpPyHpPy
	1738) 5'-W G A G A G A W-3'	ImPyImPyImPy-γ-HpPyHpPyHpPy
30	1739) 5'-W G A G A G G W-3'	ImPyImPyImIm-γ-PyPyHpPyHpPy
	1740) 5'-W G A G A G C W-3'	ImPyImPyImPy-γ-ImPyHpPyHpPy
	1741) 5'-W G A G A C T W-3'	ImPyImPyPyHp-γ-PyImHpPyHpPy
	1742) 5'-W G A G A C A W-3'	ImPyImPyPyPy-γ-HpImHpPyHpPy
	1743) 5'-W G A G A C G W-3'	ImPyImPyPyIm-γ-PyImHpPyHpPy
35	1744) 5'-W G A G A C C W-3'	ImPyImPyPyPy-γ-ImImHpPyHpPy

TABLE 101: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGAGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1745) 5'-W G A G G T T W-3'	ImPyImImHpHp-γ-PyPyPyPyHpPy
5	1746) 5'-W G A G G T A W-3'	ImPyImImHpPy-γ-HpPyPyPyHpPy
	1747) 5'-W G A G G T G W-3'	ImPyImImHpIm-γ-PyPyPyPyHpPy
	1748) 5'-W G A G G T C W-3'	ImPyImImHpPy-γ-ImPyPyPyHpPy
	1749) 5'-W G A G G A T W-3'	ImPyImImPyHp-γ-PyHpPyPyHpPy
	1750) 5'-W G A G G A A W-3'	ImPyImImPyPy-γ-HpHpPyPyHpPy
10	1751) 5'-W G A G G A G W-3'	ImPyImImPyIm-γ-PyHpPyPyHpPy
	1752) 5'-W G A G G A C W-3'	ImPyImImPyPy-γ-ImHpPyPyHpPy
	1753) 5'-W G A G G G T W-3'	ImPyImImImHp-γ-PyPyPyPyHpPy
	1754) 5'-W G A G G G A W-3'	ImPyImImImPy-γ-HpPyPyPyHpPy
	1755) 5'-W G A G G C T W-3'	ImPyImImPyHp-γ-PyImPyPyHpPy
	1756) 5'-W G A G G C A W-3'	ImPyImImPyPy-γ-HpImPyPyHpPy
	1757) 5'-W G A G C T T W-3'	ImPyImPyHpHp-γ-PyPyImPyHpPy
	1758) 5'-W G A G C T A W-3'	ImPyImPyHpPy-γ-HpPyImPyHpPy
	1759) 5'-W G A G C T G W-3'	ImPyImPyHpIm-γ-PyPyImPyHpPy
	1760) 5'-W G A G C T C W-3'	ImPyImPyHpPy-γ-ImPyImPyHpPy
20	1761) 5'-W G A G C A T W-3'	ImPyImPyPyHp-γ-PyHpImPyHpPy
	1762) 5'-W G A G C A A W-3'	ImPyImPyPyPy-γ-HpHpImPyHpPy
	1763) 5'-W G A G C A G W-3'	ImPyImPyPyIm-γ-PyHpImPyHpPy
	1764) 5'-W G A G C A C W-3'	ImPyImPyPyPy-γ-ImHpImPyHpPy
	1765) 5'-W G A G C G T W-3'	ImPyImPyImHp-γ-PyPyImPyHpPy
25	1766) 5'-W G A G C G A W-3'	ImPyImPyImPy-γ-HpPyImPyHpPy
	1767) 5'-W G A G C C T W-3'	ImPyImPyPyHp-γ-PyImImPyHpPy
	1768) 5'-W G A G C C A W-3'	ImPyImPyPyPy-γ-HpImImPyHpPy
	1769) 5'-W G A G G G G W-3'	ImPyImImImIm-γ-PyPyPyPyHpPy
	1770) 5'-W G A G G G C W-3'	ImPyImImImPy-γ-ImPyPyPyHpPy
30	1771) 5'-W G A G G C G W-3'	ImPyImImPyIm-γ-PyImPyPyHpPy
	1772) 5'-W G A G G C C W-3'	ImPyImImPyPy-γ-ImImPyPyHpPy
	1773) 5'-W G A G C G G W-3'	ImPyImPyImIm-γ-PyPyImPyHpPy
	1774) 5'-W G A G C G C W-3'	ImPyImPyImPy-γ-ImPyImPyHpPy
	1775) 5'-W G A G C C G W-3'	ImPyImPyPyIm-γ-PyImImPyHpPy
35	1776) 5'-W G A G C C C W-3'	ImPyImPyPyPy-γ-ImImImPyHpPy

TABLE 102: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGATWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1777) 5'-W G A T T T T W-3'	ImPyHpHpHpHp- γ -PyPyPyPyHpPy
5	1778) 5'-W G A T T T A W-3'	ImPyHpHpHpPy- γ -HpPyPyPyHpPy
	1779) 5'-W G A T T T G W-3'	ImPyHpHpHpIm- γ -PyPyPyPyHpPy
	1780) 5'-W G A T T T C W-3'	ImPyHpHpHpPy- γ -ImPyPyPyHpPy
	1781) 5'-W G A T T A T W-3'	ImPyHpHpPyHp- γ -PyHpPyPyHpPy
	1782) 5'-W G A T T A A W-3'	ImPyHpHpPyPy- γ -HpHpPyPyHpPy
10	1783) 5'-W G A T T A G W-3'	ImPyHpHpPyIm- γ -PyHpPyPyHpPy
	1784) 5'-W G A T T A C W-3'	ImPyHpHpPyPy- γ -ImHpPyPyHpPy
	1785) 5'-W G A T T G T W-3'	ImPyHpHpImHp- γ -PyPyPyPyHpPy
	1786) 5'-W G A T T G A W-3'	ImPyHpHpImPy- γ -HpPyPyPyHpPy
	1787) 5'-W G A T T G G W-3'	ImPyHpHpImIm- γ -PyPyPyPyHpPy
	1788) 5'-W G A T T G C W-3'	ImPyHpHpImPy- γ -ImPyPyPyHpPy
	1789) 5'-W G A T T C T W-3'	ImPyHpHpPyHp- γ -PyImPyPyHpPy
	1790) 5'-W G A T T C A W-3'	ImPyHpHpPyPy- γ -HpImPyPyHpPy
	1791) 5'-W G A T T C G W-3'	ImPyHpHpPyIm- γ -PyImPyPyHpPy
	1792) 5'-W G A T T C C W-3'	ImPyHpHpPyPy- γ -ImImPyPyHpPy
	1793) 5'-W G A T A T T W-3'	ImPyHpPyHpHp- γ -PyPyHpPyHpPy
	1794) 5'-W G A T A T A W-3'	ImPyHpPyHpPy- γ -HpPyHpPyHpPy
	1795) 5'-W G A T A T G W-3'	ImPyHpPyHpIm- γ -PyPyHpPyHpPy
	1796) 5'-W G A T A T C W-3'	ImPyHpPyHpPy- γ -ImPyHpPyHpPy
	1797) 5'-W G A T A A T W-3'	ImPyHpPyPyHp- γ -PyHpHpPyHpPy
25	1798) 5'-W G A T A A A W-3'	ImPyHpPyPyPy- γ -HpHpHpPyHpPy
	1799) 5'-W G A T A A G W-3'	ImPyHpPyPyIm- γ -PyHpHpPyHpPy
	1800) 5'-W G A T A A C W-3'	ImPyHpPyPyPy- γ -ImHpHpPyHpPy
	1801) 5'-W G A T A G T W-3'	ImPyHpPyImHp- γ -PyPyHpPyHpPy
	1802) 5'-W G A T A G A W-3'	ImPyHpPyImPy- γ -HpPyHpPyHpPy
30	1803) 5'-W G A T A G G W-3'	ImPyHpPyImIm- γ -PyPyHpPyHpPy
	1804) 5'-W G A T A G C W-3'	ImPyHpPyImPy- γ -ImPyHpPyHpPy
	1805) 5'-W G A T A C T W-3'	ImPyHpPyPyHp- γ -PyImHpPyHpPy
	1806) 5'-W G A T A C A W-3'	ImPyHpPyPyPy- γ -HpImHpPyHpPy
	1807) 5'-W G A T A C G W-3'	ImPyHpPyPyIm- γ -PyImHpPyHpPy
35	1808) 5'-W G A T A C C W-3'	ImPyHpPyPyPy- γ -ImImHpPyHpPy

TABLE 103: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGATSNW-3'

	DNA sequence	aromatic amino acid sequence
	1809) 5'-W G A T G T T W-3'	ImPyHpImHpHp- γ -PyPyPyPyHpPy
5	1810) 5'-W G A T G T A W-3'	ImPyHpImHpPy- γ -HpPyPyPyHpPy
	1811) 5'-W G A T G T G W-3'	ImPyHpImHpIm- γ -PyPyPyPyHpPy
	1812) 5'-W G A T G T C W-3'	ImPyHpImHpPy- γ -ImPyPyPyHpPy
	1813) 5'-W G A T G A T W-3'	ImPyHpImPyHp- γ -PyHpPyPyHpPy
	1814) 5'-W G A T G A A W-3'	ImPyHpImPyPy- γ -HpHpPyPyHpPy
10	1815) 5'-W G A T G A G W-3'	ImPyHpImPyIm- γ -PyHpPyPyHpPy
	1816) 5'-W G A T G A C W-3'	ImPyHpImPyPy- γ -ImHpPyPyHpPy
	1817) 5'-W G A T G G T W-3'	ImPyHpImImHp- γ -PyPyPyPyHpPy
	1818) 5'-W G A T G G A W-3'	ImPyHpImImPy- γ -HpPyPyPyHpPy
	1819) 5'-W G A T G C T W-3'	ImPyHpImPyHp- γ -PyImPyPyHpPy
	1820) 5'-W G A T G C A W-3'	ImPyHpImPyPy- γ -HpImPyPyHpPy
	1821) 5'-W G A T G G G W-3'	ImPyHpImImIm- γ -PyPyPyPyHpPy
	1822) 5'-W G A T G G C W-3'	ImPyHpImImPy- γ -ImPyPyPyHpPy
	1823) 5'-W G A T G C G W-3'	ImPyHpImPyIm- γ -PyImPyPyHpPy
	1824) 5'-W G A T G C C W-3'	ImPyHpImPyPy- γ -ImImPyPyHpPy
20	1825) 5'-W G A T C T T W-3'	ImPyHpPyHpHp- γ -PyPyImPyHpPy
	1826) 5'-W G A T C T A W-3'	ImPyHpPyHpPy- γ -HpPyImPyHpPy
	1827) 5'-W G A T C T G W-3'	ImPyHpPyHpIm- γ -PyPyImPyHpPy
	1828) 5'-W G A T C T C W-3'	ImPyHpPyHpPy- γ -ImPyImPyHpPy
	1829) 5'-W G A T C A T W-3'	ImPyHpPyPyHp- γ -PyHpImPyHpPy
25	1830) 5'-W G A T C A A W-3'	ImPyHpPyPyPy- γ -HpHpImPyHpPy
	1831) 5'-W G A T C A G W-3'	ImPyHpPyPyIm- γ -PyHpImPyHpPy
	1832) 5'-W G A T C A C W-3'	ImPyHpPyPyPy- γ -ImHpImPyHpPy
	1833) 5'-W G A T C G T W-3'	ImPyHpPyImHp- γ -PyPyImPyHpPy
	1834) 5'-W G A T C G A W-3'	ImPyHpPyImPy- γ -HpPyImPyHpPy
30	1835) 5'-W G A T C C T W-3'	ImPyHpPyPyHp- γ -PyImImPyHpPy
	1836) 5'-W G A T C C A W-3'	ImPyHpPyPyPy- γ -HpImImPyHpPy
	1837) 5'-W G A T C G G W-3'	ImPyHpPyImIm- γ -PyPyImPyHpPy
	1838) 5'-W G A T C G C W-3'	ImPyHpPyImPy- γ -ImPyImPyHpPy
	1839) 5'-W G A T C C G W-3'	ImPyHpPyPyIm- γ -PyImImPyHpPy
35	1840) 5'-W G A T C C C W-3'	ImPyHpPyPyPy- γ -ImImImPyHpPy

TABLE 104: 12-ring Harpin Polyamides for recognition of 8-bp 5'-WGAAWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	1841) 5'-W G A A T T T W-3'	ImPyPyHpHpHp-γ-PyPyPyHpHpPy
	1842) 5'-W G A A T T A W-3'	ImPyPyHpHpPy-γ-HpPyPyHpHpPy
	1843) 5'-W G A A T T G W-3'	ImPyPyHpHpIm-γ-PyPyPyHpHpPy
	1844) 5'-W G A A T T C W-3'	ImPyPyHpHpPy-γ-ImPyPyHpHpPy
	1845) 5'-W G A A T A T W-3'	ImPyPyHpPyHp-γ-PyHpPyHpHpPy
10	1846) 5'-W G A A T A A W-3'	ImPyPyHpPyPy-γ-HpHpPyHpHpPy
	1847) 5'-W G A A T A G W-3'	ImPyPyHpPyIm-γ-PyHpPyHpHpPy
	1848) 5'-W G A A T A C W-3'	ImPyPyHpPyPy-γ-ImHpPyHpHpPy
	1849) 5'-W G A A T G T W-3'	ImPyPyHpImHp-γ-PyPyPyHpHpPy
	1850) 5'-W G A A T G A W-3'	ImPyPyHpImPy-γ-HpPyPyHpHpPy
	1851) 5'-W G A A T G G W-3'	ImPyPyHpImIm-γ-PyPyPyHpHpPy
	1852) 5'-W G A A T G C W-3'	ImPyPyHpImPy-γ-ImPyPyHpHpPy
	1853) 5'-W G A A T C T W-3'	ImPyPyHpPyHp-γ-PyImPyHpHpPy
	1854) 5'-W G A A T C A W-3'	ImPyPyHpPyPy-γ-HpImPyHpHpPy
	1855) 5'-W G A A T C G W-3'	ImPyPyHpPyIm-γ-PyImPyHpHpPy
20	1856) 5'-W G A A T C C W-3'	ImPyPyHpPyPy-γ-ImImPyHpHpPy
	1857) 5'-W G A A A T T W-3'	ImPyPyPyHpHp-γ-PyPyHpHpHpPy
	1858) 5'-W G A A A T A W-3'	ImPyPyPyHpPy-γ-HpPyHpHpHpPy
	1869) 5'-W G A A A T G W-3'	ImPyPyPyHpIm-γ-PyPyHpHpHpPy
	1860) 5'-W G A A A T C W-3'	ImPyPyPyHpPy-γ-ImPyHpHpHpPy
	1861) 5'-W G A A A A T W-3'	ImPyPyPyPyHp-γ-PyHpHpHpHpPy
25	1862) 5'-W G A A A A A W-3'	ImPyPyPyPyPy-γ-HpHpHpHpHpPy
	1863) 5'-W G A A A A G W-3'	ImPyPyPyPyIm-γ-PyHpHpHpHpPy
	1864) 5'-W G A A A A C W-3'	ImPyPyPyPyPy-γ-ImHpHpHpHpPy
	1865) 5'-W G A A A G T W-3'	ImPyPyPyImHp-γ-PyPyHpHpHpPy
	1866) 5'-W G A A A G A W-3'	ImPyPyPyImPy-γ-HpPyHpHpHpPy
30	1867) 5'-W G A A A G G W-3'	ImPyPyPyImIm-γ-PyPyHpHpHpPy
	1868) 5'-W G A A A G C W-3'	ImPyPyPyImPy-γ-ImPyHpHpHpPy
	1869) 5'-W G A A A C T W-3'	ImPyPyPyPyHp-γ-PyImHpHpHpPy
	1870) 5'-W G A A A C A W-3'	ImPyPyPyPyPy-γ-HpImHpHpHpPy
	1871) 5'-W G A A A C G W-3'	ImPyPyPyPyIm-γ-PyImHpHpHpPy
35	1872) 5'-W G A A A C C W-3'	ImPyPyPyPyPy-γ-ImImHpHpHpPy

TABLE 105: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGAASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1873) 5'-W G A A G T T W-3'	ImPyPyImHpHp- γ -PyPyPyHpHpPy
5	1874) 5'-W G A A G T A W-3'	ImPyPyImHpPy- γ -HpPyPyHpHpPy
	1875) 5'-W G A A G T G W-3'	ImPyPyImHpIm- γ -PyPyPyHpHpPy
	1876) 5'-W G A A G T C W-3'	ImPyPyImHpPy- γ -ImPyPyHpHpPy
	1877) 5'-W G A A G A T W-3'	ImPyPyImPyHp- γ -PyHpPyHpHpPy
	1878) 5'-W G A A G A A W-3'	ImPyPyImPyPy- γ -HpHpPyHpHpPy
10	1879) 5'-W G A A G A G W-3'	ImPyPyImPyIm- γ -PyHpPyHpHpPy
	1880) 5'-W G A A G A C W-3'	ImPyPyImPyPy- γ -ImHpPyHpHpPy
	1881) 5'-W G A A G G T W-3'	ImPyPyImImHp- γ -PyPyPyHpHpPy
	1882) 5'-W G A A G G A W-3'	ImPyPyImImPy- γ -HpPyPyHpHpPy
	1883) 5'-W G A A G C T W-3'	ImPyPyImPyHp- γ -PyImPyHpHpPy
	1884) 5'-W G A A G C A W-3'	ImPyPyImPyPy- γ -HpImPyHpHpPy
	1885) 5'-W G A A G G G W-3'	ImPyPyImImIm- γ -PyPyPyHpHpPy
	1886) 5'-W G A A G G C W-3'	ImPyPyImImPy- γ -ImPyPyHpHpPy
	1887) 5'-W G A A G C G W-3'	ImPyPyImPyIm- γ -PyImPyHpHpPy
	1888) 5'-W G A A G C C W-3'	ImPyPyImPyPy- γ -ImImPyHpHpPy
	1889) 5'-W G A A C T T W-3'	ImPyPyPyHpHp- γ -PyPyImHpHpPy
	1890) 5'-W G A A C T A W-3'	ImPyPyPyHpPy- γ -HpPyImHpHpPy
	1891) 5'-W G A A C T G W-3'	ImPyPyPyHpIm- γ -PyPyImHpHpPy
	1892) 5'-W G A A C T C W-3'	ImPyPyPyHpPy- γ -ImPyImHpHpPy
	1893) 5'-W G A A C A T W-3'	ImPyPyPyPyHp- γ -PyHpImHpHpPy
25	1894) 5'-W G A A C A A W-3'	ImPyPyPyPyPy- γ -HpHpImHpHpPy
	1895) 5'-W G A A C A G W-3'	ImPyPyPyPyIm- γ -PyHpImHpHpPy
	1896) 5'-W G A A C A C W-3'	ImPyPyPyPyPy- γ -ImHpImHpHpPy
	1897) 5'-W G A A C G T W-3'	ImPyPyPyImHp- γ -PyPyImHpHpPy
	1898) 5'-W G A A C G A W-3'	ImPyPyPyImPy- γ -HpPyImHpHpPy
30	1899) 5'-W G A A C C T W-3'	ImPyPyPyPyHp- γ -PyImImHpHpPy
	1900) 5'-W G A A C C A W-3'	ImPyPyPyPyPy- γ -HpImImHpHpPy
	1901) 5'-W G A A C G G W-3'	ImPyPyPyImIm- γ -PyPyImHpHpPy
	1902) 5'-W G A A C G C W-3'	ImPyPyPyImPy- γ -ImPyImHpHpPy
	1903) 5'-W G A A C C G W-3'	ImPyPyPyPyIm- γ -PyImImHpHpPy
35	1904) 5'-W G A A C C C W-3'	ImPyPyPyPyPy- γ -ImImImHpHpPy

TABLE 106: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGACWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1905) 5'-W G A C T T T W-3'	ImPyPyHpHpHp-γ-PyPyPyImHpPy
5	1906) 5'-W G A C T T A W-3'	ImPyPyHpHpPy-γ-HpPyPyImHpPy
	1907) 5'-W G A C T T G W-3'	ImPyPyHpHpIm-γ-PyPyPyImHpPy
	1908) 5'-W G A C T T C W-3'	ImPyPyHpHpPy-γ-ImPyPyImHpPy
	1909) 5'-W G A C T A T W-3'	ImPyPyHpPyHp-γ-PyHpPyImHpPy
	1910) 5'-W G A C T A A W-3'	ImPyPyHpPyPy-γ-HpHpPyImHpPy
10	1911) 5'-W G A C T A G W-3'	ImPyPyHpPyIm-γ-PyHpPyImHpPy
	1912) 5'-W G A C T A C W-3'	ImPyPyHpPyPy-γ-ImHpPyImHpPy
	1913) 5'-W G A C T G T W-3'	ImPyPyHpImHp-γ-PyPyPyImHpPy
	1914) 5'-W G A C T G A W-3'	ImPyPyHpImPy-γ-HpPyPyImHpPy
	1915) 5'-W G A C T G G W-3'	ImPyPyHpImIm-γ-PyPyPyImHpPy
	1916) 5'-W G A C T G C W-3'	ImPyPyHpImPy-γ-ImPyPyImHpPy
	1917) 5'-W G A C T C T W-3'	ImPyPyHpPyHp-γ-PyImPyImHpPy
	1918) 5'-W G A C T C A W-3'	ImPyPyHpPyPy-γ-HpImPyImHpPy
	1919) 5'-W G A C T C G W-3'	ImPyPyHpPyIm-γ-PyImPyImHpPy
	1920) 5'-W G A C T C C W-3'	ImPyPyHpPyPy-γ-ImImPyImHpPy
20	1921) 5'-W G A C A T T W-3'	ImPyPyPyHpHp-γ-PyPyPyImHpPy
	1922) 5'-W G A C A T A W-3'	ImPyPyPyHpPy-γ-HpPyPyImHpPy
	1923) 5'-W G A C A T G W-3'	ImPyPyPyHpIm-γ-PyPyPyImHpPy
	1924) 5'-W G A C A T C W-3'	ImPyPyPyHpPy-γ-ImPyPyImHpPy
	1925) 5'-W G A C A A T W-3'	ImPyPyPyPyHp-γ-PyHpPyImHpPy
25	1926) 5'-W G A C A A A W-3'	ImPyPyPyPyPy-γ-HpHpPyImHpPy
	1927) 5'-W G A C A A G W-3'	ImPyPyPyPyIm-γ-PyHpPyImHpPy
	1928) 5'-W G A C A A C W-3'	ImPyPyPyPyPy-γ-ImHpPyImHpPy
	1929) 5'-W G A C A G T W-3'	ImPyPyPyImHp-γ-PyPyPyImHpPy
	1930) 5'-W G A C A G A W-3'	ImPyPyPyImPy-γ-HpPyPyImHpPy
30	1931) 5'-W G A C A G G W-3'	ImPyPyPyImIm-γ-PyPyPyImHpPy
	1932) 5'-W G A C A G C W-3'	ImPyPyPyImPy-γ-ImPyPyImHpPy
	1933) 5'-W G A C A C T W-3'	ImPyPyPyPyHp-γ-PyImPyImHpPy
	1934) 5'-W G A C A C A W-3'	ImPyPyPyPyPy-γ-HpImPyImHpPy
	1935) 5'-W G A C A C G W-3'	ImPyPyPyPyIm-γ-PyImPyImHpPy
35	1936) 5'-W G A C A C C W-3'	ImPyPyPyPyPy-γ-ImImPyImHpPy

TABLE 107: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGACSNW-3'

	DNA sequence	aromatic amino acid sequence
	1937) 5'-W G A C G T T W-3'	ImPyPyImHpHp-γ-PyPyPyImHpPy
5	1938) 5'-W G A C G T A W-3'	ImPyPyImHpPy-γ-HpPyPyImHpPy
	1939) 5'-W G A C G T G W-3'	ImPyPyImHpIm-γ-PyPyPyImHpPy
	1940) 5'-W G A C G T C W-3'	ImPyPyImHpPy-γ-ImPyPyImHpPy
	1941) 5'-W G A C G A T W-3'	ImPyPyImPyHp-γ-PyHpPyImHpPy
	1942) 5'-W G A C G A A W-3'	ImPyPyImPyPy-γ-HpHpPyImHpPy
10	1943) 5'-W G A C G A G W-3'	ImPyPyImPyIm-γ-PyHpPyImHpPy
	1944) 5'-W G A C G A C W-3'	ImPyPyImPyPy-γ-ImHpPyImHpPy
	1945) 5'-W G A C G G T W-3'	ImPyPyImImHp-γ-PyPyPyImHpPy
	1946) 5'-W G A C G G A W-3'	ImPyPyImImPy-γ-HpPyPyImHpPy
	1947) 5'-W G A C G C T W-3'	ImPyPyImPyHp-γ-PyImPyImHpPy
	1948) 5'-W G A C G C A W-3'	ImPyPyImPyPy-γ-HpImPyImHpPy
	1949) 5'-W G A C C T T W-3'	ImPyPyPyHpHp-γ-PyPyImImHpPy
	1950) 5'-W G A C C T A W-3'	ImPyPyPyHpPy-γ-HpPyImImHpPy
	1951) 5'-W G A C C T G W-3'	ImPyPyPyHpIm-γ-PyPyImImHpPy
	1952) 5'-W G A C C T C W-3'	ImPyPyPyHpPy-γ-ImPyImImHpPy
20	1953) 5'-W G A C C A T W-3'	ImPyPyPyPyHp-γ-PyHpImImHpPy
	1954) 5'-W G A C C A A W-3'	ImPyPyPyPyPy-γ-HpHpImImHpPy
	1955) 5'-W G A C C A G W-3'	ImPyPyPyPyIm-γ-PyHpImImHpPy
	1956) 5'-W G A C C A C W-3'	ImPyPyPyPyPy-γ-ImHpImImHpPy
	1957) 5'-W G A C C G T W-3'	ImPyPyPyImHp-γ-PyPyImImHpPy
25	1958) 5'-W G A C C G A W-3'	ImPyPyPyImPy-γ-HpPyImImHpPy
	1959) 5'-W G A C C C T W-3'	ImPyPyPyPyHp-γ-PyImImImHpPy
	1960) 5'-W G A C C C A W-3'	ImPyPyPyPyPy-γ-HpImImImHpPy
	1961) 5'-W G A C G G G W-3'	ImPyPyImImIm-γ-PyPyPyImHpPy
	1962) 5'-W G A C G G C W-3'	ImPyPyImImPy-γ-ImPyPyImHpPy
30	1963) 5'-W G A C G C G W-3'	ImPyPyImPyIm-γ-PyImPyImHpPy
	1964) 5'-W G A C G C C W-3'	ImPyPyImPyPy-γ-ImImPyImHpPy
	1965) 5'-W G A C C G G W-3'	ImPyPyPyImIm-γ-PyPyImImHpPy
	1966) 5'-W G A C C G C W-3'	ImPyPyPyImPy-γ-ImPyImImHpPy
	1967) 5'-W G A C C C G W-3'	ImPyPyPyPyIm-γ-PyImImImHpPy
35	1968) 5'-W G A C C C C W-3'	ImPyPyPyPyPy-γ-ImImImImHpPy

TABLE 108: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1969) 5'-W G T G T T T W-3'	ImHpImHpHpHp-γ-PyPyPyPyPyPy
5	1970) 5'-W G T G T T A W-3'	ImHpImHpHpPy-γ-HpPyPyPyPyPy
	1971) 5'-W G T G T T G W-3'	ImHpImHpHpIm-γ-PyPyPyPyPyPy
	1972) 5'-W G T G T T C W-3'	ImHpImHpHpPy-γ-ImPyPyPyPyPy
	1973) 5'-W G T G T A T W-3'	ImHpImHpPyHp-γ-PyHpPyPyPyPy
	1974) 5'-W G T G T A A W-3'	ImHpImHpPyPy-γ-HpHpPyPyPyPy
10	1975) 5'-W G T G T A G W-3'	ImHpImHpPyIm-γ-PyHpPyPyPyPy
	1976) 5'-W G T G T A C W-3'	ImHpImHpPyPy-γ-ImHpPyPyPyPy
	1977) 5'-W G T G T G T W-3'	ImHpImHpImHp-γ-PyPyPyPyPyPy
	1978) 5'-W G T G T G A W-3'	ImHpImHpImPy-γ-HpPyPyPyPyPy
	1979) 5'-W G T G T G G W-3'	ImHpImHpImIm-γ-PyPyPyPyPyPy
	1980) 5'-W G T G T G C W-3'	ImHpImHpImPy-γ-ImPyPyPyPyPy
	1981) 5'-W G T G T C T W-3'	ImHpImHpPyHp-γ-PyImPyPyPyPy
	1982) 5'-W G T G T C A W-3'	ImHpImHpPyPy-γ-HpImPyPyPyPy
	1983) 5'-W G T G T C G W-3'	ImHpImHpPyIm-γ-PyImPyPyPyPy
	1984) 5'-W G T G T C C W-3'	ImHpImHpPyPy-γ-ImImPyPyPyPy
	1985) 5'-W G T G A T T W-3'	ImHpImPyHpHp-γ-PyPyHpPyPyPy
	1986) 5'-W G T G A T A W-3'	ImHpImPyHpPy-γ-HpPyHpPyPyPy
	1987) 5'-W G T G A T G W-3'	ImHpImPyHpIm-γ-PyPyHpPyPyPy
	1988) 5'-W G T G A T C W-3'	ImHpImPyHpPy-γ-ImPyHpPyPyPy
	1989) 5'-W G T G A A T W-3'	ImHpImPyPyHp-γ-PyHpHpPyPyPy
25	1990) 5'-W G T G A A A W-3'	ImHpImPyPyPy-γ-HpHpHpPyPyPy
	1991) 5'-W G T G A A G W-3'	ImHpImPyPyIm-γ-PyHpHpPyPyPy
	1992) 5'-W G T G A A C W-3'	ImHpImPyPyPy-γ-ImHpHpPyPyPy
	1993) 5'-W G T G A G T W-3'	ImHpImPyImHp-γ-PyPyHpPyPyPy
	1994) 5'-W G T G A G A W-3'	ImHpImPyImPy-γ-HpPyHpPyPyPy
30	1995) 5'-W G T G A G G W-3'	ImHpImPyImIm-γ-PyPyHpPyPyPy
	1996) 5'-W G T G A G C W-3'	ImHpImPyImPy-γ-ImPyHpPyPyPy
	1997) 5'-W G T G A C T W-3'	ImHpImPyPyHp-γ-PyImHpPyPyPy
	1998) 5'-W G T G A C A W-3'	ImHpImPyPyPy-γ-HpImHpPyPyPy
	1999) 5'-W G T G A C G W-3'	ImHpImPyPyIm-γ-PyImHpPyPyPy
35	2000) 5'-W G T G A C C W-3'	ImHpImPyPyPy-γ-ImImHpPyPyPy

TABLE 109: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2001) 5'-W G T G G T T W-3'	ImHpImImHpHp-γ-PyPyPyPyPyPy
5	2002) 5'-W G T G G T A W-3'	ImHpImImHpPy-γ-HpPyPyPyPyPy
	2003) 5'-W G T G G T G W-3'	ImHpImImHpIm-γ-PyPyPyPyPyPy
	2004) 5'-W G T G G T C W-3'	ImHpImImHpPy-γ-ImPyPyPyPyPy
	2005) 5'-W G T G G A T W-3'	ImHpImImPyHp-γ-PyHpPyPyPyPy
	2006) 5'-W G T G G A A W-3'	ImHpImImPyPy-γ-HpHpPyPyPyPy
10	2007) 5'-W G T G G A G W-3'	ImHpImImPyIm-γ-PyHpPyPyPyPy
	2008) 5'-W G T G G A C W-3'	ImHpImImPyPy-γ-ImHpPyPyPyPy
	2009) 5'-W G T G G G T W-3'	ImHpImImImHp-γ-PyPyPyPyPyPy
	2010) 5'-W G T G G G A W-3'	ImHpImImImPy-γ-HpPyPyPyPyPy
	2011) 5'-W G T G G C T W-3'	ImHpImImPyHp-γ-PyImPyPyPyPy
	2012) 5'-W G T G G C A W-3'	ImHpImImPyPy-γ-HpImPyPyPyPy
	2013) 5'-W G T G C T T W-3'	ImHpImPyHpHp-γ-PyPyImPyPyPy
	2014) 5'-W G T G C T A W-3'	ImHpImPyHpPy-γ-HpPyImPyPyPy
	2015) 5'-W G T G C T G W-3'	ImHpImPyHpIm-γ-PyPyImPyPyPy
	2016) 5'-W G T G C T C W-3'	ImHpImPyHpPy-γ-ImPyImPyPyPy
20	2017) 5'-W G T G C A T W-3'	ImHpImPyPyHp-γ-PyHpImPyPyPy
	2018) 5'-W G T G C A A W-3'	ImHpImPyPyPy-γ-HpHpImPyPyPy
	2019) 5'-W G T G C A G W-3'	ImHpImPyPyIm-γ-PyHpImPyPyPy
	2020) 5'-W G T G C A C W-3'	ImHpImPyPyPy-γ-ImHpImPyPyPy
	2021) 5'-W G T G C G T W-3'	ImHpImPyImHp-γ-PyPyImPyPyPy
25	2022) 5'-W G T G C G A W-3'	ImHpImPyImPy-γ-HpPyImPyPyPy
	2023) 5'-W G T G C C T W-3'	ImHpImPyPyHp-γ-PyImImPyPyPy
	2024) 5'-W G T G C C A W-3'	ImHpImPyPyPy-γ-HpImImPyPyPy
	2025) 5'-W G T G G G G W-3'	ImHpImImImIm-γ-PyPyPyPyPyPy
	2026) 5'-W G T G G G C W-3'	ImHpImImImPy-γ-ImPyPyPyPyPy
30	2027) 5'-W G T G G C G W-3'	ImHpImImPyIm-γ-PyImPyPyPyPy
	2028) 5'-W G T G G C C W-3'	ImHpImImPyPy-γ-ImImPyPyPyPy
	2029) 5'-W G T G C G G W-3'	ImHpImPyImIm-γ-PyPyImPyPyPy
	2030) 5'-W G T G C G C W-3'	ImHpImPyImPy-γ-ImPyImPyPyPy
	2031) 5'-W G T G C C G W-3'	ImHpImPyPyIm-γ-PyImImPyPyPy
35	2032) 5'-W G T G C C C W-3'	ImHpImPyPyPy-γ-ImImImPyPyPy

TABLE 110: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2033) 5'-W G T T T T W-3'	ImHpHpHpHp-γ-PyPyPyPyPyPy
5	2034) 5'-W G T T T T A W-3'	ImHpHpHpHpPy-γ-HpPyPyPyPyPy
	2035) 5'-W G T T T T G W-3'	ImHpHpHpHpIm-γ-PyPyPyPyPyPy
	2036) 5'-W G T T T T C W-3'	ImHpHpHpHpPy-γ-ImPyPyPyPyPy
	2037) 5'-W G T T T A T W-3'	ImHpHpHpPyHp-γ-PyHpPyPyPyPy
	2038) 5'-W G T T T A A W-3'	ImHpHpHpPyPy-γ-HpHpPyPyPyPy
10	2039) 5'-W G T T T A G W-3'	ImHpHpHpPyIm-γ-PyHpPyPyPyPy
	2040) 5'-W G T T T A C W-3'	ImHpHpHpPyPy-γ-ImHpPyPyPyPy
	2041) 5'-W G T T T G T W-3'	ImHpHpHpImHp-γ-PyPyPyPyPyPy
	2042) 5'-W G T T T G A W-3'	ImHpHpHpImPy-γ-HpPyPyPyPyPy
	2043) 5'-W G T T T G G W-3'	ImHpHpHpImIm-γ-PyPyPyPyPyPy
	2044) 5'-W G T T T G C W-3'	ImHpHpHpImPy-γ-ImPyPyPyPyPy
	2045) 5'-W G T T T C T W-3'	ImHpHpHpPyHp-γ-PyImPyPyPyPy
	2046) 5'-W G T T T C A W-3'	ImHpHpHpPyPy-γ-HpImPyPyPyPy
	2047) 5'-W G T T T C G W-3'	ImHpHpHpPyIm-γ-PyImPyPyPyPy
	2048) 5'-W G T T T C C W-3'	ImHpHpHpPyPy-γ-ImImPyPyPyPy
	2049) 5'-W G T T A T T W-3'	ImHpHpPyHpHp-γ-PyPyHpPyPyPy
	2050) 5'-W G T T A T A W-3'	ImHpHpPyHpPy-γ-HpPyHpPyPyPy
	2051) 5'-W G T T A T G W-3'	ImHpHpPyHpIm-γ-PyPyHpPyPyPy
	2052) 5'-W G T T A T C W-3'	ImHpHpPyHpPy-γ-ImPyHpPyPyPy
	2053) 5'-W G T T A A T W-3'	ImHpHpPyPyHp-γ-PyHpHpPyPyPy
25	2054) 5'-W G T T A A A W-3'	ImHpHpPyPyPy-γ-HpHpHpPyPyPy
	2055) 5'-W G T T A A G W-3'	ImHpHpPyPyIm-γ-PyHpHpPyPyPy
	2056) 5'-W G T T A A C W-3'	ImHpHpPyPyPy-γ-ImHpHpPyPyPy
	2057) 5'-W G T T A G T W-3'	ImHpHpPyImHp-γ-PyPyHpPyPyPy
	2058) 5'-W G T T A G A W-3'	ImHpHpPyImPy-γ-HpPyHpPyPyPy
30	2059) 5'-W G T T A G G W-3'	ImHpHpPyImIm-γ-PyPyHpPyPyPy
	2060) 5'-W G T T A G C W-3'	ImHpHpPyImPy-γ-ImPyHpPyPyPy
	2061) 5'-W G T T A C T W-3'	ImHpHpPyPyHp-γ-PyImHpPyPyPy
	2062) 5'-W G T T A C A W-3'	ImHpHpPyPyPy-γ-HpImHpPyPyPy
	2063) 5'-W G T T A C G W-3'	ImHpHpPyPyIm-γ-FyImHpPyPyPy
35	2064) 5'-W G T T A C C W-3'	ImHpHpPyPyPy-γ-ImImHpPyPyPy

TABLE 111: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2065) 5'-W G T T G T T W-3'	ImHpHpImHpHp-γ-PyPyPyPyPyPy
5	2066) 5'-W G T T G T A W-3'	ImHpHpImHpPy-γ-HpPyPyPyPyPy
	2067) 5'-W G T T G T G W-3'	ImHpHpImHpIm-γ-PyPyPyPyPyPy
	2068) 5'-W G T T G T C W-3'	ImHpHpImHpPy-γ-ImPyPyPyPyPy
	2069) 5'-W G T T G A T W-3'	ImHpHpImPyHp-γ-PyHpPyPyPyPy
	2070) 5'-W G T T G A A W-3'	ImHpHpImPyPy-γ-HpHpPyPyPyPy
10	2071) 5'-W G T T G A G W-3'	ImHpHpImPyIm-γ-PyHpPyPyPyPy
	2072) 5'-W G T T G A C W-3'	ImHpHpImPyPy-γ-ImHpPyPyPyPy
	2073) 5'-W G T T G G T W-3'	ImHpHpImImHp-γ-PyPyPyPyPyPy
	2074) 5'-W G T T G G A W-3'	ImHpHpImImPy-γ-HpPyPyPyPyPy
	2075) 5'-W G T T G C T W-3'	ImHpHpImPyHp-γ-PyImPyPyPyPy
	2076) 5'-W G T T G C A W-3'	ImHpHpImPyPy-γ-HpImPyPyPyPy
	2077) 5'-W G T T G G G W-3'	ImHpHpImImIm-γ-PyPyPyPyPyPy
	2078) 5'-W G T T G G C W-3'	ImHpHpImImPy-γ-ImPyPyPyPyPy
	2079) 5'-W G T T G C G W-3'	ImHpHpImPyIm-γ-PyImPyPyPyPy
	2080) 5'-W G T T G C C W-3'	ImHpHpImPyPy-γ-ImImPyPyPyPy
20	2081) 5'-W G T T C T T W-3'	ImHpHpPyHpHp-γ-PyPyImPyPyPy
	2082) 5'-W G T T C T A W-3'	ImHpHpPyHpPy-γ-HpPyImPyPyPy
	2083) 5'-W G T T C T G W-3'	ImHpHpPyHpIm-γ-PyPyImPyPyPy
	2084) 5'-W G T T C T C W-3'	ImHpHpPyHpPy-γ-ImPyImPyPyPy
	2085) 5'-W G T T C A T W-3'	ImHpHpPyPyHp-γ-PyHpImPyPyPy
25	2086) 5'-W G T T C A A W-3'	ImHpHpPyPyPy-γ-HpHpImPyPyPy
	2087) 5'-W G T T C A G W-3'	ImHpHpPyPyIm-γ-PyHpImPyPyPy
	2088) 5'-W G T T C A C W-3'	ImHpHpPyPyPy-γ-ImHpImPyPyPy
	2089) 5'-W G T T C G T W-3'	ImHpHpPyImHp-γ-PyPyImPyPyPy
	2090) 5'-W G T T C G A W-3'	ImHpHpPyImPy-γ-HpPyImPyPyPy
30	2091) 5'-W G T T C C T W-3'	ImHpHpPyPyHp-γ-PyImImPyPyPy
	2092) 5'-W G T T C C A W-3'	ImHpHpPyPyPy-γ-HpImImPyPyPy
	2093) 5'-W G T T C G G W-3'	ImHpHpPyImIm-γ-PyPyImPyPyPy
	2094) 5'-W G T T C G C W-3'	ImHpHpPyImPy-γ-ImPyImPyPyPy
	2095) 5'-W G T T C C G W-3'	ImHpHpPyPyIm-γ-PyImImPyPyPy
35	2096) 5'-W G T T C C C W-3'	ImHpHpPyPyPy-γ-ImImImPyPyPy

TABLE 112: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2097) 5'-W G T A T T T W-3'	ImHpPyHpHpHp-γ-PyPyPyHpPyPy
5	2098) 5'-W G T A T T A W-3'	ImHpPyHpHpPy-γ-HpPyPyHpPyPy
	2099) 5'-W G T A T T G W-3'	ImHpPyHpHpIm-γ-PyPyPyHpPyPy
	2100) 5'-W G T A T T C W-3'	ImHpPyHpHpPy-γ-ImPyPyHpPyPy
	2101) 5'-W G T A T A T W-3'	ImHpPyHpPyHp-γ-PyHpPyHpPyPy
	2102) 5'-W G T A T A A W-3'	ImHpPyHpPyPy-γ-HpHpPyHpPyPy
10	2103) 5'-W G T A T A G W-3'	ImHpPyHpPyIm-γ-PyHpPyHpPyPy
	2104) 5'-W G T A T A C W-3'	ImHpPyHpPyPy-γ-ImHpPyHpPyPy
	2105) 5'-W G T A T G T W-3'	ImHpPyHpImHp-γ-PyPyPyHpPyPy
	2106) 5'-W G T A T G A W-3'	ImHpPyHpImPy-γ-HpPyPyHpPyPy
	2107) 5'-W G T A T G G W-3'	ImHpPyHpImIm-γ-PyPyPyHpPyPy
15	2108) 5'-W G T A T G C W-3'	ImHpPyHpImPy-γ-ImPyPyHpPyPy
	2109) 5'-W G T A T C T W-3'	ImHpPyHpPyHp-γ-PyImPyHpPyPy
	2110) 5'-W G T A T C A W-3'	ImHpPyHpPyPy-γ-HpImPyHpPyPy
	2111) 5'-W G T A T C G W-3'	ImHpPyHpPyIm-γ-PyImPyHpPyPy
	2112) 5'-W G T A T C C W-3'	ImHpPyHpPyPy-γ-ImImPyHpPyPy
20	2113) 5'-W G T A A T T W-3'	ImHpPyPyHpHp-γ-PyPyHpHpPyPy
	2114) 5'-W G T A A T A W-3'	ImHpPyPyHpPy-γ-HpPyHpHpPyPy
	2115) 5'-W G T A A T G W-3'	ImHpPyPyHpIm-γ-PyPyHpHpPyPy
	2116) 5'-W G T A A T C W-3'	ImHpPyPyHpPy-γ-ImPyHpHpPyPy
	2117) 5'-W G T A A A T W-3'	ImHpPyPyPyHp-γ-PyHpHpHpPyPy
25	2118) 5'-W G T A A A A W-3'	ImHpPyPyPyPy-γ-HpHpHpHpPyPy
	2119) 5'-W G T A A A G W-3'	ImHpPyPyPyIm-γ-PyHpHpHpPyPy
	2120) 5'-W G T A A A C W-3'	ImHpPyPyPyPy-γ-ImHpHpHpPyPy
	2121) 5'-W G T A A G T W-3'	ImHpPyPyImHp-γ-PyPyHpHpPyPy
	2122) 5'-W G T A A G A W-3'	ImHpPyPyImPy-γ-HpPyHpHpPyPy
30	2123) 5'-W G T A A G G W-3'	ImHpPyPyImIm-γ-PyPyHpHpPyPy
	2124) 5'-W G T A A G C W-3'	ImHpPyPyImPy-γ-ImPyHpHpPyPy
	2125) 5'-W G T A A C T W-3'	ImHpPyPyPyHp-γ-PyImHpHpPyPy
	2126) 5'-W G T A A C A W-3'	ImHpPyPyPyPy-γ-HpImHpHpPyPy
	2127) 5'-W G T A A C G W-3'	ImHpPyPyPyIm-γ-PyImHpHpPyPy
35	2128) 5'-W G T A A C C W-3'	ImHpPyPyPyPy-γ-ImImHpHpPyPy

TABLE 113: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTASNNW-3'

	DNA sequence	aromatic amino acid sequence
	2129) 5'-W G T A G T T W-3'	ImHpPyImHpHp-γ-PyPyPyHpPyPy
5	2130) 5'-W G T A G T A W-3'	ImHpPyImHpPy-γ-HpPyPyHpPyPy
	2131) 5'-W G T A G T G W-3'	ImHpPyImHpIm-γ-PyPyPyHpPyPy
	2132) 5'-W G T A G T C W-3'	ImHpPyImHpPy-γ-ImPyPyHpPyPy
	2133) 5'-W G T A G A T W-3'	ImHpPyImPyHp-γ-PyHpPyHpPyPy
	2134) 5'-W G T A G A A W-3'	ImHpPyImPyPy-γ-HpHpPyHpPyPy
10	2135) 5'-W G T A G A G W-3'	ImHpPyImPyIm-γ-PyHpPyHpPyPy
	2136) 5'-W G T A G A C W-3'	ImHpPyImPyPy-γ-ImHpPyHpPyPy
	2137) 5'-W G T A G G T W-3'	ImHpPyImImHp-γ-PyPyPyHpPyPy
	2138) 5'-W G T A G G A W-3'	ImHpPyImImPy-γ-HpPyPyHpPyPy
	2139) 5'-W G T A G C T W-3'	ImHpPyImPyHp-γ-PyImPyHpPyPy
	2140) 5'-W G T A G C A W-3'	ImHpPyImPyPy-γ-HpImPyHpPyPy
	2141) 5'-W G T A G G G W-3'	ImHpPyImImIm-γ-PyPyPyHpPyPy
	2142) 5'-W G T A G G C W-3'	ImHpPyImImPy-γ-ImPyPyHpPyPy
	2143) 5'-W G T A G C G W-3'	ImHpPyImPyIm-γ-PyImPyHpPyPy
	2144) 5'-W G T A G C C W-3'	ImHpPyImPyPy-γ-ImImPyHpPyPy
20	2145) 5'-W G T A C T T W-3'	ImHpPyPyHpHp-γ-PyPyImHpPyPy
	2146) 5'-W G T A C T A W-3'	ImHpPyPyHpPy-γ-HpPyImHpPyPy
	2147) 5'-W G T A C T G W-3'	ImHpPyPyHpIm-γ-PyPyImHpPyPy
	2148) 5'-W G T A C T C W-3'	ImHpPyPyHpPy-γ-ImPyImHpPyPy
	2149) 5'-W G T A C A T W-3'	ImHpPyPyPyHp-γ-PyHpImHpPyPy
25	2150) 5'-W G T A C A A W-3'	ImHpPyPyPyPy-γ-HpHpImHpPyPy
	2151) 5'-W G T A C A G W-3'	ImHpPyPyPyIm-γ-PyHpImHpPyPy
	2152) 5'-W G T A C A C W-3'	ImHpPyPyPyPy-γ-ImHpImHpPyPy
	2153) 5'-W G T A C G T W-3'	ImHpPyPyImHp-γ-PyPyImHpPyPy
	2154) 5'-W G T A C G A W-3'	ImHpPyPyImPy-γ-HpPyImHpPyPy
30	2155) 5'-W G T A C C T W-3'	ImHpPyPyPyHp-γ-PyImImHpPyPy
	2156) 5'-W G T A C C A W-3'	ImHpPyPyPyPy-γ-HpImImHpPyPy
	2157) 5'-W G T A C G G W-3'	ImHpPyPyImIm-γ-PyPyImHpPyPy
	2158) 5'-W G T A C G C W-3'	ImHpPyPyImPy-γ-ImPyImHpPyPy
	2159) 5'-W G T A C C G W-3'	ImHpPyPyPyIm-γ-PyImImHpPyPy
35	2160) 5'-W G T A C C C W-3'	ImHpPyPyPyPy-γ-ImImImHpPyPy

TABLE 114: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2161) 5'-W G T C T T T W-3'	ImHpPyHpHpHp-γ-PyPyPyImPyPy
5	2162) 5'-W G T C T T A W-3'	ImHpPyHpHpPy-γ-HpPyPyImPyPy
	2163) 5'-W G T C T T G W-3'	ImHpPyHpHpIm-γ-PyPyPyImPyPy
	2164) 5'-W G T C T T C W-3'	ImHpPyHpHpPy-γ-ImPyPyImPyPy
	2165) 5'-W G T C T A T W-3'	ImHpPyHpPyHp-γ-PyHpPyImPyPy
	2166) 5'-W G T C T A A W-3'	ImHpPyHpPyPy-γ-HpHpPyImPyPy
10	2167) 5'-W G T C T A G W-3'	ImHpPyHpPyIm-γ-PyHpPyImPyPy
	2168) 5'-W G T C T A C W-3'	ImHpPyHpPyPy-γ-ImHpPyImPyPy
	2169) 5'-W G T C T G T W-3'	ImHpPyHpImHp-γ-PyPyPyImPyPy
	2170) 5'-W G T C T G A W-3'	ImHpPyHpImPy-γ-HpPyPyImPyPy
	2171) 5'-W G T C T G G W-3'	ImHpPyHpImIm-γ-PyPyPyImPyPy
	2172) 5'-W G T C T G C W-3'	ImHpPyHpImPy-γ-ImPyPyImPyPy
	2173) 5'-W G T C T C T W-3'	ImHpPyHpPyHp-γ-PyImPyImPyPy
	2174) 5'-W G T C T C A W-3'	ImHpPyHpPyPy-γ-HpImPyImPyPy
	2175) 5'-W G T C T C G W-3'	ImHpPyHpPyIm-γ-PyImPyImPyPy
	2176) 5'-W G T C T C C W-3'	ImHpPyHpPyPy-γ-ImImPyImPyPy
20	2177) 5'-W G T C A T T W-3'	ImHpPyPyHpHp-γ-PyPyHpImPyPy
	2178) 5'-W G T C A T A W-3'	ImHpPyPyHpPy-γ-HpPyHpImPyPy
	2179) 5'-W G T C A T G W-3'	ImHpPyPyHpIm-γ-PyPyHpImPyPy
	2180) 5'-W G T C A T C W-3'	ImHpPyPyHpPy-γ-ImPyHpImPyPy
	2181) 5'-W G T C A A T W-3'	ImHpPyPyPyHp-γ-PyHpHpImPyPy
25	2182) 5'-W G T C A A A W-3'	ImHpPyPyPyPy-γ-HpHpHpImPyPy
	2183) 5'-W G T C A A G W-3'	ImHpPyPyPyIm-γ-PyHpHpImPyPy
	2184) 5'-W G T C A A C W-3'	ImHpPyPyPyPy-γ-ImHpHpImPyPy
	2185) 5'-W G T C A G T W-3'	ImHpPyPyImHp-γ-PyPyHpImPyPy
	2186) 5'-W G T C A G A W-3'	ImHpPyPyImPy-γ-HpPyHpImPyPy
30	2187) 5'-W G T C A G G W-3'	ImHpPyPyImIm-γ-PyPyHpImPyPy
	2188) 5'-W G T C A G C W-3'	ImHpPyPyImPy-γ-ImPyHpImPyPy
	2189) 5'-W G T C A C T W-3'	ImHpPyPyPyHp-γ-PyImHpImPyPy
	2190) 5'-W G T C A C A W-3'	ImHpPyPyPyPy-γ-HpImHpImPyPy
	2191) 5'-W G T C A C G W-3'	ImHpPyPyPyIm-γ-PyImHpImPyPy
35	2192) 5'-W G T C A C C W-3'	ImHpPyPyPyPy-γ-ImImHpImPyPy

TABLE 115: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WGTCNNW-3'

	DNA sequence	aromatic amino acid sequence
	2193) 5'-W G T C G T T W-3'	ImHpPyImHpHp- γ -PyPyPyImPyPy
5	2194) 5'-W G T C G T A W-3'	ImHpPyImHpPy- γ -HpPyPyImPyPy
	2195) 5'-W G T C G T G W-3'	ImHpPyImHpIm- γ -PyPyPyImPyPy
	2196) 5'-W G T C G T C W-3'	ImHpPyImHpPy- γ -ImPyPyImPyPy
	2197) 5'-W G T C G A T W-3'	ImHpPyImPyHp- γ -PyHpPyImPyPy
	2198) 5'-W G T C G A A W-3'	ImHpPyImPyPy- γ -HpHpPyImPyPy
10	2199) 5'-W G T C G A G W-3'	ImHpPyImPyIm- γ -PyHpPyImPyPy
	2200) 5'-W G T C G A C W-3'	ImHpPyImPyPy- γ -ImHpPyImPyPy
	2201) 5'-W G T C G G T W-3'	ImHpPyImImHp- γ -PyPyPyImPyPy
	2202) 5'-W G T C G G A W-3'	ImHpPyImImPy- γ -HpPyPyImPyPy
	2203) 5'-W G T C G C T W-3'	ImHpPyImPyHp- γ -PyImPyImPyPy
	2204) 5'-W G T C G C A W-3'	ImHpPyImPyPy- γ -HpImPyImPyPy
	2205) 5'-W G T C C T T W-3'	ImHpPyPyHpHp- γ -PyPyImImPyPy
	2206) 5'-W G T C C T A W-3'	ImHpPyPyHpPy- γ -HpPyImImPyPy
	2207) 5'-W G T C C T G W-3'	ImHpPyPyHpIm- γ -PyPyImImPyPy
	2208) 5'-W G T C C T C W-3'	ImHpPyPyHpPy- γ -ImPyImImPyPy
	2209) 5'-W G T C C A T W-3'	ImHpPyPyPyHp- γ -PyHpImImPyPy
	2210) 5'-W G T C C A A W-3'	ImHpPyPyPyPy- γ -HpHpImImPyPy
	2211) 5'-W G T C C A G W-3'	ImHpPyPyPyIm- γ -PyHpImImPyPy
	2212) 5'-W G T C C A C W-3'	ImHpPyPyPyPy- γ -ImHpImImPyPy
	2213) 5'-W G T C C G T W-3'	ImHpPyPyImHp- γ -PyPyImImPyPy
25	2214) 5'-W G T C C G A W-3'	ImHpPyPyImPy- γ -HpPyImImPyPy
	2215) 5'-W G T C C C T W-3'	ImHpPyPyPyHp- γ -PyImImImPyPy
	2216) 5'-W G T C C C A W-3'	ImHpPyPyPyPy- γ -HpImImImPyPy
	2217) 5'-W G T C G G G W-3'	ImHpPyImImIm- γ -PyPyPyImPyPy
	2218) 5'-W G T C G G C W-3'	ImHpPyImImPy- γ -ImPyPyImPyPy
30	2219) 5'-W G T C G C G W-3'	ImHpPyImPyIm- γ -PyImPyImPyPy
	2220) 5'-W G T C G C C W-3'	ImHpPyImPyPy- γ -ImImPyImPyPy
	2221) 5'-W G T C C G G W-3'	ImHpPyPyImIm- γ -PyPyImImPyPy
	2222) 5'-W G T C C G C W-3'	ImHpPyPyImPy- γ -ImPyImImPyPy
	2223) 5'-W G T C C C G W-3'	ImHpPyPyPyIm- γ -PyImImImPyPy
35	2224) 5'-W G T C C C C W-3'	ImHpPyPyPyPy- γ -ImImImImPyPy

TABLE 116: 12-ring Hairpin Polyamides for recognition of 8-bp 5'WCGGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2225) 5'W C G G T T T W-3'	PyImImHpHpHp- γ -PyPyPyPyPyIm
	2226) 5'W C G G T T A W-3'	PyImImHpHpPy- γ -HpPyPyPyPyIm
5	2227) 5'W C G G T T G W-3'	PyImImHpHpIm- γ -PyPyPyPyPyIm
	2228) 5'W C G G T T C W-3'	PyImImHpHpPy- γ -ImPyPyPyPyIm
	2229) 5'W C G G T A T W-3'	PyImImHpPyHp- γ -PyHpPyPyPyIm
	2230) 5'W C G G T A A W-3'	PyImImHpPyPy- γ -HpHpPyPyPyIm
	2231) 5'W C G G T A G W-3'	PyImImHpPyIm- γ -PyHpPyPyPyIm
10	2232) 5'W C G G T A C W-3'	PyImImHpPyPy- γ -ImHpPyPyPyIm
	2233) 5'W C G G T G T W-3'	PyImImHpImHp- γ -PyPyPyPyPyIm
	2234) 5'W C G G T G A W-3'	PyImImHpImPy- γ -HpPyPyPyPyIm
	2235) 5'W C G G T G G W-3'	PyImImHpImIm- γ -PyPyPyPyPyIm
	2236) 5'W C G G T G C W-3'	PyImImHpImPy- γ -ImPyPyPyPyIm
	2237) 5'W C G G T C T W-3'	PyImImHpPyHp- γ -PyImPyPyPyIm
	2238) 5'W C G G T C A W-3'	PyImImHpPyPy- γ -HpImPyPyPyIm
	2239) 5'W C G G T C G W-3'	PyImImHpPyIm- γ -PyImPyPyPyIm
	2240) 5'W C G G T C C W-3'	PyImImHpPyPy- γ -ImImPyPyPyIm
	2241) 5'W C G G A T T W-3'	PyImImPyHpHp- γ -PyPyHpPyPyIm
	2242) 5'W C G G A T A W-3'	PyImImPyHpPy- γ -HpPyHpPyPyIm
	2243) 5'W C G G A T G W-3'	PyImImPyHpIm- γ -PyPyHpPyPyIm
	2244) 5'W C G G A T C W-3'	PyImImPyHpPy- γ -ImPyHpPyPyIm
	2245) 5'W C G G A A T W-3'	PyImImPyPyHp- γ -PyHpHpPyPyIm
	2246) 5'W C G G A A A W-3'	PyImImPyPyPy- γ -HpHpHpPyPyIm
25	2247) 5'W C G G A A G W-3'	PyImImPyPyIm- γ -PyHpHpPyPyIm
	2248) 5'W C G G A A C W-3'	PyImImPyPyPy- γ -ImHpHpPyPyIm
	2249) 5'W C G G A G T W-3'	PyImImPyImHp- γ -PyPyHpPyPyIm
	2250) 5'W C G G A G A W-3'	PyImImPyImPy- γ -HpPyHpPyPyIm
	2251) 5'W C G G A G G W-3'	PyImImPyImIm- γ -PyPyHpPyPyIm
30	2252) 5'W C G G A G C W-3'	PyImImPyImPy- γ -ImPyHpPyPyIm
	2253) 5'W C G G A C T W-3'	PyImImPyPyHp- γ -PyImHpPyPyIm
	2254) 5'W C G G A C A W-3'	PyImImPyPyPy- γ -HpImHpPyPyIm
	2255) 5'W C G G A C G W-3'	PyImImPyPyIm- γ -PyImHpPyPyIm
	2256) 5'W C G G A C C W-3'	PyImImPyPyPy- γ -ImImHpPyPyIm

TABLE 117: 12-ring Hairpin Polyamides for recognition of 8-bp 5'WCGGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2257) 5'W C G G G T T W-3'	PyImImImHpHp- γ -PyPyPyPyPyIm
5	2258) 5'W C G G G T A W-3'	PyImImImHpPy- γ -HpPyPyPyPyIm
	2259) 5'W C G G G T G W-3'	PyImImImHpIm- γ -PyPyPyPyPyIm
	2260) 5'W C G G G T C W-3'	PyImImImHpPy- γ -ImPyPyPyPyIm
	2261) 5'W C G G G A T W-3'	PyImImImPyHp- γ -PyHpPyPyPyIm
	2262) 5'W C G G G A A W-3'	PyImImImPyPy- γ -HpHpPyPyPyIm
10	2263) 5'W C G G G A G W-3'	PyImImImPyIm- γ -PyHpPyPyPyIm
	2264) 5'W C G G G A C W-3'	PyImImImPyPy- γ -ImHpPyPyPyIm
	2265) 5'W C G G G G T W-3'	PyImImImImHp- γ -PyPyPyPyPyIm
	2266) 5'W C G G G G A W-3'	PyImImImImPy- γ -HpPyPyPyPyIm
	2267) 5'W C G G G C T W-3'	PyImImImPyHp- γ -PyImPyPyPyIm
	2268) 5'W C G G G C A W-3'	PyImImImPyPy- γ -HpImPyPyPyIm
	2269) 5'W C G G C T T W-3'	PyImImPyHpHp- γ -PyPyImPyPyIm
	2270) 5'W C G G C T A W-3'	PyImImPyHpPy- γ -HpPyImPyPyIm
	2271) 5'W C G G C T G W-3'	PyImImPyHpIm- γ -PyPyImPyPyIm
	2272) 5'W C G G C T C W-3'	PyImImPyHpPy- γ -ImPyImPyPyIm
	2273) 5'W C G G C A T W-3'	PyImImPyPyHp- γ -PyHpImPyPyIm
	2274) 5'W C G G C A A W-3'	PyImImPyPyPy- γ -HpHpImPyPyIm
	2275) 5'W C G G C A G W-3'	PyImImPyPyIm- γ -PyHpImPyPyIm
	2276) 5'W C G G C A C W-3'	PyImImPyPyPy- γ -ImHpImPyPyIm
	2277) 5'W C G G C G T W-3'	PyImImPyImHp- γ -PyPyImPyPyIm
25	2278) 5'W C G G C G A W-3'	PyImImPyImPy- γ -HpPyImPyPyIm
	2279) 5'W C G G C C T W-3'	PyImImPyPyHp- γ -PyImImPyPyIm
	2280) 5'W C G G C C A W-3'	PyImImPyPyPy- γ -HpImImPyPyIm
	G83) 5'W C G G G G G W-3'	PyImImImImIm- γ -PyPyPyPyPyIm
	G84) 5'W C G G G G C W-3'	PyImImImImPy- γ -ImPyPyPyPyIm
30	G85) 5'W C G G G C G W-3'	PyImImImPyIm- γ -PyImPyPyPyIm
	G86) 5'W C G G G C C W-3'	PyImImImPyPy- γ -ImImPyPyPyIm
	G87) 5'W C G G C G G W-3'	PyImImPyImIm- γ -PyPyImPyPyIm
	G88) 5'W C G G C G C W-3'	PyImImPyImPy- γ -ImPyImPyPyIm
	G89) 5'W C G G C C G W-3'	PyImImPyPyIm- γ -PyImImPyPyIm
35	G90) 5'W C G G C C C W-3'	PyImImPyPyPy- γ -ImImImPyPyIm

TABLE 118: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2281) 5'W C G T T T T W-3'	PyImHpHpHpHp-γ-PyPyPyPyPyIm
5	2282) 5'W C G T T T A W-3'	PyImHpHpHpPy-γ-HpPyPyPyPyIm
	2283) 5'W C G T T T G W-3'	PyImHpHpHpIm-γ-PyPyPyPyPyIm
	2284) 5'W C G T T T C W-3'	PyImHpHpHpPy-γ-ImPyPyPyPyIm
	2285) 5'W C G T T A T W-3'	PyImHpHpPyHp-γ-PyHpPyPyPyIm
	2286) 5'W C G T T A A W-3'	PyImHpHpPyPy-γ-HpHpPyPyPyIm
10	2287) 5'W C G T T A G W-3'	PyImHpHpPyIm-γ-PyHpPyPyPyIm
	2288) 5'W C G T T A C W-3'	PyImHpHpPyPy-γ-ImHpPyPyPyIm
	2289) 5'W C G T T G T W-3'	PyImHpHpImHp-γ-PyPyPyPyPyIm
	2290) 5'W C G T T G A W-3'	PyImHpHpImPy-γ-HpPyPyPyPyIm
	2291) 5'W C G T T G G W-3'	PyImHpHpImIm-γ-PyPyPyPyPyIm
	2292) 5'W C G T T G C W-3'	PyImHpHpImPy-γ-ImPyPyPyPyIm
	2293) 5'W C G T T C T W-3'	PyImHpHpPyHp-γ-PyImPyPyPyIm
	2294) 5'W C G T T C A W-3'	PyImHpHpPyPy-γ-HpImPyPyPyIm
	2295) 5'W C G T T C G W-3'	PyImHpHpPyIm-γ-PyImPyPyPyIm
	2296) 5'W C G T T C C W-3'	PyImHpHpPyPy-γ-ImImPyPyPyIm
20	2297) 5'W C G T A T T W-3'	PyImHpPyHpHp-γ-PyPyHpPyPyIm
	2298) 5'W C G T A T A W-3'	PyImHpPyHpPy-γ-HpPyHpPyPyIm
	2299) 5'W C G T A T G W-3'	PyImHpPyHpIm-γ-PyPyHpPyPyIm
	2300) 5'W C G T A T C W-3'	PyImHpPyHpPy-γ-ImPyHpPyPyIm
	2301) 5'W C G T A A T W-3'	PyImHpPyPyHp-γ-PyHpHpPyPyIm
25	2302) 5'W C G T A A A W-3'	PyImHpPyPyPy-γ-HpHpHpPyPyIm
	2303) 5'W C G T A A G W-3'	PyImHpPyPyIm-γ-PyHpHpPyPyIm
	2304) 5'W C G T A A C W-3'	PyImHpPyPyPy-γ-ImHpHpPyPyIm
	2305) 5'W C G T A G T W-3'	PyImHpPyImHp-γ-PyPyHpPyPyIm
	2306) 5'W C G T A G A W-3'	PyImHpPyImPy-γ-HpPyHpPyPyIm
30	2307) 5'W C G T A G G W-3'	PyImHpPyImIm-γ-PyPyHpPyPyIm
	2308) 5'W C G T A G C W-3'	PyImHpPyImPy-γ-ImPyHpPyPyIm
	2309) 5'W C G T A C T W-3'	PyImHpPyPyHp-γ-PyImHpPyPyIm
	2310) 5'W C G T A C A W-3'	PyImHpPyPyPy-γ-HpImHpPyPyIm
	2311) 5'W C G T A C G W-3'	PyImHpPyPyIm-γ-PyImHpPyPyIm
35	2312) 5'W C G T A C C W-3'	PyImHpPyPyPy-γ-ImImHpPyPyIm

TABLE 119: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2313) 5'W C G T G T T W-3'	PyImHpImHpHp- γ -PyPyPyPyPyIm
5	2314) 5'W C G T G T A W-3'	PyImHpImHpPy- γ -HpPyPyPyPyIm
	2315) 5'W C G T G T G W-3'	PyImHpImHpIm- γ -PyPyPyPyPyIm
	2316) 5'W C G T G T C W-3'	PyImHpImHpPy- γ -ImPyPyPyPyIm
	2317) 5'W C G T G A T W-3'	PyImHpImPyHp- γ -PyHpPyPyPyIm
	2318) 5'W C G T G A A W-3'	PyImHpImPyPy- γ -HpHpPyPyPyIm
10	2319) 5'W C G T G A G W-3'	PyImHpImPyIm- γ -PyHpPyPyPyIm
	2320) 5'W C G T G A C W-3'	PyImHpImPyPy- γ -ImHpPyPyPyIm
	2321) 5'W C G T G G T W-3'	PyImHpImImHp- γ -PyPyPyPyPyIm
	2322) 5'W C G T G G A W-3'	PyImHpImImPy- γ -HpPyPyPyPyIm
	2323) 5'W C G T G C T W-3'	PyImHpImPyHp- γ -PyImPyPyPyIm
	2324) 5'W C G T G C A W-3'	PyImHpImPyPy- γ -HpImPyPyPyIm
	2325) 5'W C G T G G G W-3'	PyImHpImImIm- γ -PyPyPyPyPyIm
	2326) 5'W C G T G G C W-3'	PyImHpImImPy- γ -ImPyPyPyPyIm
	2327) 5'W C G T G C G W-3'	PyImHpImPyIm- γ -PyImPyPyPyIm
	2328) 5'W C G T G C C W-3'	PyImHpImPyPy- γ -ImImPyPyPyIm
	2329) 5'W C G T C T T W-3'	PyImHpPyHpHp- γ -PyPyImPyPyIm
	2330) 5'W C G T C T A W-3'	PyImHpPyHpPy- γ -HpPyImPyPyIm
	2331) 5'W C G T C T G W-3'	PyImHpPyHpIm- γ -PyPyImPyPyIm
	2332) 5'W C G T C T C W-3'	PyImHpPyHpPy- γ -ImPyImPyPyIm
	2333) 5'W C G T C A T W-3'	PyImHpPyPyHp- γ -PyHpImPyPyIm
25	2334) 5'W C G T C A A W-3'	PyImHpPyPyPy- γ -HpHpImPyPyIm
	2335) 5'W C G T C A G W-3'	PyImHpPyPyIm- γ -PyHpImPyPyIm
	2336) 5'W C G T C A C W-3'	PyImHpPyPyPy- γ -ImHpImPyPyIm
	2337) 5'W C G T C G T W-3'	PyImHpPyImHp- γ -PyPyImPyPyIm
	2338) 5'W C G T C G A W-3'	PyImHpPyImPy- γ -HpPyImPyPyIm
30	2339) 5'W C G T C C T W-3'	PyImHpPyPyHp- γ -PyImImPyPyIm
	2340) 5'W C G T C C A W-3'	PyImHpPyPyPy- γ -HpImImPyPyIm
	2341) 5'W C G T C G G W-3'	PyImHpPyImIm- γ -PyPyImPyPyIm
	2342) 5'W C G T C G C W-3'	PyImHpPyImPy- γ -ImPyImPyPyIm
	2343) 5'W C G T C C G W-3'	PyImHpPyPyIm- γ -PyImImPyPyIm
35	2344) 5'W C G T C C C W-3'	PyImHpPyPyPy- γ -ImImImPyPyIm

TABLE 120: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2345) 5'W C G A T T T W-3'	PyImPyHpHpHp-γ-PyPyPyHpPyIm
5	2346) 5'W C G A T T A W-3'	PyImPyHpHpPy-γ-HpPyPyHpPyIm
	2347) 5'W C G A T T G W-3'	PyImPyHpHpIm-γ-PyPyPyHpPyIm
	2348) 5'W C G A T T C W-3'	PyImPyHpHpPy-γ-ImPyPyHpPyIm
	2349) 5'W C G A T A T W-3'	PyImPyHpPyHp-γ-PyHpPyHpPyIm
	2350) 5'W C G A T A A W-3'	PyImPyHpPyPy-γ-HpHpPyHpPyIm
10	2351) 5'W C G A T A G W-3'	PyImPyHpPyIm-γ-PyHpPyHpPyIm
	2352) 5'W C G A T A C W-3'	PyImPyHpPyPy-γ-ImHpPyHpPyIm
	2353) 5'W C G A T G T W-3'	PyImPyHpImHp-γ-PyPyPyHpPyIm
	2354) 5'W C G A T G A W-3'	PyImPyHpImPy-γ-HpPyPyHpPyIm
	2355) 5'W C G A T G G W-3'	PyImPyHpImIm-γ-PyPyPyHpPyIm
	2356) 5'W C G A T G C W-3'	PyImPyHpImPy-γ-ImPyPyHpPyIm
	2357) 5'W C G A T C T W-3'	PyImPyHpPyHp-γ-PyImPyHpPyIm
	2358) 5'W C G A T C A W-3'	PyImPyHpPyPy-γ-HpImPyHpPyIm
	2359) 5'W C G A T C G W-3'	PyImPyHpPyIm-γ-PyImPyHpPyIm
	2360) 5'W C G A T C C W-3'	PyImPyHpPyPy-γ-ImImPyHpPyIm
20	2361) 5'W C G A A T T W-3'	PyImPyPyHpHp-γ-PyPyHpHpPyIm
	2362) 5'W C G A A T A W-3'	PyImPyPyHpPy-γ-HpPyHpHpPyIm
	2363) 5'W C G A A T G W-3'	PyImPyPyHpIm-γ-PyPyHpHpPyIm
	2364) 5'W C G A A T C W-3'	PyImPyPyHpPy-γ-ImPyHpHpPyIm
	2365) 5'W C G A A A T W-3'	PyImPyPyPyHp-γ-PyHpHpHpPyIm
25	2366) 5'W C G A A A A W-3'	PyImPyPyPyPy-γ-HpHpHpHpPyIm
	2367) 5'W C G A A A G W-3'	PyImPyPyPyIm-γ-PyHpHpHpPyIm
	2368) 5'W C G A A A C W-3'	PyImPyPyPyPy-γ-ImHpHpHpPyIm
	2369) 5'W C G A A G T W-3'	PyImPyPyImHp-γ-PyPyHpHpPyIm
	2370) 5'W C G A A G A W-3'	PyImPyPyImPy-γ-HpPyHpHpPyIm
30	2371) 5'W C G A A G G W-3'	PyImPyPyImIm-γ-PyPyHpHpPyIm
	2372) 5'W C G A A G C W-3'	PyImPyPyImPy-γ-ImPyHpHpPyIm
	2373) 5'W C G A A C T W-3'	PyImPyPyPyHp-γ-PyImHpHpPyIm
	2374) 5'W C G A A C A W-3'	PyImPyPyPyPy-γ-HpImHpHpPyIm
	2375) 5'W C G A A C G W-3'	PyImPyPyPyIm-γ-PyImHpHpPyIm
35	2376) 5'W C G A A C C W-3'	PyImPyPyPyPy-γ-ImImHpHpPyIm

TABLE 121: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGASNNW-3'

	DNA sequence	aromatic amino acid sequence
	2377) 5'W C G A G T T W-3'	PyImPyImHpHp- γ -PyPyPyHpPyIm
5	2378) 5'W C G A G T A W-3'	PyImPyImHpPy- γ -HpPyPyHpPyIm
	2379) 5'W C G A G T G W-3'	PyImPyImHpIm- γ -PyPyPyHpPyIm
	2380) 5'W C G A G T C W-3'	PyImPyImHpPy- γ -ImPyPyHpPyIm
	2381) 5'W C G A G A T W-3'	PyImPyImPyHp- γ -PyHpPyHpPyIm
	2382) 5'W C G A G A A W-3'	PyImPyImPyPy- γ -HpHpPyHpPyIm
10	2383) 5'W C G A G A G W-3'	PyImPyImPyIm- γ -PyHpPyHpPyIm
	2384) 5'W C G A G A C W-3'	PyImPyImPyPy- γ -ImHpPyHpPyIm
	2385) 5'W C G A G G T W-3'	PyImPyImImHp- γ -PyPyPyHpPyIm
	2386) 5'W C G A G G A W-3'	PyImPyImImPy- γ -HpPyPyHpPyIm
	2387) 5'W C G A G C T W-3'	PyImPyImPyHp- γ -PyImPyHpPyIm
	2388) 5'W C G A G C A W-3'	PyImPyImPyPy- γ -HpImPyHpPyIm
	2389) 5'W C G A G G G W-3'	PyImPyImImIm- γ -PyPyPyHpPyIm
	2390) 5'W C G A G G C W-3'	PyImPyImImPy- γ -ImPyPyHpPyIm
	2391) 5'W C G A G C G W-3'	PyImPyImPyIm- γ -PyImPyHpPyIm
	2392) 5'W C G A G C C W-3'	PyImPyImPyPy- γ -ImImPyHpPyIm
	2393) 5'W C G A C T T W-3'	PyImPyPyHpHp- γ -PyPyImHpPyIm
	2394) 5'W C G A C T A W-3'	PyImPyPyHpPy- γ -HpPyImHpPyIm
	2395) 5'W C G A C T G W-3'	PyImPyPyHpIm- γ -PyPyImHpPyIm
	2396) 5'W C G A C T C W-3'	PyImPyPyHpPy- γ -ImPyImHpPyIm
	2397) 5'W C G A C A T W-3'	PyImPyPyPyHp- γ -PyHpImHpPyIm
25	2398) 5'W C G A C A A W-3'	PyImPyPyPyPy- γ -HpHpImHpPyIm
	2399) 5'W C G A C A G W-3'	PyImPyPyPyIm- γ -PyHpImHpPyIm
	2400) 5'W C G A C A C W-3'	PyImPyPyPyPy- γ -ImHpImHpPyIm
	2401) 5'W C G A C G T W-3'	PyImPyPyImHp- γ -PyPyImHpPyIm
	2402) 5'W C G A C G A W-3'	PyImPyPyImPy- γ -HpPyImHpPyIm
30	2403) 5'W C G A C C T W-3'	PyImPyPyPyHp- γ -PyImImHpPyIm
	2404) 5'W C G A C C A W-3'	PyImPyPyPyPy- γ -HpImImHpPyIm
	2405) 5'W C G A C G G W-3'	PyImPyPyImIm- γ -PyPyImHpPyIm
	2406) 5'W C G A C G C W-3'	PyImPyPyImPy- γ -ImPyImHpPyIm
	2407) 5'W C G A C C G W-3'	PyImPyPyPyIm- γ -PyImImHpPyIm
35	2408) 5'W C G A C C C W-3'	PyImPyPyPyPy- γ -ImImImHpPyIm

TABLE 122: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2409) 5'W C G C T T T W-3'	PyImPyHpHpHp-γ-PyPyPyImPyIm
5	2410) 5'W C G C T T A W-3'	PyImPyHpHpPy-γ-HpPyPyImPyIm
	2411) 5'W C G C T T G W-3'	PyImPyHpHpIm-γ-PyPyPyImPyIm
	2412) 5'W C G C T T C W-3'	PyImPyHpHpPy-γ-ImPyPyImPyIm
	2413) 5'W C G C T A T W-3'	PyImPyHpPyHp-γ-PyHpPyImPyIm
	2414) 5'W C G C T A A W-3'	PyImPyHpPyPy-γ-HpHpPyImPyIm
10	2415) 5'W C G C T A G W-3'	PyImPyHpPyIm-γ-PyHpPyImPyIm
	2416) 5'W C G C T A C W-3'	PyImPyHpPyPy-γ-ImHpPyImPyIm
	2417) 5'W C G C T G T W-3'	PyImPyHpImHp-γ-PyPyPyImPyIm
	2418) 5'W C G C T G A W-3'	PyImPyHpImPy-γ-HpPyPyImPyIm
	2419) 5'W C G C T G G W-3'	PyImPyHpImIm-γ-PyPyPyImPyIm
	2420) 5'W C G C T G C W-3'	PyImPyHpImPy-γ-ImPyPyImPyIm
	2421) 5'W C G C T C T W-3'	PyImPyHpPyHp-γ-PyImPyImPyIm
	2422) 5'W C G C T C A W-3'	PyImPyHpPyPy-γ-HpImPyImPyIm
	2423) 5'W C G C T C G W-3'	PyImPyHpPyIm-γ-PyImPyImPyIm
	2424) 5'W C G C T C C W-3'	PyImPyHpPyPy-γ-ImImPyImPyIm
20	2425) 5'W C G C A T T W-3'	PyImPyPyHpHp-γ-PyPyHpImPyIm
	2426) 5'W C G C A T A W-3'	PyImPyPyHpPy-γ-HpPyHpImPyIm
	2427) 5'W C G C A T G W-3'	PyImPyPyHpIm-γ-PyPyHpImPyIm
	2428) 5'W C G C A T C W-3'	PyImPyPyHpPy-γ-ImPyHpImPyIm
	2429) 5'W C G C A A T W-3'	PyImPyPyPyHp-γ-PyHpHpImPyIm
25	2430) 5'W C G C A A A W-3'	PyImPyPyPyPy-γ-HpHpHpImPyIm
	2431) 5'W C G C A A G W-3'	PyImPyPyPyIm-γ-PyHpHpImPyIm
	2432) 5'W C G C A A C W-3'	PyImPyPyPyPy-γ-ImHpHpImPyIm
	2433) 5'W C G C A G T W-3'	PyImPyPyImHp-γ-PyPyHpImPyIm
	2434) 5'W C G C A G A W-3'	PyImPyPyImPy-γ-HpPyHpImPyIm
30	2435) 5'W C G C A G G W-3'	PyImPyPyImIm-γ-PyPyHpImPyIm
	2436) 5'W C G C A G C W-3'	PyImPyPyImPy-γ-ImPyHpImPyIm
	2437) 5'W C G C A C T W-3'	PyImPyPyPyHp-γ-PyImHpImPyIm
	2438) 5'W C G C A C A W-3'	PyImPyPyPyPy-γ-HpImHpImPyIm
	2439) 5'W C G C A C G W-3'	PyImPyPyPyIm-γ-PyImHpImPyIm
35	2440) 5'W C G C A C C W-3'	PyImPyPyPyPy-γ-ImImHpImPyIm

TABLE 123: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCGCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2441) 5'W C G C G T T W-3'	PyImPyImHpHp-γ-PyPyPyImPyIm
5	2442) 5'W C G C G T A W-3'	PyImPyImHpPy-γ-HpPyPyImPyIm
	2443) 5'W C G C G T G W-3'	PyImPyImHpIm-γ-PyPyPyImPyIm
	2444) 5'W C G C G T C W-3'	PyImPyImHpPy-γ-ImPyPyImPyIm
	2445) 5'W C G C G A T W-3'	PyImPyImPyHp-γ-PyHpPyImPyIm
	2446) 5'W C G C G A A W-3'	PyImPyImPyPy-γ-HpHpPyImPyIm
10	2447) 5'W C G C G A G W-3'	PyImPyImPyIm-γ-PyHpPyImPyIm
	2448) 5'W C G C G A C W-3'	PyImPyImPyPy-γ-ImHpPyImPyIm
	2449) 5'W C G C G G T W-3'	PyImPyImImHp-γ-PyPyPyImPyIm
	2450) 5'W C G C G G A W-3'	PyImPyImImPy-γ-HpPyPyImPyIm
	2451) 5'W C G C G C T W-3'	PyImPyImPyHp-γ-PyImPyImPyIm
15	2452) 5'W C G C G C A W-3'	PyImPyImPyPy-γ-HpImPyImPyIm
	2453) 5'W C G C C T T W-3'	PyImPyPyHpHp-γ-PyPyImImPyIm
	2454) 5'W C G C C T A W-3'	PyImPyPyHpPy-γ-HpPyImImPyIm
	2455) 5'W C G C C T G W-3'	PyImPyPyHpIm-γ-PyPyImImPyIm
	2456) 5'W C G C C T C W-3'	PyImPyPyHpPy-γ-ImPyImImPyIm
20	2457) 5'W C G C C A T W-3'	PyImPyPyPyHp-γ-PyHpImImPyIm
	2458) 5'W C G C C A A W-3'	PyImPyPyPyPy-γ-HpHpImImPyIm
	2459) 5'W C G C C A G W-3'	PyImPyPyPyIm-γ-PyHpImImPyIm
	2460) 5'W C G C C A C W-3'	PyImPyPyPyPy-γ-ImHpImImPyIm
	2461) 5'W C G C C G T W-3'	PyImPyPyImHp-γ-PyPyImImPyIm
25	2462) 5'W C G C C G A W-3'	PyImPyPyImPy-γ-HpPyImImPyIm
	2463) 5'W C G C C C T W-3'	PyImPyPyPyHp-γ-PyImImImPyIm
	2464) 5'W C G C C C A W-3'	PyImPyPyPyPy-γ-HpImImImPyIm
	G91) 5'W C G C G G G W-3'	PyImPyImImIm-γ-PyPyPyImPyIm
	G92) 5'W C G C G G C W-3'	PyImPyImImPy-γ-ImPyPyImPyIm
30	G93) 5'W C G C G C G W-3'	PyImPyImPyIm-γ-PyImPyImPyIm
	G94) 5'W C G C G C C W-3'	PyImPyImPyPy-γ-ImImPyImPyIm
	G95) 5'W C G C C G G W-3'	PyImPyPyImIm-γ-PyPyImImPyIm
	G96) 5'W C G C C G C W-3'	PyImPyPyImPy-γ-ImPyImImPyIm
	G97) 5'W C G C C C G W-3'	PyImPyPyPyIm-γ-PyImImImPyIm
35	G98) 5'W C G C C C C W-3'	PyImPyPyPyPy-γ-ImImImImPyIm

TABLE 124: 12-ring Harpin Polyamides for recognition of 8-bp 5'-WCCGWNW-3'

	DNA sequence	aromatic amino acid sequence
	2465) 5'W C C G T T T W-3'	PyPyImHpHpHp-γ-PyPyPyPyImIm
5	2466) 5'W C C G T T A W-3'	PyPyImHpHpPy-γ-HpPyPyPyImIm
	2467) 5'W C C G T T G W-3'	PyPyImHpHpIm-γ-PyPyPyPyImIm
	2468) 5'W C C G T T C W-3'	PyPyImHpHpPy-γ-ImPyPyPyImIm
	2469) 5'W C C G T A T W-3'	PyPyImHpPyHp-γ-PyHpPyPyImIm
	2470) 5'W C C G T A A W-3'	PyPyImHpPyPy-γ-HpHpPyPyImIm
10	2471) 5'W C C G T A G W-3'	PyPyImHpPyIm-γ-PyHpPyPyImIm
	2472) 5'W C C G T A C W-3'	PyPyImHpPyPy-γ-ImHpPyPyImIm
	2473) 5'W C C G T G T W-3'	PyPyImHpImHp-γ-PyPyPyPyImIm
	2474) 5'W C C G T G A W-3'	PyPyImHpImPy-γ-HpPyPyPyImIm
	2475) 5'W C C G T G G W-3'	PyPyImHpImIm-γ-PyPyPyPyImIm
15	2476) 5'W C C G T G C W-3'	PyPyImHpImPy-γ-ImPyPyPyImIm
	2477) 5'W C C G T C T W-3'	PyPyImHpPyHp-γ-PyImPyPyImIm
	2478) 5'W C C G T C A W-3'	PyPyImHpPyPy-γ-HpImPyPyImIm
	2479) 5'W C C G T C G W-3'	PyPyImHpPyIm-γ-PyImPyPyImIm
	2480) 5'W C C G T C C W-3'	PyPyImHpPyPy-γ-ImImPyPyImIm
20	2481) 5'W C C G A T T W-3'	PyPyImPyHpHp-γ-PyPyHpPyImIm
	2482) 5'W C C G A T A W-3'	PyPyImPyHpPy-γ-HpPyHpPyImIm
	2483) 5'W C C G A T G W-3'	PyPyImPyHpIm-γ-PyPyHpPyImIm
	2484) 5'W C C G A T C W-3'	PyPyImPyHpPy-γ-ImPyHpPyImIm
	2485) 5'W C C G A A T W-3'	PyPyImPyPyHp-γ-PyHpHpPyImIm
25	2486) 5'W C C G A A A W-3'	PyPyImPyPyPy-γ-HpHpHpPyImIm
	2487) 5'W C C G A A G W-3'	PyPyImPyPyIm-γ-PyHpHpPyImIm
	2488) 5'W C C G A A C W-3'	PyPyImPyPyPy-γ-ImHpHpPyImIm
	2489) 5'W C C G A G T W-3'	PyPyImPyImHp-γ-PyPyHpPyImIm
	2490) 5'W C C G A G A W-3'	PyPyImPyImPy-γ-HpPyHpPyImIm
30	2491) 5'W C C G A G G W-3'	PyPyImPyImIm-γ-PyPyHpPyImIm
	2492) 5'W C C G A G C W-3'	PyPyImPyImPy-γ-ImPyHpPyImIm
	2493) 5'W C C G A C T W-3'	PyPyImPyPyHp-γ-PyImHpPyImIm
	2494) 5'W C C G A C A W-3'	PyPyImPyPyPy-γ-HpImHpPyImIm
	2495) 5'W C C G A C G W-3'	PyPyImPyPyIm-γ-PyImHpPyImIm
35	2496) 5'W C C G A C C W-3'	PyPyImPyPyPy-γ-ImImHpPyImIm

TABLE 125: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2497) 5'W C C G G T T W-3'	PyPyImImHpHp-γ-PyPyPyPyImIm
5	2498) 5'W C C G G T A W-3'	PyPyImImHpPy-γ-HpPyPyPyImIm
	2499) 5'W C C G G T G W-3'	PyPyImImHpIm-γ-PyPyPyPyImIm
	2500) 5'W C C G G T C W-3'	PyPyImImHpPy-γ-ImPyPyPyImIm
	2501) 5'W C C G G A T W-3'	PyPyImImPyHp-γ-PyHpPyPyImIm
	2502) 5'W C C G G A A W-3'	PyPyImImPyPy-γ-HpHpPyPyImIm
10	2503) 5'W C C G G A G W-3'	PyPyImImPyIm-γ-PyHpPyPyImIm
	2504) 5'W C C G G A C W-3'	PyPyImImPyPy-γ-ImHpPyPyImIm
	2505) 5'W C C G G G T W-3'	PyPyImImImHp-γ-PyPyPyPyImIm
	2506) 5'W C C G G G A W-3'	PyPyImImImPy-γ-HpPyPyPyImIm
	2507) 5'W C C G G C T W-3'	PyPyImImPyHp-γ-PyImPyPyImIm
	2508) 5'W C C G G C A W-3'	PyPyImImPyPy-γ-HpImPyPyImIm
	2509) 5'W C C G C T T W-3'	PyPyImPyHpHp-γ-PyPyImPyImIm
	2510) 5'W C C G C T A W-3'	PyPyImPyHpPy-γ-HpPyImPyImIm
	2511) 5'W C C G C T G W-3'	PyPyImPyHpIm-γ-PyPyImPyImIm
	2512) 5'W C C G C T C W-3'	PyPyImPyHpPy-γ-ImPyImPyImIm
20	2513) 5'W C C G C A T W-3'	PyPyImPyPyHp-γ-PyHpImPyImIm
	2514) 5'W C C G C A A W-3'	PyPyImPyPyPy-γ-HpHpImPyImIm
	2515) 5'W C C G C A G W-3'	PyPyImPyPyIm-γ-PyHpImPyImIm
	2516) 5'W C C G C A C W-3'	PyPyImPyPyPy-γ-ImHpImPyImIm
	2517) 5'W C C G C G T W-3'	PyPyImPyImHp-γ-PyPyImPyImIm
25	2518) 5'W C C G C G A W-3'	PyPyImPyImPy-γ-HpPyImPyImIm
	2519) 5'W C C G C C T W-3'	PyPyImPyPyHp-γ-PyImImPyImIm
	2520) 5'W C C G C C A W-3'	PyPyImPyPyPy-γ-HpImImPyImIm
	G99) 5'W C C G G G G W-3'	PyPyImImImIm-γ-PyPyPyPyImIm
	G100) 5'W C C G G G C W-3'	PyPyImImImPy-γ-ImPyPyPyImIm
30	G101) 5'W C C G G C G W-3'	PyPyImImPyIm-γ-PyImPyPyImIm
	G102) 5'W C C G G C C W-3'	PyPyImImPyPy-γ-ImImPyPyImIm
	G103) 5'W C C G C G G W-3'	PyPyImPyImIm-γ-PyPyImPyImIm
	G104) 5'W C C G C G C W-3'	PyPyImPyImPy-γ-ImPyImPyImIm
	G105) 5'W C C G C C G W-3'	PyPyImPyPyIm-γ-PyImImPyImIm
35	G106) 5'W C C G C C C W-3'	PyPyImPyPyPy-γ-ImImImPyImIm

TABLE 126: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCTWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	2521) 5'W C C T T T T W-3'	PyPyHpHpHpHp-γ-PyPyPyPyImIm
	2522) 5'W C C T T T A W-3'	PyPyHpHpHpPy-γ-HpPyPyPyImIm
	2523) 5'W C C T T T G W-3'	PyPyHpHpHpIm-γ-PyPyPyPyImIm
	2524) 5'W C C T T T C W-3'	PyPyHpHpHpPy-γ-ImPyPyPyImIm
10	2525) 5'W C C T T A T W-3'	PyPyHpHpPyHp-γ-PyHpPyPyImIm
	2526) 5'W C C T T A A W-3'	PyPyHpHpPyPy-γ-HpHpPyPyImIm
	2527) 5'W C C T T A G W-3'	PyPyHpHpPyIm-γ-PyHpPyPyImIm
	2528) 5'W C C T T A C W-3'	PyPyHpHpPyPy-γ-ImHpPyPyImIm
	2529) 5'W C C T T G T W-3'	PyPyHpHpImHp-γ-PyPyPyPyImIm
	2530) 5'W C C T T G A W-3'	PyPyHpHpImPy-γ-HpPyPyPyImIm
	2531) 5'W C C T T G G W-3'	PyPyHpHpImIm-γ-PyPyPyPyImIm
	2532) 5'W C C T T G C W-3'	PyPyHpHpImPy-γ-ImPyPyPyImIm
	2533) 5'W C C T T C T W-3'	PyPyHpHpPyHp-γ-PyImPyPyImIm
	2534) 5'W C C T T C A W-3'	PyPyHpHpPyPy-γ-HpImPyPyImIm
20	2535) 5'W C C T T C G W-3'	PyPyHpHpPyIm-γ-PyImPyPyImIm
	2536) 5'W C C T T C C W-3'	PyPyHpHpPyPy-γ-ImImPyPyImIm
	2537) 5'W C C T A T T W-3'	PyPyHpPyHpHp-γ-PyPyHpPyImIm
	2538) 5'W C C T A T A W-3'	PyPyHpPyHpPy-γ-HpPyHpPyImIm
	2539) 5'W C C T A T G W-3'	PyPyHpPyHpIm-γ-PyPyHpPyImIm
	2540) 5'W C C T A T C W-3'	PyPyHpPyHpPy-γ-ImPyHpPyImIm
	2541) 5'W C C T A A T W-3'	PyPyHpPyPyHp-γ-PyHpHpPyImIm
	2542) 5'W C C T A A A W-3'	PyPyHpPyPyPy-γ-HpHpHpPyImIm
	2543) 5'W C C T A A G W-3'	PyPyHpPyPyIm-γ-PyHpHpPyImIm
	2544) 5'W C C T A A C W-3'	PyPyHpPyPyPy-γ-ImHpHpPyImIm
30	2545) 5'W C C T A G T W-3'	PyPyHpPyImHp-γ-PyPyHpPyImIm
	2546) 5'W C C T A G A W-3'	PyPyHpPyImPy-γ-HpPyHpPyImIm
	2547) 5'W C C T A G G W-3'	PyPyHpPyImIm-γ-PyPyHpPyImIm
	2548) 5'W C C T A G C W-3'	PyPyHpPyImPy-γ-ImPyHpPyImIm
	2549) 5'W C C T A C T W-3'	PyPyHpPyPyHp-γ-PyImHpPyImIm
	2550) 5'W C C T A C A W-3'	PyPyHpPyPyPy-γ-HpImHpPyImIm
	2551) 5'W C C T A C G W-3'	PyPyHpPyPyIm-γ-PyImHpPyImIm
	2552) 5'W C C T A C C W-3'	PyPyHpPyPyPy-γ-ImImHpPyImIm

TABLE 127: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCTSNW-3'

DNA sequence		aromatic amino acid sequence
2553)	5'W C C T G T T W-3'	PyPyHpImHpHp-γ-PyPyPyPyImIm
2554)	5'W C C T G T A W-3'	PyPyHpImHpPy-γ-HpPyPyPyImIm
2555)	5'W C C T G T G W-3'	PyPyHpImHpIm-γ-PyPyPyPyImIm
2556)	5'W C C T G T C W-3'	PyPyHpImHpPy-γ-ImPyPyPyImIm
2557)	5'W C C T G A T W-3'	PyPyHpImPyHp-γ-PyHpPyPyImIm
2558)	5'W C C T G A A W-3'	PyPyHpImPyPy-γ-HpHpPyPyImIm
2559)	5'W C C T G A G W-3'	PyPyHpImPyIm-γ-PyHpPyPyImIm
2560)	5'W C C T G A C W-3'	PyPyHpImPyPy-γ-ImHpPyPyImIm
2561)	5'W C C T G G T W-3'	PyPyHpImImHp-γ-PyPyPyPyImIm
2562)	5'W C C T G G A W-3'	PyPyHpImImPy-γ-HpPyPyPyImIm
2563)	5'W C C T G C T W-3'	PyPyHpImPyHp-γ-PyImPyPyImIm
2564)	5'W C C T G C A W-3'	PyPyHpImPyPy-γ-HpImPyPyImIm
2565)	5'W C C T G G G W-3'	PyPyHpImImIm-γ-PyPyPyPyImIm
2566)	5'W C C T G G C W-3'	PyPyHpImImPy-γ-ImPyPyPyImIm
2567)	5'W C C T G C G W-3'	PyPyHpImPyIm-γ-PyImPyPyImIm
2568)	5'W C C T G C C W-3'	PyPyHpImPyPy-γ-ImImPyPyImIm
2569)	5'W C C T C T T W-3'	PyPyHpPyHpHp-γ-PyPyImPyImIm
2570)	5'W C C T C T A W-3'	PyPyHpPyHpPy-γ-HpPyImPyImIm
2571)	5'W C C T C T G W-3'	PyPyHpPyHpIm-γ-PyPyImPyImIm
2572)	5'W C C T C T C W-3'	PyPyHpPyHpPy-γ-ImPyImPyImIm
2573)	5'W C C T C A T W-3'	PyPyHpPyPyHp-γ-PyHpImPyImIm
2574)	5'W C C T C A A W-3'	PyPyHpPyPyPy-γ-HpHpImPyImIm
2575)	5'W C C T C A G W-3'	PyPyHpPyPyIm-γ-PyHpImPyImIm
2576)	5'W C C T C A C W-3'	PyPyHpPyPyPy-γ-ImHpImPyImIm
2577)	5'W C C T C G T W-3'	PyPyHpPyImHp-γ-PyPyImPyImIm
2578)	5'W C C T C G A W-3'	PyPyHpPyImPy-γ-HpPyImPyImIm
2579)	5'W C C T C C T W-3'	PyPyHpPyPyHp-γ-PyImImPyImIm
2580)	5'W C C T C C A W-3'	PyPyHpPyPyPy-γ-HpImImPyImIm
2581)	5'W C C T C G G W-3'	PyPyHpPyImIm-γ-PyPyImPyImIm
2582)	5'W C C T C G C W-3'	PyPyHpPyImPy-γ-ImPyImPyImIm
2583)	5'W C C T C C G W-3'	PyPyHpPyPyIm-γ-PyImImPyImIm
2584)	5'W C C T C C C W-3'	PyPyHpPyPyPy-γ-ImImImPyImIm

TABLE 128: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCA WNNW-3'

	DNA sequence	aromatic amino acid sequence
	2585) 5'W C C A T T T W-3'	PyPyPyHpHpHp-γ-PyPyPyHpImIm
5	2586) 5'W C C A T T A W-3'	PyPyPyHpHpPy-γ-HpPyPyHpImIm
	2587) 5'W C C A T T G W-3'	PyPyPyHpHpIm-γ-PyPyPyHpImIm
	2588) 5'W C C A T T C W-3'	PyPyPyHpHpPy-γ-ImPyPyHpImIm
	2589) 5'W C C A T A T W-3'	PyPyPyHpPyHp-γ-PyHpPyHpImIm
	2590) 5'W C C A T A A W-3'	PyPyPyHpPyPy-γ-HpHpPyHpImIm
10	2591) 5'W C C A T A G W-3'	PyPyPyHpPyIm-γ-PyHpPyHpImIm
	2592) 5'W C C A T A C W-3'	PyPyPyHpPyPy-γ-ImHpPyHpImIm
	2593) 5'W C C A T G T W-3'	PyPyPyHpImHp-γ-PyPyPyHpImIm
	2594) 5'W C C A T G A W-3'	PyPyPyHpImPy-γ-HpPyPyHpImIm
	2595) 5'W C C A T G G W-3'	PyPyPyHpImIm-γ-PyPyPyHpImIm
	2596) 5'W C C A T G C W-3'	PyPyPyHpImPy-γ-ImPyPyHpImIm
	2597) 5'W C C A T C T W-3'	PyPyPyHpPyHp-γ-PyImPyHpImIm
	2598) 5'W C C A T C A W-3'	PyPyPyHpPyPy-γ-HpImPyHpImIm
	2599) 5'W C C A T C G W-3'	PyPyPyHpPyIm-γ-PyImPyHpImIm
	2600) 5'W C C A T C C W-3'	PyPyPyHpPyPy-γ-ImImPyHpImIm
20	2601) 5'W C C A A T T W-3'	PyPyPyPyHpHp-γ-PyPyHpHpImIm
	2602) 5'W C C A A T A W-3'	PyPyPyPyHpPy-γ-HpPyHpHpImIm
	2603) 5'W C C A A T G W-3'	PyPyPyPyHpIm-γ-PyPyHpHpImIm
	2604) 5'W C C A A T C W-3'	PyPyPyPyHpPy-γ-ImPyHpHpImIm
	2605) 5'W C C A A A T W-3'	PyPyPyPyPyHp-γ-PyHpHpHpImIm
25	2606) 5'W C C A A A A W-3'	PyPyPyPyPyPy-γ-HpHpHpHpImIm
	2607) 5'W C C A A A G W-3'	PyPyPyPyPyIm-γ-PyHpHpHpImIm
	2608) 5'W C C A A A C W-3'	PyPyPyPyPyPy-γ-ImHpHpHpImIm
	2609) 5'W C C A A G T W-3'	PyPyPyPyImHp-γ-PyPyHpHpImIm
	2610) 5'W C C A A G A W-3'	PyPyPyPyImPy-γ-HpPyHpHpImIm
30	2611) 5'W C C A A G G W-3'	PyPyPyPyImIm-γ-PyPyHpHpImIm
	2612) 5'W C C A A G C W-3'	PyPyPyPyImPy-γ-ImPyHpHpImIm
	2613) 5'W C C A A C T W-3'	PyPyPyPyPyHp-γ-PyImHpHpImIm
	2614) 5'W C C A A C A W-3'	PyPyPyPyPyPy-γ-HpImHpHpImIm
	2615) 5'W C C A A C G W-3'	PyPyPyPyPyIm-γ-PyImHpHpImIm
35	2616) 5'W C C A A C C W-3'	PyPyPyPyPyPy-γ-ImImHpHpImIm

TABLE 129: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCASNNW-3'

	DNA sequence	aromatic amino acid sequence
	2617) 5'W C C A G T T W-3'	PyPyPyImHpHp- γ -PyPyPyHpImIm
5	2618) 5'W C C A G T A W-3'	PyPyPyImHpPy- γ -HpPyPyHpImIm
	2619) 5'W C C A G T G W-3'	PyPyPyImHpIm- γ -PyPyPyHpImIm
	2620) 5'W C C A G T C W-3'	PyPyPyImHpPy- γ -ImPyPyHpImIm
	2621) 5'W C C A G A T W-3'	PyPyPyImPyHp- γ -PyHpPyHpImIm
	2622) 5'W C C A G A A W-3'	PyPyPyImPyPy- γ -HpHpPyHpImIm
10	2623) 5'W C C A G A G W-3'	PyPyPyImPyIm- γ -PyHpPyHpImIm
	2624) 5'W C C A G A C W-3'	PyPyPyImPyPy- γ -ImHpPyHpImIm
	2625) 5'W C C A G G T W-3'	PyPyPyImImHp- γ -PyPyPyHpImIm
	2626) 5'W C C A G G A W-3'	PyPyPyImImPy- γ -HpPyPyHpImIm
	2627) 5'W C C A G C T W-3'	PyPyPyImPyHp- γ -PyImPyHpImIm
	2628) 5'W C C A G C A W-3'	PyPyPyImPyPy- γ -HpImPyHpImIm
	2629) 5'W C C A G G G W-3'	PyPyPyImImIm- γ -PyPyPyHpImIm
	2630) 5'W C C A G G C W-3'	PyPyPyImImPy- γ -ImPyPyHpImIm
	2631) 5'W C C A G C G W-3'	PyPyPyImPyIm- γ -PyImPyHpImIm
	2632) 5'W C C A G C C W-3'	PyPyPyImPyPy- γ -ImImPyHpImIm
20	2633) 5'W C C A C T T W-3'	PyPyPyPyHpHp- γ -PyPyImHpImIm
	2634) 5'W C C A C T A W-3'	PyPyPyPyHpPy- γ -HpPyImHpImIm
	2635) 5'W C C A C T G W-3'	PyPyPyPyHpIm- γ -PyPyImHpImIm
	2636) 5'W C C A C T C W-3'	PyPyPyPyHpPy- γ -ImPyImHpImIm
	2637) 5'W C C A C A T W-3'	PyPyPyPyPyHp- γ -PyHpImHpImIm
25	2638) 5'W C C A C A A W-3'	PyPyPyPyPyPy- γ -HpHpImHpImIm
	2639) 5'W C C A C A G W-3'	PyPyPyPyPyIm- γ -PyHpImHpImIm
	2640) 5'W C C A C A C W-3'	PyPyPyPyPyPy- γ -ImHpImHpImIm
	2641) 5'W C C A C G T W-3'	PyPyPyPyImHp- γ -PyPyImHpImIm
	2642) 5'W C C A C G A W-3'	PyPyPyPyImPy- γ -HpPyImHpImIm
30	2643) 5'W C C A C C T W-3'	PyPyPyPyPyHp- γ -PyImImHpImIm
	2644) 5'W C C A C C A W-3'	PyPyPyPyPyPy- γ -HpImImHpImIm
	2645) 5'W C C A C G G W-3'	PyPyPyPyImIm- γ -PyPyImHpImIm
	2646) 5'W C C A C G C W-3'	PyPyPyPyImPy- γ -ImPyImHpImIm
	2647) 5'W C C A C C G W-3'	PyPyPyPyPyIm- γ -PyImImHpImIm
35	2648) 5'W C C A C C C W-3'	PyPyPyPyPyPy- γ -ImImImHpImIm

TABLE 130: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2649) 5'W C C C T T T W-3'	PyPyPyHpHpHp-γ-PyPyPyImImIm
5	2650) 5'W C C C T T A W-3'	PyPyPyHpHpPy-γ-HpPyPyImImIm
	2651) 5'W C C C T T G W-3'	PyPyPyHpHpIm-γ-PyPyPyImImIm
	2652) 5'W C C C T T C W-3'	PyPyPyHpHpPy-γ-ImPyPyImImIm
	2653) 5'W C C C T A T W-3'	PyPyPyHpPyHp-γ-PyHpPyImImIm
	2654) 5'W C C C T A A W-3'	PyPyPyHpPyPy-γ-HpHpPyImImIm
10	2655) 5'W C C C T A G W-3'	PyPyPyHpPyIm-γ-PyHpPyImImIm
	2656) 5'W C C C T A C W-3'	PyPyPyHpPyPy-γ-ImHpPyImImIm
	2657) 5'W C C C T G T W-3'	PyPyPyHpImHp-γ-PyPyPyImImIm
	2658) 5'W C C C T G A W-3'	PyPyPyHpImPy-γ-HpPyPyImImIm
	2659) 5'W C C C T G G W-3'	PyPyPyHpImIm-γ-PyPyPyImImIm
	2660) 5'W C C C T G C W-3'	PyPyPyHpImPy-γ-ImPyPyImImIm
	2661) 5'W C C C T C T W-3'	PyPyPyHpPyHp-γ-PyImPyImImIm
	2662) 5'W C C C T C A W-3'	PyPyPyHpPyPy-γ-HpImPyImImIm
	2663) 5'W C C C T C G W-3'	PyPyPyHpPyIm-γ-PyImPyImImIm
	2664) 5'W C C C T C C W-3'	PyPyPyHpPyPy-γ-ImImPyImImIm
20	2665) 5'W C C C A T T W-3'	PyPyPyPyHpHp-γ-PyPyHpImImIm
	2666) 5'W C C C A T A W-3'	PyPyPyPyHpPy-γ-HpPyHpImImIm
	2667) 5'W C C C A T G W-3'	PyPyPyPyHpIm-γ-PyPyHpImImIm
	2668) 5'W C C C A T C W-3'	PyPyPyPyHpPy-γ-ImPyHpImImIm
	2669) 5'W C C C A A T W-3'	PyPyPyPyPyHp-γ-PyHpHpImImIm
25	2670) 5'W C C C A A A W-3'	PyPyPyPyPyPy-γ-HpHpHpImImIm
	2671) 5'W C C C A A G W-3'	PyPyPyPyPyIm-γ-PyHpHpImImIm
	2672) 5'W C C C A A C W-3'	PyPyPyPyPyPy-γ-ImHpHpImImIm
	2673) 5'W C C C A G T W-3'	PyPyPyPyImHp-γ-PyPyHpImImIm
	2674) 5'W C C C A G A W-3'	PyPyPyPyImPy-γ-HpPyHpImImIm
30	2675) 5'W C C C A G G W-3'	PyPyPyPyImIm-γ-PyPyHpImImIm
	2676) 5'W C C C A G C W-3'	PyPyPyPyImPy-γ-ImPyHpImImIm
	2677) 5'W C C C A C T W-3'	PyPyPyPyPyHp-γ-PyImHpImImIm
	2678) 5'W C C C A C A W-3'	PyPyPyPyPyPy-γ-HpImHpImImIm
	2679) 5'W C C C A C G W-3'	PyPyPyPyPyIm-γ-PyImHpImImIm
35	2680) 5'W C C C A C C W-3'	PyPyPyPyPyPy-γ-ImImHpImImIm

TABLE 131: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCCCSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	2681) 5'W C C C G T T W-3'	PyPyPyImHpHp-γ-PyPyPyImImIm
	2682) 5'W C C C G T A W-3'	PyPyPyImHpPy-γ-HpPyPyImImIm
	2683) 5'W C C C G T G W-3'	PyPyPyImHpIm-γ-PyPyPyImImIm
	2684) 5'W C C C G T C W-3'	PyPyPyImHpPy-γ-ImPyPyImImIm
	2685) 5'W C C C G A T W-3'	PyPyPyImPyHp-γ-PyHpPyImImIm
10	2686) 5'W C C C G A A W-3'	PyPyPyImPyPy-γ-HpHpPyImImIm
	2687) 5'W C C C G A G W-3'	PyPyPyImPyIm-γ-PyHpPyImImIm
	2688) 5'W C C C G A C W-3'	PyPyPyImPyPy-γ-ImHpPyImImIm
	2689) 5'W C C C G G T W-3'	PyPyPyImImHp-γ-PyPyPyImImIm
	2690) 5'W C C C G G A W-3'	PyPyPyImImPy-γ-HpPyPyImImIm
	2691) 5'W C C C G C T W-3'	PyPyPyImPyHp-γ-PyImPyImImIm
	2692) 5'W C C C G C A W-3'	PyPyPyImPyPy-γ-HpImPyImImIm
	2693) 5'W C C C C T T W-3'	PyPyPyPyHpHp-γ-PyPyImImImIm
	2694) 5'W C C C C T A W-3'	PyPyPyPyHpPy-γ-HpPyImImImIm
	2695) 5'W C C C C T G W-3'	PyPyPyPyHpIm-γ-PyPyImImImIm
20	2696) 5'W C C C C T C W-3'	PyPyPyPyHpPy-γ-ImPyImImImIm
	2697) 5'W C C C C A T W-3'	PyPyPyPyPyHp-γ-PyHpImImImIm
	2698) 5'W C C C C A A W-3'	PyPyPyPyPyPy-γ-HpHpImImImIm
	2699) 5'W C C C C A G W-3'	PyPyPyPyPyIm-γ-PyHpImImImIm
	2690) 5'W C C C C A C W-3'	PyPyPyPyPyPy-γ-ImHpImImImIm
	2701) 5'W C C C C G T W-3'	PyPyPyPyImHp-γ-PyPyImImImIm
	2702) 5'W C C C C G A W-3'	PyPyPyPyImPy-γ-HpPyImImImIm
	2703) 5'W C C C C C T W-3'	PyPyPyPyPyHp-γ-PyImImImImIm
	2704) 5'W C C C C C A W-3'	PyPyPyPyPyPy-γ-HpImImImImIm
	G107) 5'W C C C G G G W-3'	PyPyPyImImIm-γ-PyPyPyImImIm
30	G108) 5'W C C C G G C W-3'	PyPyPyImImPy-γ-ImPyPyImImIm
	G109) 5'W C C C G C G W-3'	PyPyPyImPyIm-γ-PyImPyImImIm
	G110) 5'W C C C G C C W-3'	PyPyPyImPyPy-γ-ImImPyImImIm
	G111) 5'W C C C C G G W-3'	PyPyPyPyImIm-γ-PyPyImImImIm
	G112) 5'W C C C C G C W-3'	PyPyPyPyImPy-γ-ImPyImImImIm
35	G113) 5'W C C C C C G W-3'	PyPyPyPyPyIm-γ-PyImImImImIm
	G114) 5'W C C C C C C W-3'	PyPyPyPyPyPy-γ-ImImImImImIm

TABLE 132: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCAGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2705) 5'W C A G T T T W-3'	PyPyImHpHpHp-γ-PyPyPyPyHpIm
5	2706) 5'W C A G T T A W-3'	PyPyImHpHpPy-γ-HpPyPyPyHpIm
	2707) 5'W C A G T T G W-3'	PyPyImHpHpIm-γ-PyPyPyPyHpIm
	2708) 5'W C A G T T C W-3'	PyPyImHpHpPy-γ-ImPyPyPyHpIm
	2709) 5'W C A G T A T W-3'	PyPyImHpPyHp-γ-PyHpPyPyHpIm
	2700) 5'W C A G T A A W-3'	PyPyImHpPyPy-γ-HpHpPyPyHpIm
10	2711) 5'W C A G T A G W-3'	PyPyImHpPyIm-γ-PyHpPyPyHpIm
	2712) 5'W C A G T A C W-3'	PyPyImHpPyPy-γ-ImHpPyPyHpIm
	2713) 5'W C A G T G T W-3'	PyPyImHpImHp-γ-PyPyPyPyHpIm
	2714) 5'W C A G T G A W-3'	PyPyImHpImPy-γ-HpPyPyPyHpIm
	2715) 5'W C A G T G G W-3'	PyPyImHpImIm-γ-PyPyPyPyHpIm
	2716) 5'W C A G T G C W-3'	PyPyImHpImPy-γ-ImPyPyPyHpIm
	2717) 5'W C A G T C T W-3'	PyPyImHpPyHp-γ-PyImPyPyHpIm
	2718) 5'W C A G T C A W-3'	PyPyImHpPyPy-γ-HpImPyPyHpIm
	2719) 5'W C A G T C G W-3'	PyPyImHpPyIm-γ-PyImPyPyHpIm
	2720) 5'W C A G T C C W-3'	PyPyImHpPyPy-γ-ImImPyPyHpIm
	2721) 5'W C A G A T T W-3'	PyPyImPyHpHp-γ-PyPyHpPyHpIm
	2722) 5'W C A G A T A W-3'	PyPyImPyHpPy-γ-HpPyHpPyHpIm
	2723) 5'W C A G A T G W-3'	PyPyImPyHpIm-γ-PyPyHpPyHpIm
	2724) 5'W C A G A T C W-3'	PyPyImPyHpPy-γ-ImPyHpPyHpIm
	2725) 5'W C A G A A T W-3'	PyPyImPyPyHp-γ-PyHpHpPyHpIm
25	2726) 5'W C A G A A A W-3'	PyPyImPyPyPy-γ-HpHpHpPyHpIm
	2727) 5'W C A G A A G W-3'	PyPyImPyPyIm-γ-PyHpHpPyHpIm
	2728) 5'W C A G A A C W-3'	PyPyImPyPyPy-γ-ImHpHpPyHpIm
	2729) 5'W C A G A G T W-3'	PyPyImPyImHp-γ-PyPyHpPyHpIm
	2730) 5'W C A G A G A W-3'	PyPyImPyImPy-γ-HpPyHpPyHpIm
30	2731) 5'W C A G A G G W-3'	PyPyImPyImIm-γ-PyPyHpPyHpIm
	2732) 5'W C A G A G C W-3'	PyPyImPyImPy-γ-ImPyHpPyHpIm
	2733) 5'W C A G A C T W-3'	PyPyImPyPyHp-γ-PyImHpPyHpIm
	2734) 5'W C A G A C A W-3'	PyPyImPyPyPy-γ-HpImHpPyHpIm
	2735) 5'W C A G A C G W-3'	PyPyImPyPyIm-γ-PyImHpPyHpIm
35	2736) 5'W C A G A C C W-3'	PyPyImPyPyPy-γ-ImImHpPyHpIm

TABLE 133: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCAGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2737) 5'W C A G G T T W-3'	PyPyImImHpHp-γ-PyPyPyPyHpIm
5	2738) 5'W C A G G T A W-3'	PyPyImImHpPy-γ-HpPyPyPyHpIm
	2739) 5'W C A G G T G W-3'	PyPyImImHpIm-γ-PyPyPyPyHpIm
	2740) 5'W C A G G T C W-3'	PyPyImImHpPy-γ-ImPyPyPyHpIm
	2741) 5'W C A G G A T W-3'	PyPyImImPyHp-γ-PyHpPyPyHpIm
	2742) 5'W C A G G A A W-3'	PyPyImImPyPy-γ-HpHpPyPyHpIm
10	2743) 5'W C A G G A G W-3'	PyPyImImPyIm-γ-PyHpPyPyHpIm
	2744) 5'W C A G G A C W-3'	PyPyImImPyPy-γ-ImHpPyPyHpIm
	2745) 5'W C A G G G T W-3'	PyPyImImImHp-γ-PyPyPyPyHpIm
	2746) 5'W C A G G G A W-3'	PyPyImImImPy-γ-HpPyPyPyHpIm
	2747) 5'W C A G G C T W-3'	PyPyImImPyHp-γ-PyImPyPyHpIm
	2748) 5'W C A G G C A W-3'	PyPyImImPyPy-γ-HpImPyPyHpIm
	2749) 5'W C A G C T T W-3'	PyPyImPyHpHp-γ-PyPyImPyHpIm
	2750) 5'W C A G C T A W-3'	PyPyImPyHpPy-γ-HpPyImPyHpIm
	2751) 5'W C A G C T G W-3'	PyPyImPyHpIm-γ-PyPyImPyHpIm
	2752) 5'W C A G C T C W-3'	PyPyImPyHpPy-γ-ImPyImPyHpIm
	2753) 5'W C A G C A T W-3'	PyPyImPyPyHp-γ-PyHpImPyHpIm
	2754) 5'W C A G C A A W-3'	PyPyImPyPyPy-γ-HpHpImPyHpIm
	2755) 5'W C A G C A G W-3'	PyPyImPyPyIm-γ-PyHpImPyHpIm
	2756) 5'W C A G C A C W-3'	PyPyImPyPyPy-γ-ImHpImPyHpIm
	2757) 5'W C A G C G T W-3'	PyPyImPyImHp-γ-PyPyImPyHpIm
25	2758) 5'W C A G C G A W-3'	PyPyImPyImPy-γ-HpPyImPyHpIm
	2759) 5'W C A G C C T W-3'	PyPyImPyPyHp-γ-PyImImPyHpIm
	2760) 5'W C A G C C A W-3'	PyPyImPyPyPy-γ-HpImImPyHpIm
	2761) 5'W C A G G G G W-3'	PyPyImImImIm-γ-PyPyPyPyHpIm
	2762) 5'W C A G G G C W-3'	PyPyImImImPy-γ-ImPyPyPyHpIm
30	2763) 5'W C A G G C G W-3'	PyPyImImPyIm-γ-PyImPyPyHpIm
	2764) 5'W C A G G C C W-3'	PyPyImImPyPy-γ-ImImPyPyHpIm
	2765) 5'W C A G C G G W-3'	PyPyImPyImIm-γ-PyPyImPyHpIm
	2766) 5'W C A G C G C W-3'	PyPyImPyImPy-γ-ImPyImPyHpIm
	2767) 5'W C A G C C G W-3'	PyPyImPyPyIm-γ-PyImImPyHpIm
35	2768) 5'W C A G C C C W-3'	PyPyImPyPyPy-γ-ImImImPyHpIm

TABLE 134: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCATWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2769) 5'W C A T T T T W-3'	PyPyHpHpHpHp-γ-PyPyPyPyHpIm
5	2770) 5'W C A T T T A W-3'	PyPyHpHpHpPy-γ-HpPyPyPyHpIm
	2771) 5'W C A T T T G W-3'	PyPyHpHpHpIm-γ-PyPyPyPyHpIm
	2772) 5'W C A T T T C W-3'	PyPyHpHpHpPy-γ-ImPyPyPyHpIm
	2773) 5'W C A T T A T W-3'	PyPyHpHpPyHp-γ-PyHpPyPyHpIm
	2774) 5'W C A T T A A W-3'	PyPyHpHpPyPy-γ-HpHpPyPyHpIm
10	2775) 5'W C A T T A G W-3'	PyPyHpHpPyIm-γ-PyHpPyPyHpIm
	2776) 5'W C A T T A C W-3'	PyPyHpHpPyPy-γ-ImHpPyPyHpIm
	2777) 5'W C A T T G T W-3'	PyPyHpHpImHp-γ-PyPyPyPyHpIm
	2778) 5'W C A T T G A W-3'	PyPyHpHpImPy-γ-HpPyPyPyHpIm
	2779) 5'W C A T T G G W-3'	PyPyHpHpImIm-γ-PyPyPyPyHpIm
	2780) 5'W C A T T G C W-3'	PyPyHpHpImPy-γ-ImPyPyPyHpIm
	2781) 5'W C A T T C T W-3'	PyPyHpHpPyHp-γ-PyImPyPyHpIm
	2782) 5'W C A T T C A W-3'	PyPyHpHpPyPy-γ-HpImPyPyHpIm
	2783) 5'W C A T T C G W-3'	PyPyHpHpPyIm-γ-PyImPyPyHpIm
	2784) 5'W C A T T C C W-3'	PyPyHpHpPyPy-γ-ImImPyPyHpIm
20	2785) 5'W C A T A T T W-3'	PyPyHpPyHpHp-γ-PyPyHpPyHpIm
	2786) 5'W C A T A T A W-3'	PyPyHpPyHpPy-γ-HpPyHpPyHpIm
	2787) 5'W C A T A T G W-3'	PyPyHpPyHpIm-γ-PyPyHpPyHpIm
	2788) 5'W C A T A T C W-3'	PyPyHpPyHpPy-γ-ImPyHpPyHpIm
	2789) 5'W C A T A A T W-3'	PyPyHpPyPyHp-γ-PyHpHpPyHpIm
25	2790) 5'W C A T A A A W-3'	PyPyHpPyPyPy-γ-HpHpHpPyHpIm
	2791) 5'W C A T A A G W-3'	PyPyHpPyPyIm-γ-PyHpHpPyHpIm
	2792) 5'W C A T A A C W-3'	PyPyHpPyPyPy-γ-ImHpHpPyHpIm
	2793) 5'W C A T A G T W-3'	PyPyHpPyImHp-γ-PyPyHpPyHpIm
	2794) 5'W C A T A G A W-3'	PyPyHpPyImPy-γ-HpPyHpPyHpIm
30	2795) 5'W C A T A G G W-3'	PyPyHpPyImIm-γ-PyPyHpPyHpIm
	2796) 5'W C A T A G C W-3'	PyPyHpPyImPy-γ-ImPyHpPyHpIm
	2797) 5'W C A T A C T W-3'	PyPyHpPyPyHp-γ-PyImHpPyHpIm
	2798) 5'W C A T A C A W-3'	PyPyHpPyPyPy-γ-HpImHpPyHpIm
	2799) 5'W C A T A C G W-3'	PyPyHpPyPyIm-γ-PyImHpPyHpIm
35	2800) 5'W C A T A C C W-3'	PyPyHpPyPyPy-γ-ImImHpPyHpIm

TABLE 135: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCATSNW-3'

	DNA sequence	aromatic amino acid sequence
	2801) 5'W C A T G T T W-3'	PyPyHpImHpHp- γ -PyPyPyPyHpIm
5	2802) 5'W C A T G T A W-3'	PyPyHpImHpPy- γ -HpPyPyPyHpIm
	2803) 5'W C A T G T G W-3'	PyPyHpImHpIm- γ -PyPyPyPyHpIm
	2804) 5'W C A T G T C W-3'	PyPyHpImHpPy- γ -ImPyPyPyHpIm
	2805) 5'W C A T G A T W-3'	PyPyHpImPyHp- γ -PyHpPyPyHpIm
	2806) 5'W C A T G A A W-3'	PyPyHpImPyPy- γ -HpHpPyPyHpIm
10	2807) 5'W C A T G A G W-3'	PyPyHpImPyIm- γ -PyHpPyPyHpIm
	2808) 5'W C A T G A C W-3'	PyPyHpImPyPy- γ -ImHpPyPyHpIm
	2809) 5'W C A T G G T W-3'	PyPyHpImImHp- γ -PyPyPyPyHpIm
	2810) 5'W C A T G G A W-3'	PyPyHpImImPy- γ -HpPyPyPyHpIm
	2811) 5'W C A T G C T W-3'	PyPyHpImPyHp- γ -PyImPyPyHpIm
	2812) 5'W C A T G C A W-3'	PyPyHpImPyPy- γ -HpImPyPyHpIm
	2813) 5'W C A T G G G W-3'	PyPyHpImImIm- γ -PyPyPyPyHpIm
	2814) 5'W C A T G G C W-3'	PyPyHpImImPy- γ -ImPyPyPyHpIm
	2815) 5'W C A T G C G W-3'	PyPyHpImPyIm- γ -PyImPyPyHpIm
	2816) 5'W C A T G C C W-3'	PyPyHpImPyPy- γ -ImImPyPyHpIm
	2817) 5'W C A T C T T W-3'	PyPyHpPyHpHp- γ -PyPyImPyHpIm
	2818) 5'W C A T C T A W-3'	PyPyHpPyHpPy- γ -HpPyImPyHpIm
	2819) 5'W C A T C T G W-3'	PyPyHpPyHpIm- γ -PyPyImPyHpIm
	2820) 5'W C A T C T C W-3'	PyPyHpPyHpPy- γ -ImPyImPyHpIm
	2821) 5'W C A T C A T W-3'	PyPyHpPyPyHp- γ -PyHpImPyHpIm
25	2822) 5'W C A T C A A W-3'	PyPyHpPyPyPy- γ -HpHpImPyHpIm
	2823) 5'W C A T C A G W-3'	PyPyHpPyPyIm- γ -PyHpImPyHpIm
	2824) 5'W C A T C A C W-3'	PyPyHpPyPyPy- γ -ImHpImPyHpIm
	2825) 5'W C A T C G T W-3'	PyPyHpPyImHp- γ -PyPyImPyHpIm
	2826) 5'W C A T C G A W-3'	PyPyHpPyImPy- γ -HpPyImPyHpIm
30	2827) 5'W C A T C C T W-3'	PyPyHpPyPyHp- γ -PyImImPyHpIm
	2828) 5'W C A T C C A W-3'	PyPyHpPyPyPy- γ -HpImImPyHpIm
	2829) 5'W C A T C G G W-3'	PyPyHpPyImIm- γ -PyPyImPyHpIm
	2830) 5'W C A T C G C W-3'	PyPyHpPyImPy- γ -ImPyImPyHpIm
	2831) 5'W C A T C C G W-3'	PyPyHpPyPyIm- γ -PyImImPyHpIm
35	2832) 5'W C A T C C C W-3'	PyPyHpPyPyPy- γ -ImImImPyHpIm

TABLE 136: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCAAWNNW-3'

	DNA sequence	aromatic amino acid sequence
5	2833) 5'W C A A T T T W-3'	PyPyPyHpHpHp- γ -PyPyPyHpHpIm
	2834) 5'W C A A T T A W-3'	PyPyPyHpHpPy- γ -HpPyPyHpHpIm
	2835) 5'W C A A T T G W-3'	PyPyPyHpHpIm- γ -PyPyPyHpHpIm
	2836) 5'W C A A T T C W-3'	PyPyPyHpHpPy- γ -ImPyPyHpHpIm
	2837) 5'W C A A T A T W-3'	PyPyPyHpPyHp- γ -PyHpPyHpHpIm
10	2838) 5'W C A A T A A W-3'	PyPyPyHpPyPy- γ -HpHpPyHpHpIm
	2839) 5'W C A A T A G W-3'	PyPyPyHpPyIm- γ -PyHpPyHpHpIm
	2840) 5'W C A A T A C W-3'	PyPyPyHpPyPy- γ -ImHpPyHpHpIm
	2841) 5'W C A A T G T W-3'	PyPyPyHpImHp- γ -PyPyPyHpHpIm
	2842) 5'W C A A T G A W-3'	PyPyPyHpImPy- γ -HpPyPyHpHpIm
	2843) 5'W C A A T G G W-3'	PyPyPyHpImIm- γ -PyPyPyHpHpIm
	2844) 5'W C A A T G C W-3'	PyPyPyHpImPy- γ -ImPyPyHpHpIm
	2845) 5'W C A A T C T W-3'	PyPyPyHpPyHp- γ -PyImPyHpHpIm
	2846) 5'W C A A T C A W-3'	PyPyPyHpPyPy- γ -HpImPyHpHpIm
	2847) 5'W C A A T C G W-3'	PyPyPyHpPyIm- γ -PyImPyHpHpIm
20	2848) 5'W C A A T C C W-3'	PyPyPyHpPyPy- γ -ImImPyHpHpIm
	2849) 5'W C A A A T T W-3'	PyPyPyPyHpHp- γ -PyPyHpHpHpIm
	2850) 5'W C A A A T A W-3'	PyPyPyPyHpPy- γ -HpPyHpHpHpIm
	2851) 5'W C A A A T G W-3'	PyPyPyPyHpIm- γ -PyPyHpHpHpIm
	2852) 5'W C A A A T C W-3'	PyPyPyPyHpPy- γ -ImPyHpHpHpIm
	2853) 5'W C A A A A T W-3'	PyPyPyPyPyHp- γ -PyHpHpHpHpIm
	2854) 5'W C A A A A A W-3'	PyPyPyPyPyPy- γ -HpHpHpHpHpIm
	2855) 5'W C A A A A G W-3'	PyPyPyPyPyIm- γ -PyHpHpHpHpIm
	2856) 5'W C A A A A C W-3'	PyPyPyPyPyPy- γ -ImHpHpHpHpIm
	2857) 5'W C A A A G T W-3'	PyPyPyPyImHp- γ -PyPyHpHpHpIm
30	2858) 5'W C A A A G A W-3'	PyPyPyPyImPy- γ -HpPyHpHpHpIm
	2859) 5'W C A A A G G W-3'	PyPyPyPyImIm- γ -PyPyHpHpHpIm
	2860) 5'W C A A A G C W-3'	PyPyPyPyImPy- γ -ImPyHpHpHpIm
	2861) 5'W C A A A C T W-3'	PyPyPyPyPyHp- γ -PyImHpHpHpIm
	2862) 5'W C A A A C A W-3'	PyPyPyPyPyPy- γ -HpImHpHpHpIm
	2863) 5'W C A A A C G W-3'	PyPyPyPyPyIm- γ -PyImHpHpHpIm
	2864) 5'W C A A A C C W-3'	PyPyPyPyPyPy- γ -ImImHpHpHpIm
35		

TABLE 137: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCAASNNW-3'

	DNA sequence	aromatic amino acid sequence
	2865) 5'W C A A G T T W-3'	PyPyPyImHpHp- γ -PyPyPyHpHpIm
5	2866) 5'W C A A G T A W-3'	PyPyPyImHpPy- γ -HpPyPyHpHpIm
	2867) 5'W C A A G T G W-3'	PyPyPyImHpIm- γ -PyPyPyHpHpIm
	2868) 5'W C A A G T C W-3'	PyPyPyImHpPy- γ -ImPyPyHpHpIm
	2869) 5'W C A A G A T W-3'	PyPyPyImPyHp- γ -PyHpPyHpHpIm
	2870) 5'W C A A G A A W-3'	PyPyPyImPyPy- γ -HpHpPyHpHpIm
10	2871) 5'W C A A G A G W-3'	PyPyPyImPyIm- γ -PyHpPyHpHpIm
	2872) 5'W C A A G A C W-3'	PyPyPyImPyPy- γ -ImHpPyHpHpIm
	2873) 5'W C A A G G T W-3'	PyPyPyImImHp- γ -PyPyPyHpHpIm
	2874) 5'W C A A G G A W-3'	PyPyPyImImPy- γ -HpPyPyHpHpIm
	2875) 5'W C A A G C T W-3'	PyPyPyImPyHp- γ -PyImPyHpHpIm
	2876) 5'W C A A G C A W-3'	PyPyPyImPyPy- γ -HpImPyHpHpIm
	2877) 5'W C A A G G G W-3'	PyPyPyImImIm- γ -PyPyPyHpHpIm
	2878) 5'W C A A G G C W-3'	PyPyPyImImPy- γ -ImPyPyHpHpIm
	2879) 5'W C A A G C G W-3'	PyPyPyImPyIm- γ -PyImPyHpHpIm
	2880) 5'W C A A G C C W-3'	PyPyPyImPyPy- γ -ImImPyHpHpIm
20	2881) 5'W C A A C T T W-3'	PyPyPyPyHpHp- γ -PyPyImHpHpIm
	2882) 5'W C A A C T A W-3'	PyPyPyPyHpPy- γ -HpPyImHpHpIm
	2883) 5'W C A A C T G W-3'	PyPyPyPyHpIm- γ -PyPyImHpHpIm
	2884) 5'W C A A C T C W-3'	PyPyPyPyHpPy- γ -ImPyImHpHpIm
	2885) 5'W C A A C A T W-3'	PyPyPyPyPyHp- γ -PyHpImHpHpIm
25	2886) 5'W C A A C A A W-3'	PyPyPyPyPyPy- γ -HpHpImHpHpIm
	2887) 5'W C A A C A G W-3'	PyPyPyPyPyIm- γ -PyHpImHpHpIm
	2888) 5'W C A A C A C W-3'	PyPyPyPyPyPy- γ -ImHpImHpHpIm
	2889) 5'W C A A C G T W-3'	PyPyPyPyImHp- γ -PyPyImHpHpIm
	2890) 5'W C A A C G A W-3'	PyPyPyPyImPy- γ -HpPyImHpHpIm
30	2891) 5'W C A A C C T W-3'	PyPyPyPyPyHp- γ -PyImImHpHpIm
	2892) 5'W C A A C C A W-3'	PyPyPyPyPyPy- γ -HpImImHpHpIm
	2893) 5'W C A A C G G W-3'	PyPyPyPyImIm- γ -PyPyImHpHpIm
	2894) 5'W C A A C G C W-3'	PyPyPyPyImPy- γ -ImPyImHpHpIm
	2895) 5'W C A A C C G W-3'	PyPyPyPyPyIm- γ -PyImImHpHpIm
35	2896) 5'W C A A C C C W-3'	PyPyPyPyPyPy- γ -ImImImHpHpIm

TABLE 138: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCACWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2897) 5'W C A C T T T W-3'	PyPyPyHpHpHp- γ -PyPyPyImHpIm
5	2898) 5'W C A C T T A W-3'	PyPyPyHpHpPy- γ -HpPyPyImHpIm
	2899) 5'W C A C T T G W-3'	PyPyPyHpHpIm- γ -PyPyPyImHpIm
	2900) 5'W C A C T T C W-3'	PyPyPyHpHpPy- γ -ImPyPyImHpIm
	2901) 5'W C A C T A T W-3'	PyPyPyHpPyHp- γ -PyHpPyImHpIm
	2902) 5'W C A C T A A W-3'	PyPyPyHpPyPy- γ -HpHpPyImHpIm
10	2903) 5'W C A C T A G W-3'	PyPyPyHpPyIm- γ -PyHpPyImHpIm
	2904) 5'W C A C T A C W-3'	PyPyPyHpPyPy- γ -ImHpPyImHpIm
	2905) 5'W C A C T G T W-3'	PyPyPyHpImHp- γ -PyPyPyImHpIm
	2906) 5'W C A C T G A W-3'	PyPyPyHpImPy- γ -HpPyPyImHpIm
	2907) 5'W C A C T G G W-3'	PyPyPyHpImIm- γ -PyPyPyImHpIm
	2908) 5'W C A C T G C W-3'	PyPyPyHpImPy- γ -ImPyPyImHpIm
	2909) 5'W C A C T C T W-3'	PyPyPyHpPyHp- γ -PyImPyImHpIm
	2910) 5'W C A C T C A W-3'	PyPyPyHpPyPy- γ -HpImPyImHpIm
	2911) 5'W C A C T C G W-3'	PyPyPyHpPyIm- γ -PyImPyImHpIm
	2912) 5'W C A C T C C W-3'	PyPyPyHpPyPy- γ -ImImPyImHpIm
20	2913) 5'W C A C A T T W-3'	PyPyPyPyHpHp- γ -PyPyHpImHpIm
	2914) 5'W C A C A T A W-3'	PyPyPyPyHpPy- γ -HpPyHpImHpIm
	2915) 5'W C A C A T G W-3'	PyPyPyPyHpIm- γ -PyPyHpImHpIm
	2916) 5'W C A C A T C W-3'	PyPyPyPyHpPy- γ -ImPyHpImHpIm
	2917) 5'W C A C A A T W-3'	PyPyPyPyPyHp- γ -PyHpHpImHpIm
25	2918) 5'W C A C A A A W-3'	PyPyPyPyPyPy- γ -HpHpHpImHpIm
	2919) 5'W C A C A A G W-3'	PyPyPyPyPyIm- γ -PyHpHpImHpIm
	2920) 5'W C A C A A C W-3'	PyPyPyPyPyPy- γ -ImHpHpImHpIm
	2921) 5'W C A C A G T W-3'	PyPyPyPyImHp- γ -PyPyHpImHpIm
	2922) 5'W C A C A G A W-3'	PyPyPyPyImPy- γ -HpPyHpImHpIm
30	2923) 5'W C A C A G G W-3'	PyPyPyPyImIm- γ -PyPyHpImHpIm
	2924) 5'W C A C A G C W-3'	PyPyPyPyImPy- γ -ImPyHpImHpIm
	2925) 5'W C A C A C T W-3'	PyPyPyPyPyHp- γ -PyImHpImHpIm
	2926) 5'W C A C A C A W-3'	PyPyPyPyPyPy- γ -HpImHpImHpIm
	2927) 5'W C A C A C G W-3'	PyPyPyPyPyIm- γ -PyImHpImHpIm
35	2928) 5'W C A C A C C W-3'	PyPyPyPyPyPy- γ -ImImHpImHpIm

TABLE 139: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCACNNW-3'

	DNA sequence	aromatic amino acid sequence
	2929) 5'W C A C G T T W-3'	PyPyPyImHpHp-γ-PyPyPyImHpIm
5	2930) 5'W C A C G T A W-3'	PyPyPyImHpPy-γ-HpPyPyImHpIm
	2931) 5'W C A C G T G W-3'	PyPyPyImHpIm-γ-PyPyPyImHpIm
	2932) 5'W C A C G T C W-3'	PyPyPyImHpPy-γ-ImPyPyImHpIm
	2933) 5'W C A C G A T W-3'	PyPyPyImPyHp-γ-PyHpPyImHpIm
	2934) 5'W C A C G A A W-3'	PyPyPyImPyPy-γ-HpHpPyImHpIm
10	2935) 5'W C A C G A G W-3'	PyPyPyImPyIm-γ-PyHpPyImHpIm
	2936) 5'W C A C G A C W-3'	PyPyPyImPyPy-γ-ImHpPyImHpIm
	2937) 5'W C A C G G T W-3'	PyPyPyImImHp-γ-PyPyPyImHpIm
	2938) 5'W C A C G G A W-3'	PyPyPyImImPy-γ-HpPyPyImHpIm
	2939) 5'W C A C G C T W-3'	PyPyPyImPyHp-γ-PyImPyImHpIm
	2940) 5'W C A C G C A W-3'	PyPyPyImPyPy-γ-HpImPyImHpIm
	2941) 5'W C A C C T T W-3'	PyPyPyPyHpHp-γ-PyPyImImHpIm
	2942) 5'W C A C C T A W-3'	PyPyPyPyHpPy-γ-HpPyImImHpIm
	2943) 5'W C A C C T G W-3'	PyPyPyPyHpIm-γ-PyPyImImHpIm
	2944) 5'W C A C C T C W-3'	PyPyPyPyHpPy-γ-ImPyImImHpIm
20	2945) 5'W C A C C A T W-3'	PyPyPyPyPyHp-γ-PyHpImImHpIm
	2946) 5'W C A C C A A W-3'	PyPyPyPyPyPy-γ-HpHpImImHpIm
	2947) 5'W C A C C A G W-3'	PyPyPyPyPyIm-γ-PyHpImImHpIm
	2948) 5'W C A C C A C W-3'	PyPyPyPyPyPy-γ-ImHpImImHpIm
	2949) 5'W C A C C G T W-3'	PyPyPyPyImHp-γ-PyPyImImHpIm
25	2950) 5'W C A C C G A W-3'	PyPyPyPyImPy-γ-HpPyImImHpIm
	2951) 5'W C A C C C T W-3'	PyPyPyPyPyHp-γ-PyImImImHpIm
	2952) 5'W C A C C C A W-3'	PyPyPyPyPyPy-γ-HpImImImHpIm
	2953) 5'W C A C G G G W-3'	PyPyPyImImIm-γ-PyPyPyImHpIm
	2954) 5'W C A C G G C W-3'	PyPyPyImImPy-γ-ImPyPyImHpIm
30	2955) 5'W C A C G C G W-3'	PyPyPyImPyIm-γ-PyImPyImHpIm
	2956) 5'W C A C G C C W-3'	PyPyPyImPyPy-γ-ImImPyImHpIm
	2957) 5'W C A C C G G W-3'	PyPyPyPyImIm-γ-PyPyImImHpIm
	2958) 5'W C A C C G C W-3'	PyPyPyPyImPy-γ-ImPyImImHpIm
	2959) 5'W C A C C C G W-3'	PyPyPyPyPyIm-γ-PyImImImHpIm
35	2960) 5'W C A C C C C W-3'	PyPyPyPyPyPy-γ-ImImImImHpIm

TABLE 140: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2961) 5'W C T G T T T W-3'	PyHpImHpHpHp-γ-PyPyPyPyPyIm
5	2962) 5'W C T G T T A W-3'	PyHpImHpHpPy-γ-HpPyPyPyPyIm
	2963) 5'W C T G T T G W-3'	PyHpImHpHpIm-γ-PyPyPyPyPyIm
	2964) 5'W C T G T T C W-3'	PyHpImHpHpPy-γ-ImPyPyPyPyIm
	2965) 5'W C T G T A T W-3'	PyHpImHpPyHp-γ-PyHpPyPyPyIm
	2966) 5'W C T G T A A W-3'	PyHpImHpPyPy-γ-HpHpPyPyPyIm
10	2967) 5'W C T G T A G W-3'	PyHpImHpPyIm-γ-PyHpPyPyPyIm
	2968) 5'W C T G T A C W-3'	PyHpImHpPyPy-γ-ImHpPyPyPyIm
	2969) 5'W C T G T G T W-3'	PyHpImHpImHp-γ-PyPyPyPyPyIm
	2970) 5'W C T G T G A W-3'	PyHpImHpImPy-γ-HpPyPyPyPyIm
	2971) 5'W C T G T G G W-3'	PyHpImHpImIm-γ-PyPyPyPyPyIm
15	2972) 5'W C T G T G C W-3'	PyHpImHpImPy-γ-ImPyPyPyPyIm
	2973) 5'W C T G T C T W-3'	PyHpImHpPyHp-γ-PyImPyPyPyIm
	2974) 5'W C T G T C A W-3'	PyHpImHpPyPy-γ-HpImPyPyPyIm
	2975) 5'W C T G T C G W-3'	PyHpImHpPyIm-γ-PyImPyPyPyIm
	2976) 5'W C T G T C C W-3'	PyHpImHpPyPy-γ-ImImPyPyPyIm
20	2977) 5'W C T G A T T W-3'	PyHpImPyHpHp-γ-PyPyHpPyPyIm
	2978) 5'W C T G A T A W-3'	PyHpImPyHpPy-γ-HpPyHpPyPyIm
	2979) 5'W C T G A T G W-3'	PyHpImPyHpIm-γ-PyPyHpPyPyIm
	2980) 5'W C T G A T C W-3'	PyHpImPyHpPy-γ-ImPyHpPyPyIm
	2981) 5'W C T G A A T W-3'	PyHpImPyPyHp-γ-PyHpHpPyPyIm
25	2982) 5'W C T G A A A W-3'	PyHpImPyPyPy-γ-HpHpHpPyPyIm
	2983) 5'W C T G A A G W-3'	PyHpImPyPyIm-γ-PyHpHpPyPyIm
	2984) 5'W C T G A A C W-3'	PyHpImPyPyPy-γ-ImHpHpPyPyIm
	2985) 5'W C T G A G T W-3'	PyHpImPyImHp-γ-PyPyHpPyPyIm
	2986) 5'W C T G A G A W-3'	PyHpImPyImPy-γ-HpPyHpPyPyIm
30	2987) 5'W C T G A G G W-3'	PyHpImPyImIm-γ-PyPyHpPyPyIm
	2988) 5'W C T G A G C W-3'	PyHpImPyImPy-γ-ImPyHpPyPyIm
	2989) 5'W C T G A C T W-3'	PyHpImPyPyHp-γ-PyImHpPyPyIm
	2990) 5'W C T G A C A W-3'	PyHpImPyPyPy-γ-HpImHpPyPyIm
	2991) 5'W C T G A C G W-3'	PyHpImPyPyIm-γ-PyImHpPyPyIm
35	2992) 5'W C T G A C C W-3'	PyHpImPyPyPy-γ-ImImHpPyPyIm

TABLE 141: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTGSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	2993) 5'W C T G G T T W-3'	PyHpImImHpHp-γ-PyPyPyPyPyIm
	2994) 5'W C T G G T A W-3'	PyHpImImHpPy-γ-HpPyPyPyPyIm
	2995) 5'W C T G G T G W-3'	PyHpImImHpIm-γ-PyPyPyPyPyIm
	2996) 5'W C T G G T C W-3'	PyHpImImHpPy-γ-ImPyPyPyPyIm
	2997) 5'W C T G G A T W-3'	PyHpImImPyHp-γ-PyHpPyPyPyIm
	2998) 5'W C T G G A A W-3'	PyHpImImPyPy-γ-HpHpPyPyPyIm
10	2999) 5'W C T G G A G W-3'	PyHpImImPyIm-γ-PyHpPyPyPyIm
	3000) 5'W C T G G A C W-3'	PyHpImImPyPy-γ-ImHpPyPyPyIm
	3001) 5'W C T G G G T W-3'	PyHpImImImHp-γ-PyPyPyPyPyIm
	3002) 5'W C T G G G A W-3'	PyHpImImImPy-γ-HpPyPyPyPyIm
	3003) 5'W C T G G C T W-3'	PyHpImImPyHp-γ-PyImPyPyPyIm
	3004) 5'W C T G G C A W-3'	PyHpImImPyPy-γ-HpImPyPyPyIm
	3005) 5'W C T G C T T W-3'	PyHpImPyHpHp-γ-PyPyImPyPyIm
	3006) 5'W C T G C T A W-3'	PyHpImPyHpPy-γ-HpPyImPyPyIm
	3007) 5'W C T G C T G W-3'	PyHpImPyHpIm-γ-PyPyImPyPyIm
	3008) 5'W C T G C T C W-3'	PyHpImPyHpPy-γ-ImPyImPyPyIm
20	3009) 5'W C T G C A T W-3'	PyHpImPyPyHp-γ-PyHpImPyPyIm
	3010) 5'W C T G C A A W-3'	PyHpImPyPyPy-γ-HpHpImPyPyIm
	3011) 5'W C T G C A G W-3'	PyHpImPyPyIm-γ-PyHpImPyPyIm
	3012) 5'W C T G C A C W-3'	PyHpImPyPyPy-γ-ImHpImPyPyIm
	3013) 5'W C T G C G T W-3'	PyHpImPyImHp-γ-PyPyImPyPyIm
25	3014) 5'W C T G C G A W-3'	PyHpImPyImPy-γ-HpPyImPyPyIm
	3015) 5'W C T G C C T W-3'	PyHpImPyPyHp-γ-PyImImPyPyIm
	3016) 5'W C T G C C A W-3'	PyHpImPyPyPy-γ-HpImImPyPyIm
	3017) 5'W C T G G G G W-3'	PyHpImImImIm-γ-PyPyPyPyPyIm
	3018) 5'W C T G G G C W-3'	PyHpImImImPy-γ-ImPyPyPyPyIm
30	3019) 5'W C T G G C G W-3'	PyHpImImPyIm-γ-PyImPyPyPyIm
	3020) 5'W C T G G C C W-3'	PyHpImImPyPy-γ-ImImPyPyPyIm
	3021) 5'W C T G C G G W-3'	PyHpImPyImIm-γ-PyPyImPyPyIm
	3022) 5'W C T G C G C W-3'	PyHpImPyImPy-γ-ImPyImPyPyIm
	3023) 5'W C T G C C G W-3'	PyHpImPyPyIm-γ-PyImImPyPyIm
35	3024) 5'W C T G C C C W-3'	PyHpImPyPyPy-γ-ImImImPyPyIm

TABLE 142: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	3025) 5'W C T T T T T W-3'	PyHpHpHpHpHp-γ-PyPyPyPyPyIm
5	3026) 5'W C T T T T A W-3'	PyHpHpHpHpPy-γ-HpPyPyPyPyIm
	3027) 5'W C T T T T G W-3'	PyHpHpHpHpIm-γ-PyPyPyPyPyIm
	3028) 5'W C T T T T C W-3'	PyHpHpHpHpPy-γ-ImPyPyPyPyIm
	3029) 5'W C T T T A T W-3'	PyHpHpHpPyHp-γ-PyHpPyPyPyIm
	3030) 5'W C T T T A A W-3'	PyHpHpHpPyPy-γ-HpHpPyPyPyIm
10	3031) 5'W C T T T A G W-3'	PyHpHpHpPyIm-γ-PyHpPyPyPyIm
	3032) 5'W C T T T A C W-3'	PyHpHpHpPyPy-γ-ImHpPyPyPyIm
	3033) 5'W C T T T G T W-3'	PyHpHpHpImHp-γ-PyPyPyPyPyIm
	3034) 5'W C T T T G A W-3'	PyHpHpHpImPy-γ-HpPyPyPyPyIm
	3035) 5'W C T T T G G W-3'	PyHpHpHpImIm-γ-PyPyPyPyPyIm
15	3036) 5'W C T T T G C W-3'	PyHpHpHpImPy-γ-ImPyPyPyPyIm
	3037) 5'W C T T T C T W-3'	PyHpHpHpPyHp-γ-PyImPyPyPyIm
	3038) 5'W C T T T C A W-3'	PyHpHpHpPyPy-γ-HpImPyPyPyIm
	3039) 5'W C T T T C G W-3'	PyHpHpHpPyIm-γ-PyImPyPyPyIm
	3040) 5'W C T T T C C W-3'	PyHpHpHpPyPy-γ-ImImPyPyPyIm
20	3041) 5'W C T T A T T W-3'	PyHpHpPyHpHp-γ-PyPyHpPyPyIm
	3042) 5'W C T T A T A W-3'	PyHpHpPyHpPy-γ-HpPyHpPyPyIm
	3043) 5'W C T T A T G W-3'	PyHpHpPyHpIm-γ-PyPyHpPyPyIm
	3044) 5'W C T T A T C W-3'	PyHpHpPyHpPy-γ-ImPyHpPyPyIm
	3045) 5'W C T T A A T W-3'	PyHpHpPyPyHp-γ-PyHpHpPyPyIm
25	3046) 5'W C T T A A A W-3'	PyHpHpPyPyPy-γ-HpHpHpPyPyIm
	3047) 5'W C T T A A G W-3'	PyHpHpPyPyIm-γ-PyHpHpPyPyIm
	3048) 5'W C T T A A C W-3'	PyHpHpPyPyPy-γ-ImHpHpPyPyIm
	3049) 5'W C T T A G T W-3'	PyHpHpPyImHp-γ-PyPyHpPyPyIm
	3050) 5'W C T T A G A W-3'	PyHpHpPyImPy-γ-HpPyHpPyPyIm
30	3051) 5'W C T T A G G W-3'	PyHpHpPyImIm-γ-PyPyHpPyPyIm
	3052) 5'W C T T A G C W-3'	PyHpHpPyImPy-γ-ImPyHpPyPyIm
	3053) 5'W C T T A C T W-3'	PyHpHpPyPyHp-γ-PyImHpPyPyIm
	3054) 5'W C T T A C A W-3'	PyHpHpPyPyPy-γ-HpImHpPyPyIm
	3055) 5'W C T T A C G W-3'	PyHpHpPyPyIm-γ-PyImHpPyPyIm
35	3056) 5'W C T T A C C W-3'	PyHpHpPyPyPy-γ-ImImHpPyPyIm

TABLE 143: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	3057) 5'W C T T G T T W-3'	PyHpHpImHpHp- γ -PyPyPyPyPyIm
5	3058) 5'W C T T G T A W-3'	PyHpHpImHpPy- γ -HpPyPyPyPyIm
	3059) 5'W C T T G T G W-3'	PyHpHpImHpIm- γ -PyPyPyPyPyIm
	3060) 5'W C T T G T C W-3'	PyHpHpImHpPy- γ -ImPyPyPyPyIm
	3061) 5'W C T T G A T W-3'	PyHpHpImPyHp- γ -PyHpPyPyPyIm
	3062) 5'W C T T G A A W-3'	PyHpHpImPyPy- γ -HpHpPyPyPyIm
10	3063) 5'W C T T G A G W-3'	PyHpHpImPyIm- γ -PyHpPyPyPyIm
	3064) 5'W C T T G A C W-3'	PyHpHpImPyPy- γ -ImHpPyPyPyIm
	3065) 5'W C T T G G T W-3'	PyHpHpImImHp- γ -PyPyPyPyPyIm
	3066) 5'W C T T G G A W-3'	PyHpHpImImPy- γ -HpPyPyPyPyIm
	3067) 5'W C T T G C T W-3'	PyHpHpImPyHp- γ -PyImPyPyPyIm
	3068) 5'W C T T G C A W-3'	PyHpHpImPyPy- γ -HpImPyPyPyIm
	3069) 5'W C T T G G G W-3'	PyHpHpImImIm- γ -PyPyPyPyPyIm
	3070) 5'W C T T G G C W-3'	PyHpHpImImPy- γ -ImPyPyPyPyIm
	3071) 5'W C T T G C G W-3'	PyHpHpImPyIm- γ -PyImPyPyPyIm
	3072) 5'W C T T G C C W-3'	PyHpHpImPyPy- γ -ImImPyPyPyIm
20	3073) 5'W C T T C T T W-3'	PyHpHpPyHpHp- γ -PyPyImPyPyIm
	3074) 5'W C T T C T A W-3'	PyHpHpPyHpPy- γ -HpPyImPyPyIm
	3075) 5'W C T T C T G W-3'	PyHpHpPyHpIm- γ -PyPyImPyPyIm
	3076) 5'W C T T C T C W-3'	PyHpHpPyHpPy- γ -ImPyImPyPyIm
	3077) 5'W C T T C A T W-3'	PyHpHpPyPyHp- γ -PyHpImPyPyIm
25	3078) 5'W C T T C A A W-3'	PyHpHpPyPyPy- γ -HpHpImPyPyIm
	3079) 5'W C T T C A G W-3'	PyHpHpPyPyIm- γ -PyHpImPyPyIm
	3080) 5'W C T T C A C W-3'	PyHpHpPyPyPy- γ -ImHpImPyPyIm
	3081) 5'W C T T C G T W-3'	PyHpHpPyImHp- γ -PyPyImPyPyIm
	3082) 5'W C T T C G A W-3'	PyHpHpPyImPy- γ -HpPyImPyPyIm
30	3083) 5'W C T T C C T W-3'	PyHpHpPyPyHp- γ -PyImImPyPyIm
	3084) 5'W C T T C C A W-3'	PyHpHpPyPyPy- γ -HpImImPyPyIm
	3085) 5'W C T T C G G W-3'	PyHpHpPyImIm- γ -PyPyImPyPyIm
	3086) 5'W C T T C G C W-3'	PyHpHpPyImPy- γ -ImPyImPyPyIm
	3087) 5'W C T T C C G W-3'	PyHpHpPyPyIm- γ -PyImImPyPyIm
35	3088) 5'W C T T C C C W-3'	PyHpHpPyPyPy- γ -ImImImPyPyIm

TABLE 144: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	3089) 5'W C T A T T T W-3'	PyHpPyHpHpHp-γ-PyPyPyHpPyIm
5	3090) 5'W C T A T T A W-3'	PyHpPyHpHpPy-γ-HpPyPyHpPyIm
	3091) 5'W C T A T T G W-3'	PyHpPyHpHpIm-γ-PyPyPyHpPyIm
	3092) 5'W C T A T T C W-3'	PyHpPyHpHpPy-γ-ImPyPyHpPyIm
	3093) 5'W C T A T A T W-3'	PyHpPyHpPyHp-γ-PyHpPyHpPyIm
	3094) 5'W C T A T A A W-3'	PyHpPyHpPyPy-γ-HpHpPyHpPyIm
10	3095) 5'W C T A T A G W-3'	PyHpPyHpPyIm-γ-PyHpPyHpPyIm
	3096) 5'W C T A T A C W-3'	PyHpPyHpPyPy-γ-ImHpPyHpPyIm
	3097) 5'W C T A T G T W-3'	PyHpPyHpImHp-γ-PyPyPyHpPyIm
	3098) 5'W C T A T G A W-3'	PyHpPyHpImPy-γ-HpPyPyHpPyIm
	3099) 5'W C T A T G G W-3'	PyHpPyHpImIm-γ-PyPyPyHpPyIm
	3100) 5'W C T A T G C W-3'	PyHpPyHpImPy-γ-ImPyPyHpPyIm
	3101) 5'W C T A T C T W-3'	PyHpPyHpPyHp-γ-PyImPyHpPyIm
	3102) 5'W C T A T C A W-3'	PyHpPyHpPyPy-γ-HpImPyHpPyIm
	3103) 5'W C T A T C G W-3'	PyHpPyHpPyIm-γ-PyImPyHpPyIm
	3104) 5'W C T A T C C W-3'	PyHpPyHpPyPy-γ-ImImPyHpPyIm
20	3105) 5'W C T A A T T W-3'	PyHpPyPyHpHp-γ-PyPyHpHpPyIm
	3106) 5'W C T A A T A W-3'	PyHpPyPyHpPy-γ-HpPyHpHpPyIm
	3107) 5'W C T A A T G W-3'	PyHpPyPyHpIm-γ-PyPyHpHpPyIm
	3108) 5'W C T A A T C W-3'	PyHpPyPyHpPy-γ-ImPyHpHpPyIm
	3109) 5'W C T A A A T W-3'	PyHpPyPyPyHp-γ-PyHpHpHpPyIm
25	3110) 5'W C T A A A A W-3'	PyHpPyPyPyPy-γ-HpHpHpHpPyIm
	3111) 5'W C T A A A G W-3'	PyHpPyPyPyIm-γ-PyHpHpHpPyIm
	3112) 5'W C T A A A C W-3'	PyHpPyPyPyPy-γ-ImHpHpHpPyIm
	3113) 5'W C T A A G T W-3'	PyHpPyPyImHp-γ-PyPyHpHpPyIm
	3114) 5'W C T A A G A W-3'	PyHpPyPyImPy-γ-HpPyHpHpPyIm
30	3115) 5'W C T A A G G W-3'	PyHpPyPyImIm-γ-PyPyHpHpPyIm
	3116) 5'W C T A A G C W-3'	PyHpPyPyImPy-γ-ImPyHpHpPyIm
	3117) 5'W C T A A C T W-3'	PyHpPyPyPyHp-γ-PyImHpHpPyIm
	3118) 5'W C T A A C A W-3'	PyHpPyPyPyPy-γ-HpImHpHpPyIm
	3119) 5'W C T A A C G W-3'	PyHpPyPyPyIm-γ-PyImHpHpPyIm
35	3120) 5'W C T A A C C W-3'	PyHpPyPyPyPy-γ-ImImHpHpPyIm

TABLE 145: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTASNNW-3'

	DNA sequence	aromatic amino acid sequence
	3121) 5'W C T A G T T W-3'	PyHpPyImHpHp- γ -PyPyPyHpPyIm
5	3122) 5'W C T A G T A W-3'	PyHpPyImHpPy- γ -HpPyPyHpPyIm
	3123) 5'W C T A G T G W-3'	PyHpPyImHpIm- γ -PyPyPyHpPyIm
	3124) 5'W C T A G T C W-3'	PyHpPyImHpPy- γ -ImPyPyHpPyIm
	3125) 5'W C T A G A T W-3'	PyHpPyImPyHp- γ -PyHpPyHpPyIm
	3126) 5'W C T A G A A W-3'	PyHpPyImPyPy- γ -HpHpPyHpPyIm
10	3127) 5'W C T A G A G W-3'	PyHpPyImPyIm- γ -PyHpPyHpPyIm
	3128) 5'W C T A G A C W-3'	PyHpPyImPyPy- γ -ImHpPyHpPyIm
	3129) 5'W C T A G G T W-3'	PyHpPyImImHp- γ -PyPyPyHpPyIm
	3130) 5'W C T A G G A W-3'	PyHpPyImImPy- γ -HpPyPyHpPyIm
	3131) 5'W C T A G C T W-3'	PyHpPyImPyHp- γ -PyImPyHpPyIm
	3132) 5'W C T A G C A W-3'	PyHpPyImPyPy- γ -HpImPyHpPyIm
	3133) 5'W C T A G G G W-3'	PyHpPyImImIm- γ -PyPyPyHpPyIm
	3134) 5'W C T A G G C W-3'	PyHpPyImImPy- γ -ImPyPyHpPyIm
	3135) 5'W C T A G C G W-3'	PyHpPyImPyIm- γ -PyImPyHpPyIm
	3136) 5'W C T A G C C W-3'	PyHpPyImPyPy- γ -ImImPyHpPyIm
20	3137) 5'W C T A C T T W-3'	PyHpPyPyHpHp- γ -PyPyImHpPyIm
	3138) 5'W C T A C T A W-3'	PyHpPyPyHpPy- γ -HpPyImHpPyIm
	3139) 5'W C T A C T G W-3'	PyHpPyPyHpIm- γ -PyPyImHpPyIm
	3140) 5'W C T A C T C W-3'	PyHpPyPyHpPy- γ -ImPyImHpPyIm
	3141) 5'W C T A C A T W-3'	PyHpPyPyPyHp- γ -PyHpImHpPyIm
25	3142) 5'W C T A C A A W-3'	PyHpPyPyPyPy- γ -HpHpImHpPyIm
	3143) 5'W C T A C A G W-3'	PyHpPyPyPyIm- γ -PyHpImHpPyIm
	3144) 5'W C T A C A C W-3'	PyHpPyPyPyPy- γ -ImHpImHpPyIm
	3145) 5'W C T A C G T W-3'	PyHpPyPyImHp- γ -PyPyImHpPyIm
	3146) 5'W C T A C G A W-3'	PyHpPyPyImPy- γ -HpPyImHpPyIm
30	3147) 5'W C T A C C T W-3'	PyHpPyPyPyHp- γ -PyImImHpPyIm
	3148) 5'W C T A C C A W-3'	PyHpPyPyPyPy- γ -HpImImHpPyIm
	3149) 5'W C T A C G G W-3'	PyHpPyPyImIm- γ -PyPyImHpPyIm
	3150) 5'W C T A C G C W-3'	PyHpPyPyImPy- γ -ImPyImHpPyIm
	3151) 5'W C T A C C G W-3'	PyHpPyPyPyIm- γ -PyImImHpPyIm
35	3152) 5'W C T A C C C W-3'	PyHpPyPyPyPy- γ -ImImImHpPyIm

TABLE 146: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	3153) 5'W C T C T T T W-3'	PyHpPyHpHpHp-γ-PyPyPyImPyIm
5	3154) 5'W C T C T T A W-3'	PyHpPyHpHpPy-γ-HpPyPyImPyIm
	3155) 5'W C T C T T G W-3'	PyHpPyHpHpIm-γ-PyPyPyImPyIm
	3156) 5'W C T C T T C W-3'	PyHpPyHpHpPy-γ-ImPyPyImPyIm
	3157) 5'W C T C T A T W-3'	PyHpPyHpPyHp-γ-PyHpPyImPyIm
	3158) 5'W C T C T A A W-3'	PyHpPyHpPyPy-γ-HpHpPyImPyIm
10	3159) 5'W C T C T A G W-3'	PyHpPyHpPyIm-γ-PyHpPyImPyIm
	3160) 5'W C T C T A C W-3'	PyHpPyHpPyPy-γ-ImHpPyImPyIm
	3161) 5'W C T C T G T W-3'	PyHpPyHpImHp-γ-PyPyPyImPyIm
	3162) 5'W C T C T G A W-3'	PyHpPyHpImPy-γ-HpPyPyImPyIm
	3163) 5'W C T C T G G W-3'	PyHpPyHpImIm-γ-PyPyPyImPyIm
	3164) 5'W C T C T G C W-3'	PyHpPyHpImPy-γ-ImPyPyImPyIm
	3165) 5'W C T C T C T W-3'	PyHpPyHpPyHp-γ-PyImPyImPyIm
	3166) 5'W C T C T C A W-3'	PyHpPyHpPyPy-γ-HpImPyImPyIm
	3167) 5'W C T C T C G W-3'	PyHpPyHpPyIm-γ-PyImPyImPyIm
	3168) 5'W C T C T C C W-3'	PyHpPyHpPyPy-γ-ImImPyImPyIm
20	3169) 5'W C T C A T T W-3'	PyHpPyPyHpHp-γ-PyPyHpImPyIm
	3170) 5'W C T C A T A W-3'	PyHpPyPyHpPy-γ-HpPyHpImPyIm
	3171) 5'W C T C A T G W-3'	PyHpPyPyHpIm-γ-PyPyHpImPyIm
	3172) 5'W C T C A T C W-3'	PyHpPyPyHpPy-γ-ImPyHpImPyIm
	3173) 5'W C T C A A T W-3'	PyHpPyPyPyHp-γ-PyHpHpImPyIm
25	3174) 5'W C T C A A A W-3'	PyHpPyPyPyPy-γ-HpHpHpImPyIm
	3175) 5'W C T C A A G W-3'	PyHpPyPyPyIm-γ-PyHpHpImPyIm
	3176) 5'W C T C A A C W-3'	PyHpPyPyPyPy-γ-ImHpHpImPyIm
	3177) 5'W C T C A G T W-3'	PyHpPyPyImHp-γ-PyPyHpImPyIm
	3178) 5'W C T C A G A W-3'	PyHpPyPyImPy-γ-HpPyHpImPyIm
30	3179) 5'W C T C A G G W-3'	PyHpPyPyImIm-γ-PyPyHpImPyIm
	3180) 5'W C T C A G C W-3'	PyHpPyPyImPy-γ-ImPyHpImPyIm
	3181) 5'W C T C A C T W-3'	PyHpPyPyPyHp-γ-PyImHpImPyIm
	3182) 5'W C T C A C A W-3'	PyHpPyPyPyPy-γ-HpImHpImPyIm
	3183) 5'W C T C A C G W-3'	PyHpPyPyPyIm-γ-PyImHpImPyIm
35	3184) 5'W C T C A C C W-3'	PyHpPyPyPyPy-γ-ImImHpImPyIm

TABLE 147: 12-ring Hairpin Polyamides for recognition of 8-bp 5'-WCTCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	3185) 5'W C T C G T T W-3'	PyHpPyImHpHp- γ -PyPyPyImPyIm
5	3186) 5'W C T C G T A W-3'	PyHpPyImHpPy- γ -HpPyPyImPyIm
	3187) 5'W C T C G T G W-3'	PyHpPyImHpIm- γ -PyPyPyImPyIm
	3188) 5'W C T C G T C W-3'	PyHpPyImHpPy- γ -ImPyPyImPyIm
	3189) 5'W C T C G A T W-3'	PyHpPyImPyHp- γ -PyHpPyImPyIm
	3190) 5'W C T C G A A W-3'	PyHpPyImPyPy- γ -HpHpPyImPyIm
10	3191) 5'W C T C G A G W-3'	PyHpPyImPyIm- γ -PyHpPyImPyIm
	3192) 5'W C T C G A C W-3'	PyHpPyImPyPy- γ -ImHpPyImPyIm
	3193) 5'W C T C G G T W-3'	PyHpPyImImHp- γ -PyPyPyImPyIm
	3194) 5'W C T C G G A W-3'	PyHpPyImImPy- γ -HpPyPyImPyIm
	3195) 5'W C T C G C T W-3'	PyHpPyImPyHp- γ -PyImPyImPyIm
15	3196) 5'W C T C G C A W-3'	PyHpPyImPyPy- γ -HpImPyImPyIm
	3197) 5'W C T C C T T W-3'	PyHpPyPyHpHp- γ -PyPyImImPyIm
	3198) 5'W C T C C T A W-3'	PyHpPyPyHpPy- γ -HpPyImImPyIm
	3199) 5'W C T C C T G W-3'	PyHpPyPyHpIm- γ -PyPyImImPyIm
	3200) 5'W C T C C T C W-3'	PyHpPyPyHpPy- γ -ImPyImImPyIm
20	3201) 5'W C T C C A T W-3'	PyHpPyPyPyHp- γ -PyHpImImPyIm
	3202) 5'W C T C C A A W-3'	PyHpPyPyPyPy- γ -HpHpImImPyIm
	3203) 5'W C T C C A G W-3'	PyHpPyPyPyIm- γ -PyHpImImPyIm
	3204) 5'W C T C C A C W-3'	PyHpPyPyPyPy- γ -ImHpImImPyIm
	3205) 5'W C T C C G T W-3'	PyHpPyPyImHp- γ -PyPyImImPyIm
25	3206) 5'W C T C C G A W-3'	PyHpPyPyImPy- γ -HpPyImImPyIm
	3207) 5'W C T C C C T W-3'	PyHpPyPyPyHp- γ -PyImImImPyIm
	3208) 5'W C T C C C A W-3'	PyHpPyPyPyPy- γ -HpImImImPyIm
	3209) 5'W C T C G G G W-3'	PyHpPyImImIm- γ -PyPyPyImPyIm
	3210) 5'W C T C G G C W-3'	PyHpPyImImPy- γ -ImPyPyImPyIm
30	3211) 5'W C T C G C G W-3'	PyHpPyImPyIm- γ -PyImPyImPyIm
	3212) 5'W C T C G C C W-3'	PyHpPyImPyPy- γ -ImImPyImPyIm
	3213) 5'W C T C C G G W-3'	PyHpPyPyImIm- γ -PyPyImImPyIm
	3214) 5'W C T C C G C W-3'	PyHpPyPyImPy- γ -ImPyImImPyIm
	3215) 5'W C T C C C G W-3'	PyHpPyPyPyIm- γ -PyImImImPyIm
35	3216) 5'W C T C C C C W-3'	PyHpPyPyPyPy- γ -ImImImImPyIm

TABLE 148: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1233 β) 5'-W G G G T T T W-3'	ImImIm- β -HpHp- γ -PyPy- β -PyPyPy
5	1234 β) 5'-W G G G T T A W-3'	ImImIm- β -HpPy- γ -HpPy- β -PyPyPy
	1235 β) 5'-W G G G T T G W-3'	ImImIm- β -HpIm- γ -PyPy- β -PyPyPy
	1236 β) 5'-W G G G T T C W-3'	ImImIm- β -HpPy- γ -ImPy- β -PyPyPy
	1237 β) 5'-W G G G T A T W-3'	ImImIm- β -PyHp- γ -PyHp- β -PyPyPy
	1238 β) 5'-W G G G T A A W-3'	ImImIm- β -PyPy- γ -HpHp- β -PyPyPy
10	1239 β) 5'-W G G G T A G W-3'	ImImIm- β -PyIm- γ -PyHp- β -PyPyPy
	1240 β) 5'-W G G G T A C W-3'	ImImIm- β -PyPy- γ -ImHp- β -PyPyPy
	1241 β) 5'-W G G G T G T W-3'	ImImIm- β -ImHp- γ -PyPy- β -PyPyPy
	1242 β) 5'-W G G G T G A W-3'	ImImIm- β -ImPy- γ -HpPy- β -PyPyPy
	1243 β) 5'-W G G G T G G W-3'	ImImIm- β -ImIm- γ -PyPy- β -PyPyPy
	1244 β) 5'-W G G G T G C W-3'	ImImIm- β -ImPy- γ -ImPy- β -PyPyPy
	1245 β) 5'-W G G G T C T W-3'	ImImIm- β -PyHp- γ -PyIm- β -PyPyPy
	1246 β) 5'-W G G G T C A W-3'	ImImIm- β -PyPy- γ -HpIm- β -PyPyPy
	1247 β) 5'-W G G G T C G W-3'	ImImIm- β -PyIm- γ -PyIm- β -PyPyPy
	1248 β) 5'-W G G G T C C W-3'	ImImIm- β -PyPy- γ -ImIm- β -PyPyPy
	1249 β) 5'-W G G G A T T W-3'	ImImIm- β -HpHp- γ -PyPy- β -PyPyPy
	1250 β) 5'-W G G G A T A W-3'	ImImIm- β -HpPy- γ -HpPy- β -PyPyPy
	1251 β) 5'-W G G G A T G W-3'	ImImIm- β -HpIm- γ -PyPy- β -PyPyPy
	1252 β) 5'-W G G G A T C W-3'	ImImIm- β -HpPy- γ -ImPy- β -PyPyPy
	1253 β) 5'-W G G G A A T W-3'	ImImIm- β -PyHp- γ -PyHp- β -PyPyPy
25	1254 β) 5'-W G G G A A A W-3'	ImImIm- β -PyPy- γ -HpHp- β -PyPyPy
	1255 β) 5'-W G G G A A G W-3'	ImImIm- β -PyIm- γ -PyHp- β -PyPyPy
	1256 β) 5'-W G G G A A C W-3'	ImImIm- β -PyPy- γ -ImHp- β -PyPyPy
	1257 β) 5'-W G G G A G T W-3'	ImImIm- β -ImHp- γ -PyPy- β -PyPyPy
	1258 β) 5'-W G G G A G A W-3'	ImImIm- β -ImPy- γ -HpPy- β -PyPyPy
30	1259 β) 5'-W G G G A G G W-3'	ImImIm- β -ImIm- γ -PyPy- β -PyPyPy
	1260 β) 5'-W G G G A G C W-3'	ImImIm- β -ImPy- γ -ImPy- β -PyPyPy
	1261 β) 5'-W G G G A C T W-3'	ImImIm- β -PyHp- γ -PyIm- β -PyPyPy
	1262 β) 5'-W G G G A C A W-3'	ImImIm- β -PyPy- γ -HpIm- β -PyPyPy
	1263 β) 5'-W G G G A C G W-3'	ImImIm- β -PyIm- γ -PyIm- β -PyPyPy
35	1264 β) 5'-W G G G A C C W-3'	ImImIm- β -PyPy- γ -ImIm- β -PyPyPy

TABLE 149: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGGSNNW-3'

	DNA sequence	aromatic amino acid sequence
5	1265 β) 5'-W G G G G T T W-3'	ImImImIm- β -Hp- γ -Py- β -PyPyPyPy
	1266 β) 5'-W G G G G T A W-3'	ImImImIm- β -Py- γ -Hp- β -PyPyPyPy
	1267 β) 5'-W G G G G T G W-3'	ImImImIm- β -Im- γ -Py- β -PyPyPyPy
	1268 β) 5'-W G G G G T C W-3'	ImImImIm- β -Py- γ -Im- β -PyPyPyPy
	1269 β) 5'-W G G G G A T W-3'	ImImImIm- β -Hp- γ -Py- β -PyPyPyPy
10	1270 β) 5'-W G G G G A A W-3'	ImImImIm- β -Py- γ -Hp- β -PyPyPyPy
	1271 β) 5'-W G G G G A G W-3'	ImImImIm- β -Im- γ -Py- β -PyPyPyPy
	1272 β) 5'-W G G G G A C W-3'	ImImImIm- β -Py- γ -Im- β -PyPyPyPy
	1275 β) 5'-W G G G G C T W-3'	ImImImIm- β -Hp- γ -PyImPy- β -PyPy
	1276 β) 5'-W G G G G C A W-3'	ImImImIm- β -Py- γ -HpImPy- β -PyPy
	1277 β) 5'-W G G G C T T W-3'	ImImIm- β -HpHp- γ -PyPyIm- β -PyPy
	1278 β) 5'-W G G G C T A W-3'	ImImIm- β -HpPy- γ -HpPyIm- β -PyPy
	1279 β) 5'-W G G G C T G W-3'	ImImIm- β -HpIm- γ -PyPyIm- β -PyPy
	1280 β) 5'-W G G G C T C W-3'	ImImIm- β -HpPy- γ -ImPyIm- β -PyPy
	1281 β) 5'-W G G G C A T W-3'	ImImIm- β -PyHp- γ -PyHpIm- β -PyPy
20	1282 β) 5'-W G G G C A A W-3'	ImImIm- β -PyPy- γ -HpHpIm- β -PyPy
	1283 β) 5'-W G G G C A G W-3'	ImImIm- β -PyIm- γ -PyHpIm- β -PyPy
	1284 β) 5'-W G G G C A C W-3'	ImImIm- β -PyPy- γ -ImHpIm- β -PyPy
	1285 β) 5'-W G G G C G T W-3'	ImImIm- β -ImHp- γ -PyPyIm- β -PyPy
	1286 β) 5'-W G G G C G A W-3'	ImImIm- β -ImPy- γ -HpPyIm- β -PyPy
	1287 β) 5'-W G G G C C T W-3'	ImImIm- β -PyHp- γ -PyImIm- β -PyPy
	1288 β) 5'-W G G G C C A W-3'	ImImIm- β -PyPy- γ -HpImIm- β -PyPy
	G52 β) 5'-W G G G G C C W-3'	ImImImIm- β -Py- γ -ImImPy- β -PyPy
	G53 β) 5'-W G G G C G G W-3'	ImImIm- β -ImIm- γ -PyPyIm- β -PyPy
	G54 β) 5'-W G G G C G C W-3'	ImImIm- β -ImPy- γ -ImPyIm- β -PyPy
30	G55 β) 5'-W G G G C C G W-3'	ImImIm- β -PyIm- γ -PyImIm- β -PyPy
	G56 β) 5'-W G G G C C C W-3'	ImImIm- β -PyPy- γ -ImImIm- β -PyPy

TABLE 150: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1289 β) 5'-W G G T T T W-3'	ImIm- β -HpHpHp- γ -PyPyPy- β -PyPy
	1290 β) 5'-W G G T T T A W-3'	ImIm- β -HpHpPy- γ -HpPyPy- β -PyPy
5	1291 β) 5'-W G G T T T G W-3'	ImIm- β -HpHpIm- γ -PyPyPy- β -PyPy
	1292 β) 5'-W G G T T T C W-3'	ImIm- β -HpHpPy- γ -ImPyPy- β -PyPy
	1293 β) 5'-W G G T T A T W-3'	ImIm- β -HpPyHp- γ -PyHpPy- β -PyPy
	1294 β) 5'-W G G T T A A W-3'	ImIm- β -HpPyPy- γ -HpHpPy- β -PyPy
	1295 β) 5'-W G G T T A G W-3'	ImIm- β -HpPyIm- γ -PyHpPy- β -PyPy
10	1296 β) 5'-W G G T T A C W-3'	ImIm- β -HpPyPy- γ -ImHpPy- β -PyPy
	1297 β) 5'-W G G T T G T W-3'	ImIm- β -HpImHp- γ -PyPyPy- β -PyPy
	1298 β) 5'-W G G T T G A W-3'	ImIm- β -HpImPy- γ -HpPyPy- β -PyPy
	1299 β) 5'-W G G T T G G W-3'	ImIm- β -HpImIm- γ -PyPyPy- β -PyPy
	1300 β) 5'-W G G T T G C W-3'	ImIm- β -HpImPy- γ -ImPyPy- β -PyPy
	1301 β) 5'-W G G T T C T W-3'	ImIm- β -HpPyHp- γ -PyImPy- β -PyPy
	1302 β) 5'-W G G T T C A W-3'	ImIm- β -HpPyPy- γ -HpImPy- β -PyPy
	1303 β) 5'-W G G T T C G W-3'	ImIm- β -HpPyIm- γ -PyImPy- β -PyPy
	1304 β) 5'-W G G T T C C W-3'	ImIm- β -HpPyPy- γ -ImImPy- β -PyPy
	1305 β) 5'-W G G T A T T W-3'	ImIm- β -PyHpHp- γ -PyPyHp- β -PyPy
	1306 β) 5'-W G G T A T A W-3'	ImIm- β -PyHpPy- γ -HpPyHp- β -PyPy
	1307 β) 5'-W G G T A T G W-3'	ImIm- β -PyHpIm- γ -PyPyHp- β -PyPy
	1308 β) 5'-W G G T A T C W-3'	ImIm- β -PyHpPy- γ -ImPyHp- β -PyPy
	1309 β) 5'-W G G T A A T W-3'	ImIm- β -PyPyHp- γ -PyHpHp- β -PyPy
	1310 β) 5'-W G G T A A A W-3'	ImIm- β -PyPyPy- γ -HpHpHp- β -PyPy
25	1311 β) 5'-W G G T A A G W-3'	ImIm- β -PyPyIm- γ -PyHpHp- β -PyPy
	1312 β) 5'-W G G T A A C W-3'	ImIm- β -PyPyPy- γ -ImHpHp- β -PyPy
	1313 β) 5'-W G G T A G T W-3'	ImIm- β -PyImHp- γ -PyPyHp- β -PyPy
	1314 β) 5'-W G G T A G A W-3'	ImIm- β -PyImPy- γ -HpPyHp- β -PyPy
	1315 β) 5'-W G G T A G G W-3'	ImIm- β -PyImIm- γ -PyPyHp- β -PyPy
30	1316 β) 5'-W G G T A G C W-3'	ImIm- β -PyImPy- γ -ImPyHp- β -PyPy
	1317 β) 5'-W G G T A C T W-3'	ImIm- β -PyPyHp- γ -PyImHp- β -PyPy
	1318 β) 5'-W G G T A C A W-3'	ImIm- β -PyPyPy- γ -HpImHp- β -PyPy
	1319 β) 5'-W G G T A C G W-3'	ImIm- β -PyPyIm- γ -PyImHp- β -PyPy
	1320 β) 5'-W G G T A C C W-3'	ImIm- β -PyPyPy- γ -ImImHp- β -PyPy

TABLE 151: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1321 β) 5'-W G G T G T T W-3'	ImIm- β -ImHpHp- γ -PyPyPy- β -PyPy
5	1322 β) 5'-W G G T G T A W-3'	ImIm- β -ImHpPy- γ -HpPyPy- β -PyPy
	1323 β) 5'-W G G T G T G W-3'	ImIm- β -ImHpIm- γ -PyPyPy- β -PyPy
	1324 β) 5'-W G G T G T C W-3'	ImIm- β -ImHpPy- γ -ImPyPy- β -PyPy
	1325 β) 5'-W G G T G A T W-3'	ImIm- β -ImPyHp- γ -PyHpPy- β -PyPy
	1326 β) 5'-W G G T G A A W-3'	ImIm- β -ImPyPy- γ -HpHpPy- β -PyPy
10	1327 β) 5'-W G G T G A G W-3'	ImIm- β -ImPyIm- γ -PyHpPy- β -PyPy
	1328 β) 5'-W G G T G A C W-3'	ImIm- β -ImPyPy- γ -ImHpPy- β -PyPy
	1329 β) 5'-W G G T G G T W-3'	ImIm- β -ImImHp- γ -PyPyPy- β -PyPy
	1330 β) 5'-W G G T G G A W-3'	ImIm- β -ImImPy- γ -HpPyPy- β -PyPy
	1331 β) 5'-W G G T G C T W-3'	ImIm- β -ImPyHp- γ -PyImPy- β -PyPy
	1332 β) 5'-W G G T G C A W-3'	ImIm- β -ImPyPy- γ -HpImPy- β -PyPy
	1333 β) 5'-W G G T G G G W-3'	ImIm- β -ImImIm- γ -PyPyPy- β -PyPy
	1334 β) 5'-W G G T G G C W-3'	ImIm- β -ImImPy- γ -ImPyPy- β -PyPy
	1335 β) 5'-W G G T G C G W-3'	ImIm- β -ImPyIm- γ -PyImPy- β -PyPy
	1336 β) 5'-W G G T G C C W-3'	ImIm- β -ImPyPy- γ -ImImPy- β -PyPy
20	1337 β) 5'-W G G T C T T W-3'	ImIm- β -PyHpHp- γ -PyPyIm- β -PyPy
	1338 β) 5'-W G G T C T A W-3'	ImIm- β -PyHpPy- γ -HpPyIm- β -PyPy
	1339 β) 5'-W G G T C T G W-3'	ImIm- β -PyHpIm- γ -PyPyIm- β -PyPy
	1340 β) 5'-W G G T C T C W-3'	ImIm- β -PyHpPy- γ -ImPyIm- β -PyPy
	1341 β) 5'-W G G T C A T W-3'	ImIm- β -PyPyHp- γ -PyHpIm- β -PyPy
25	1342 β) 5'-W G G T C A A W-3'	ImIm- β -PyPyPy- γ -HpHpIm- β -PyPy
	1343 β) 5'-W G G T C A G W-3'	ImIm- β -PyPyIm- γ -PyHpIm- β -PyPy
	1344 β) 5'-W G G T C A C W-3'	ImIm- β -PyPyPy- γ -ImHpIm- β -PyPy
	1345 β) 5'-W G G T C G T W-3'	ImIm- β -PyImHp- γ -PyPyIm- β -PyPy
	1346 β) 5'-W G G T C G A W-3'	ImIm- β -PyImPy- γ -HpPyIm- β -PyPy
30	1347 β) 5'-W G G T C C T W-3'	ImIm- β -PyPyHp- γ -PyImIm- β -PyPy
	1348 β) 5'-W G G T C C A W-3'	ImIm- β -PyPyPy- γ -HpImIm- β -PyPy
	1349 β) 5'-W G G T C G G W-3'	ImIm- β -PyImIm- γ -PyPyIm- β -PyPy
	1350 β) 5'-W G G T C G C W-3'	ImIm- β -PyImPy- γ -ImPyIm- β -PyPy
	1351 β) 5'-W G G T C C G W-3'	ImIm- β -PyPyIm- γ -PyImIm- β -PyPy
35	1352 β) 5'-W G G T C C C W-3'	ImIm- β -PyPyPy- γ -ImImIm- β -PyPy

TABLE 152: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1353 β) 5'-W G G A T T T W-3'	ImIm- β -HpHpHp- γ -PyPyPy- β -PyPy
5	1354 β) 5'-W G G A T T A W-3'	ImIm- β -HpHpPy- γ -HpPyPy- β -PyPy
	1355 β) 5'-W G G A T T G W-3'	ImIm- β -HpHpIm- γ -PyPyPy- β -PyPy
	1356 β) 5'-W G G A T T C W-3'	ImIm- β -HpHpPy- γ -ImPyPy- β -PyPy
	1357 β) 5'-W G G A T A T W-3'	ImIm- β -HpPyHp- γ -PyHpPy- β -PyPy
	1358 β) 5'-W G G A T A A W-3'	ImIm- β -HpPyPy- γ -HpHpPy- β -PyPy
10	1359 β) 5'-W G G A T A G W-3'	ImIm- β -HpPyIm- γ -PyHpPy- β -PyPy
	1360 β) 5'-W G G A T A C W-3'	ImIm- β -HpPyPy- γ -ImHpPy- β -PyPy
	1361 β) 5'-W G G A T G T W-3'	ImIm- β -HpImHp- γ -PyPyPy- β -PyPy
	1362 β) 5'-W G G A T G A W-3'	ImIm- β -HpImPy- γ -HpPyPy- β -PyPy
	1363 β) 5'-W G G A T G G W-3'	ImIm- β -HpImIm- γ -PyPyPy- β -PyPy
	1364 β) 5'-W G G A T G C W-3'	ImIm- β -HpImPy- γ -ImPyPy- β -PyPy
	1365 β) 5'-W G G A T C T W-3'	ImIm- β -HpPyHp- γ -PyImPy- β -PyPy
	1366 β) 5'-W G G A T C A W-3'	ImIm- β -HpPyPy- γ -HpImPy- β -PyPy
	1367 β) 5'-W G G A T C G W-3'	ImIm- β -HpPyIm- γ -PyImPy- β -PyPy
	1368 β) 5'-W G G A T C C W-3'	ImIm- β -HpPyPy- γ -ImImPy- β -PyPy
20	1369 β) 5'-W G G A A T T W-3'	ImIm- β -PyHpHp- γ -PyPyHp- β -PyPy
	1370 β) 5'-W G G A A T A W-3'	ImIm- β -PyHpPy- γ -HpPyHp- β -PyPy
	1371 β) 5'-W G G A A T G W-3'	ImIm- β -PyHpIm- γ -PyPyHp- β -PyPy
	1372 β) 5'-W G G A A T C W-3'	ImIm- β -PyHpPy- γ -ImPyHp- β -PyPy
	1373 β) 5'-W G G A A A T W-3'	ImIm- β -PyPyHp- γ -PyHpHp- β -PyPy
25	1374 β) 5'-W G G A A A A W-3'	ImIm- β -PyPyPy- γ -HpHpHp- β -PyPy
	1375 β) 5'-W G G A A A G W-3'	ImIm- β -PyPyIm- γ -PyHpHp- β -PyPy
	1376 β) 5'-W G G A A A C W-3'	ImIm- β -PyPyPy- γ -ImHpHp- β -PyPy
	1377 β) 5'-W G G A A G T W-3'	ImIm- β -PyImHp- γ -PyPyHp- β -PyPy
	1378 β) 5'-W G G A A G A W-3'	ImIm- β -PyImPy- γ -HpPyHp- β -PyPy
30	1379 β) 5'-W G G A A G G W-3'	ImIm- β -PyImIm- γ -PyPyHp- β -PyPy
	1380 β) 5'-W G G A A G C W-3'	ImIm- β -PyImPy- γ -ImPyHp- β -PyPy
	1381 β) 5'-W G G A A C T W-3'	ImIm- β -PyPyHp- γ -PyImHp- β -PyPy
	1382 β) 5'-W G G A A C A W-3'	ImIm- β -PyPyPy- γ -HpImHp- β -PyPy
	1383 β) 5'-W G G A A C G W-3'	ImIm- β -PyPyIm- γ -PyImHp- β -PyPy
35	1384 β) 5'-W G G A A C C W-3'	ImIm- β -PyPyPy- γ -ImImHp- β -PyPy

TABLE 153: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1385 β) 5'-W G G A G T T W-3'	ImIm- β -ImHpHp- γ -PyPyPy- β -PyPy
5	1386 β) 5'-W G G A G T A W-3'	ImIm- β -ImHpPy- γ -HpPyPy- β -PyPy
	1387 β) 5'-W G G A G T G W-3'	ImIm- β -ImHpIm- γ -PyPyPy- β -PyPy
	1388 β) 5'-W G G A G T C W-3'	ImIm- β -ImHpPy- γ -ImPyPy- β -PyPy
	1389 β) 5'-W G G A G A T W-3'	ImIm- β -ImPyHp- γ -PyHpPy- β -PyPy
	1390 β) 5'-W G G A G A A W-3'	ImIm- β -ImPyPy- γ -HpHpPy- β -PyPy
10	1391 β) 5'-W G G A G A G W-3'	ImIm- β -ImPyIm- γ -PyHpPy- β -PyPy
	1392 β) 5'-W G G A G A C W-3'	ImIm- β -ImPyPy- γ -ImHpPy- β -PyPy
	1393 β) 5'-W G G A G G T W-3'	ImIm- β -ImImHp- γ -PyPyPy- β -PyPy
	1394 β) 5'-W G G A G G A W-3'	ImIm- β -ImImPy- γ -HpPyPy- β -PyPy
	1395 β) 5'-W G G A G C T W-3'	ImIm- β -ImPyHp- γ -PyImPy- β -PyPy
15	1396 β) 5'-W G G A G C A W-3'	ImIm- β -ImPyPy- γ -HpImPy- β -PyPy
	1397 β) 5'-W G G A G G G W-3'	ImIm- β -ImImIm- γ -PyPyPy- β -PyPy
	1398 β) 5'-W G G A G G C W-3'	ImIm- β -ImImPy- γ -ImPyPy- β -PyPy
	1399 β) 5'-W G G A G C G W-3'	ImIm- β -ImPyIm- γ -PyImPy- β -PyPy
	1400 β) 5'-W G G A G C C W-3'	ImIm- β -ImPyPy- γ -ImImPy- β -PyPy
20	1401 β) 5'-W G G A C T T W-3'	ImIm- β -PyHpHp- γ -PyPyIm- β -PyPy
	1402 β) 5'-W G G A C T A W-3'	ImIm- β -PyHpPy- γ -HpPyIm- β -PyPy
	1403 β) 5'-W G G A C T G W-3'	ImIm- β -PyHpIm- γ -PyPyIm- β -PyPy
	1404 β) 5'-W G G A C T C W-3'	ImIm- β -PyHpPy- γ -ImPyIm- β -PyPy
	1405 β) 5'-W G G A C A T W-3'	ImIm- β -PyPyHp- γ -PyHpIm- β -PyPy
25	1406 β) 5'-W G G A C A A W-3'	ImIm- β -PyPyPy- γ -HpHpIm- β -PyPy
	1407 β) 5'-W G G A C A G W-3'	ImIm- β -PyPyIm- γ -PyHpIm- β -PyPy
	1408 β) 5'-W G G A C A C W-3'	ImIm- β -PyPyPy- γ -ImHpIm- β -PyPy
	1409 β) 5'-W G G A C G T W-3'	ImIm- β -PyImHp- γ -PyPyIm- β -PyPy
	1410 β) 5'-W G G A C G A W-3'	ImIm- β -PyImPy- γ -HpPyIm- β -PyPy
30	1411 β) 5'-W G G A C C T W-3'	ImIm- β -PyPyHp- γ -PyImIm- β -PyPy
	1412 β) 5'-W G G A C C A W-3'	ImIm- β -PyPyPy- γ -HpImIm- β -PyPy
	1413 β) 5'-W G G A C G G W-3'	ImIm- β -PyImIm- γ -PyPyIm- β -PyPy
	1414 β) 5'-W G G A C G C W-3'	ImIm- β -PyImPy- γ -ImPyIm- β -PyPy
	1415 β) 5'-W G G A C C G W-3'	ImIm- β -PyPyIm- γ -PyImIm- β -PyPy
35	1416 β) 5'-W G G A C C C W-3'	ImIm- β -PyPyPy- γ -ImImIm- β -PyPy

TABLE 154: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1417 β) 5'-W G G C T T T W-3'	ImImPy- β -HpHp- γ -PyPy- β -ImPyPy
5	1418 β) 5'-W G G C T T A W-3'	ImImPy- β -HpPy- γ -HpPy- β -ImPyPy
	1419 β) 5'-W G G C T T G W-3'	ImImPy- β -HpIm- γ -PyPy- β -ImPyPy
	1420 β) 5'-W G G C T T C W-3'	ImImPy- β -HpPy- γ -ImPy- β -ImPyPy
	1421 β) 5'-W G G C T A T W-3'	ImImPy- β -PyHp- γ -PyHp- β -ImPyPy
	1422 β) 5'-W G G C T A A W-3'	ImImPy- β -PyPy- γ -HpHp- β -ImPyPy
10	1423 β) 5'-W G G C T A G W-3'	ImImPy- β -PyIm- γ -PyHp- β -ImPyPy
	1424 β) 5'-W G G C T A C W-3'	ImImPy- β -PyPy- γ -ImHp- β -ImPyPy
	1425 β) 5'-W G G C T G T W-3'	ImImPy- β -ImHp- γ -PyPy- β -ImPyPy
	1426 β) 5'-W G G C T G A W-3'	ImImPy- β -ImPy- γ -HpPy- β -ImPyPy
	1427 β) 5'-W G G C T G G W-3'	ImImPy- β -ImIm- γ -PyPy- β -ImPyPy
	1428 β) 5'-W G G C T G C W-3'	ImImPy- β -ImPy- γ -ImPy- β -ImPyPy
	1429 β) 5'-W G G C T C T W-3'	ImImPy- β -PyHp- γ -PyIm- β -ImPyPy
	1430 β) 5'-W G G C T C A W-3'	ImImPy- β -PyPy- γ -HpIm- β -ImPyPy
	1431 β) 5'-W G G C T C G W-3'	ImImPy- β -PyIm- γ -PyIm- β -ImPyPy
	1432 β) 5'-W G G C T C C W-3'	ImImPy- β -PyPy- γ -ImIm- β -ImPyPy
20	1433 β) 5'-W G G C A T T W-3'	ImImPy- β -HpHp- γ -PyPy- β -ImPyPy
	1434 β) 5'-W G G C A T A W-3'	ImImPy- β -HpPy- γ -HpPy- β -ImPyPy
	1435 β) 5'-W G G C A T G W-3'	ImImPy- β -HpIm- γ -PyPy- β -ImPyPy
	1436 β) 5'-W G G C A T C W-3'	ImImPy- β -HpPy- γ -ImPy- β -ImPyPy
	1437 β) 5'-W G G C A A T W-3'	ImImPy- β -PyHp- γ -PyHp- β -ImPyPy
25	1438 β) 5'-W G G C A A A W-3'	ImImPy- β -PyPy- γ -HpHp- β -ImPyPy
	1439 β) 5'-W G G C A A G W-3'	ImImPy- β -PyIm- γ -PyHp- β -ImPyPy
	1440 β) 5'-W G G C A A C W-3'	ImImPy- β -PyPy- γ -ImHp- β -ImPyPy
	1441 β) 5'-W G G C A G T W-3'	ImImPy- β -ImHp- γ -PyPy- β -ImPyPy
	1442 β) 5'-W G G C A G A W-3'	ImImPy- β -ImPy- γ -HpPy- β -ImPyPy
30	1443 β) 5'-W G G C A G G W-3'	ImImPy- β -ImIm- γ -PyPy- β -ImPyPy
	1444 β) 5'-W G G C A G C W-3'	ImImPy- β -ImPy- γ -ImPy- β -ImPyPy
	1445 β) 5'-W G G C A C T W-3'	ImImPy- β -PyHp- γ -PyIm- β -ImPyPy
	1446 β) 5'-W G G C A C A W-3'	ImImPy- β -PyPy- γ -HpIm- β -ImPyPy
	1447 β) 5'-W G G C A C G W-3'	ImImPy- β -PyIm- γ -PyIm- β -ImPyPy
35	1448 β) 5'-W G G C A C C W-3'	ImImPy- β -PyPy- γ -ImIm- β -ImPyPy

TABLE 155: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGGCSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1449 β) 5'-W G G C G T T W-3'	ImIm- β -ImHpHp- γ -PyPy- β -ImPyPy
5	1450 β) 5'-W G G C G T A W-3'	ImIm- β -ImHpPy- γ -HpPy- β -ImPyPy
	1451 β) 5'-W G G C G T G W-3'	ImIm- β -ImHpIm- γ -PyPy- β -ImPyPy
	1452 β) 5'-W G G C G T C W-3'	ImIm- β -ImHpPy- γ -ImPy- β -ImPyPy
	1453 β) 5'-W G G C G A T W-3'	ImIm- β -ImPyHp- γ -PyHp- β -ImPyPy
	1454 β) 5'-W G G C G A A W-3'	ImIm- β -ImPyPy- γ -HpHp- β -ImPyPy
10	1455 β) 5'-W G G C G A G W-3'	ImIm- β -ImPyIm- γ -PyHp- β -ImPyPy
	1456 β) 5'-W G G C G A C W-3'	ImIm- β -ImPyPy- γ -ImHp- β -ImPyPy
	1457 β) 5'-W G G C G G T W-3'	ImIm- β -ImImHp- γ -PyPy- β -ImPyPy
	1458 β) 5'-W G G C G G A W-3'	ImIm- β -ImImPy- γ -HpPy- β -ImPyPy
	1459 β) 5'-W G G C G C T W-3'	ImIm- β -ImPyHp- γ -PyIm- β -ImPyPy
15	1460 β) 5'-W G G C G C A W-3'	ImIm- β -ImPyPy- γ -HpIm- β -ImPyPy
	1461 β) 5'-W G G C C T T W-3'	ImIm- β -PyHpHp- γ -Py- β -ImImPyPy
	1462 β) 5'-W G G C C T A W-3'	ImIm- β -PyHpPy- γ -Hp- β -ImImPyPy
	1463 β) 5'-W G G C C T G W-3'	ImIm- β -PyHpIm- γ -Py- β -ImImPyPy
	1464 β) 5'-W G G C C T C W-3'	ImIm- β -PyHpPy- γ -Im- β -ImImPyPy
20	1465 β) 5'-W G G C C A T W-3'	ImIm- β -PyPyHp- γ -Py- β -ImImPyPy
	1466 β) 5'-W G G C C A A W-3'	ImIm- β -PyPyPy- γ -Hp- β -ImImPyPy
	1467 β) 5'-W G G C C A G W-3'	ImIm- β -PyPyIm- γ -Py- β -ImImPyPy
	1468 β) 5'-W G G C C A C W-3'	ImIm- β -PyPyPy- γ -Im- β -ImImPyPy
	1469 β) 5'-W G G C C G T W-3'	ImIm- β -PyImHp- γ -Py- β -ImImPyPy
25	1470 β) 5'-W G G C C G A W-3'	ImIm- β -PyImPy- γ -Hp- β -ImImPyPy
	1471 β) 5'-W G G C C C T W-3'	ImIm- β -PyPyHp- γ -PyImImIm- β -Py
	1472 β) 5'-W G G C C C A W-3'	ImIm- β -PyPyPy- γ -HpImImIm- β -Py
	G57 β) 5'-W G G C G G G W-3'	ImIm- β -ImImIm- γ -PyPy- β -ImPyPy
	G58 β) 5'-W G G C G G C W-3'	ImIm- β -ImImPy- γ -ImPy- β -ImPyPy
30	G59 β) 5'-W G G C G C G W-3'	ImIm- β -ImPyIm- γ -PyIm- β -ImPyPy
	G60 β) 5'-W G G C G C C W-3'	ImIm- β -ImPyPy- γ -ImIm- β -ImPyPy
	G61 β) 5'-W G G C C G G W-3'	ImIm- β -PyImIm- γ -Py- β -ImImPyPy
	G62 β) 5'-W G G C C G C W-3'	ImIm- β -PyImPy- γ -Im- β -ImImPyPy
	G63 β) 5'-W G G C C C G W-3'	ImIm- β -PyPyIm- γ -PyImImIm- β -Py
35	G64 β) 5'-W G G C C C C W-3'	ImIm- β -PyPyPy- γ -ImImImIm- β -Py

TABLE 156: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1473 β) 5'-W G C G T T T W-3'	ImPyIm- β -HpHp- γ -PyPyPy- β -ImPy
5	1474 β) 5'-W G C G T T A W-3'	ImPyIm- β -HpPy- γ -HpPyPy- β -ImPy
	1475 β) 5'-W G C G T T G W-3'	ImPyIm- β -HpIm- γ -PyPyPy- β -ImPy
	1476 β) 5'-W G C G T T C W-3'	ImPyIm- β -HpPy- γ -ImPyPy- β -ImPy
	1477 β) 5'-W G C G T A T W-3'	ImPyIm- β -PyHp- γ -PyHpPy- β -ImPy
	1478 β) 5'-W G C G T A A W-3'	ImPyIm- β -PyPy- γ -HpHpPy- β -ImPy
10	1479 β) 5'-W G C G T A G W-3'	ImPyIm- β -PyIm- γ -PyHpPy- β -ImPy
	1480 β) 5'-W G C G T A C W-3'	ImPyIm- β -PyPy- γ -ImHpPy- β -ImPy
	1481 β) 5'-W G C G T G T W-3'	ImPyIm- β -ImHp- γ -PyPyPy- β -ImPy
	1482 β) 5'-W G C G T G A W-3'	ImPyIm- β -ImPy- γ -HpPyPy- β -ImPy
	1483 β) 5'-W G C G T G G W-3'	ImPyIm- β -ImIm- γ -PyPyPy- β -ImPy
15	1484 β) 5'-W G C G T G C W-3'	ImPyIm- β -ImPy- γ -ImPyPy- β -ImPy
	1485 β) 5'-W G C G T C T W-3'	ImPyIm- β -PyHp- γ -PyImPy- β -ImPy
	1486 β) 5'-W G C G T C A W-3'	ImPyIm- β -PyPy- γ -HpImPy- β -ImPy
	1487 β) 5'-W G C G T C G W-3'	ImPyIm- β -PyIm- γ -PyImPy- β -ImPy
	1488 β) 5'-W G C G T C C W-3'	ImPyIm- β -PyPy- γ -ImImPy- β -ImPy
20	1489 β) 5'-W G C G A T T W-3'	ImPyIm- β -HpHp- γ -PyPyHp- β -ImPy
	1490 β) 5'-W G C G A T A W-3'	ImPyIm- β -HpPy- γ -HpPyHp- β -ImPy
	1491 β) 5'-W G C G A T G W-3'	ImPyIm- β -HpIm- γ -PyPyHp- β -ImPy
	1492 β) 5'-W G C G A T C W-3'	ImPyIm- β -HpPy- γ -ImPyHp- β -ImPy
	1493 β) 5'-W G C G A A T W-3'	ImPyIm- β -PyHp- γ -PyHpHp- β -ImPy
25	1494 β) 5'-W G C G A A A W-3'	ImPyIm- β -PyPy- γ -HpHpHp- β -ImPy
	1495 β) 5'-W G C G A A G W-3'	ImPyIm- β -PyIm- γ -PyHpHp- β -ImPy
	1496 β) 5'-W G C G A A C W-3'	ImPyIm- β -PyPy- γ -ImHpHp- β -ImPy
	1497 β) 5'-W G C G A G T W-3'	ImPyIm- β -ImHp- γ -PyPyHp- β -ImPy
	1498 β) 5'-W G C G A G A W-3'	ImPyIm- β -ImPy- γ -HpPyHp- β -ImPy
30	1499 β) 5'-W G C G A G G W-3'	ImPyIm- β -ImIm- γ -PyPyHp- β -ImPy
	1490 β) 5'-W G C G A G C W-3'	ImPyIm- β -ImPy- γ -ImPyHp- β -ImPy
	1501 β) 5'-W G C G A C T W-3'	ImPyIm- β -PyHp- γ -PyImHp- β -ImPy
	1502 β) 5'-W G C G A C A W-3'	ImPyIm- β -PyPy- γ -HpImHp- β -ImPy
	1503 β) 5'-W G C G A C G W-3'	ImPyIm- β -PyIm- γ -PyImHp- β -ImPy
35	1504 β) 5'-W G C G A C C W-3'	ImPyIm- β -PyPy- γ -ImImHp- β -ImPy

TABLE 157: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1505 β) 5'-W G C G G T T W-3'	Im- β -ImImHpHp- γ -PyPyPy- β -ImPy
5	1506 β) 5'-W G C G G T A W-3'	Im- β -ImImHpPy- γ -HpPyPy- β -ImPy
	1507 β) 5'-W G C G G T G W-3'	Im- β -ImImHpIm- γ -PyPyPy- β -ImPy
	1508 β) 5'-W G C G G T C W-3'	Im- β -ImImHpPy- γ -ImPyPy- β -ImPy
	1509 β) 5'-W G C G G A T W-3'	Im- β -ImImPyHp- γ -PyHpPy- β -ImPy
	1510 β) 5'-W G C G G A A W-3'	Im- β -ImImPyPy- γ -HpHpPy- β -ImPy
10	1511 β) 5'-W G C G G A G W-3'	Im- β -ImImPyIm- γ -PyHpPy- β -ImPy
	1512 β) 5'-W G C G G A C W-3'	Im- β -ImImPyPy- γ -ImHpPy- β -ImPy
	1513 β) 5'-W G C G G G T W-3'	Im- β -ImImImHp- γ -PyPyPy- β -ImPy
	1514 β) 5'-W G C G G G A W-3'	Im- β -ImImImPy- γ -HpPyPy- β -ImPy
	1515 β) 5'-W G C G G C T W-3'	Im- β -ImImPyHp- γ -PyImPy- β -ImPy
	1516 β) 5'-W G C G G C A W-3'	Im- β -ImImPyPy- γ -HpImPy- β -ImPy
	1517 β) 5'-W G C G C T T W-3'	ImPyIm- β -HpHp- γ -PyPyIm- β -ImPy
	1518 β) 5'-W G C G C T A W-3'	ImPyIm- β -HpPy- γ -HpPyIm- β -ImPy
	1519 β) 5'-W G C G C T G W-3'	ImPyIm- β -HpIm- γ -PyPyIm- β -ImPy
	1520 β) 5'-W G C G C T C W-3'	ImPyIm- β -HpPy- γ -ImPyIm- β -ImPy
20	1521 β) 5'-W G C G C A T W-3'	ImPyIm- β -PyHp- γ -PyHpIm- β -ImPy
	1522 β) 5'-W G C G C A A W-3'	ImPyIm- β -PyPy- γ -HpHpIm- β -ImPy
	1523 β) 5'-W G C G C A G W-3'	ImPyIm- β -PyIm- γ -PyHpIm- β -ImPy
	1524 β) 5'-W G C G C A C W-3'	ImPyIm- β -PyPy- γ -ImHpIm- β -ImPy
	1525 β) 5'-W G C G C G T W-3'	ImPyIm- β -ImHp- γ -PyPyIm- β -ImPy
25	1526 β) 5'-W G C G C G A W-3'	ImPyIm- β -ImPy- γ -HpPyIm- β -ImPy
	1527 β) 5'-W G C G C C T W-3'	ImPyIm- β -PyHp- γ -PyImIm- β -ImPy
	1528 β) 5'-W G C G C C A W-3'	ImPyIm- β -PyPy- γ -HpImIm- β -ImPy
	G65 β) 5'-W G C G G G G W-3'	Im- β -ImImImIm- γ -PyPyPy- β -ImPy
	G66 β) 5'-W G C G G G C W-3'	Im- β -ImImImPy- γ -ImPyPy- β -ImPy
30	G67 β) 5'-W G C G G C G W-3'	Im- β -ImImPyIm- γ -PyImPy- β -ImPy
	G68 β) 5'-W G C G G C C W-3'	Im- β -ImImPyPy- γ -ImImPy- β -ImPy
	G69 β) 5'-W G C G C G G W-3'	ImPyIm- β -ImIm- γ -PyPyIm- β -ImPy
	G70 β) 5'-W G C G C G C W-3'	ImPyIm- β -ImPy- γ -ImPyIm- β -ImPy
	G71 β) 5'-W G C G C C G W-3'	ImPyIm- β -PyIm- γ -PyImIm- β -ImPy
35	G72 β) 5'-W G C G C C C W-3'	ImPyIm- β -PyPy- γ -ImImIm- β -ImPy

TABLE 158: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1529 β) 5'-W G C T T T T W-3'	ImPy- β -HpHpHp- γ -PyPyPy- β -ImPy
5	1530 β) 5'-W G C T T T A W-3'	ImPy- β -HpHpPy- γ -HpPyPy- β -ImPy
	1531 β) 5'-W G C T T T G W-3'	ImPy- β -HpHpIm- γ -PyPyPy- β -ImPy
	1532 β) 5'-W G C T T T C W-3'	ImPy- β -HpHpPy- γ -ImPyPy- β -ImPy
	1533 β) 5'-W G C T T A T W-3'	ImPy- β -HpPyHp- γ -PyHpPy- β -ImPy
	1534 β) 5'-W G C T T A A W-3'	ImPy- β -HpPyPy- γ -HpHpPy- β -ImPy
10	1535 β) 5'-W G C T T A G W-3'	ImPy- β -HpPyIm- γ -PyHpPy- β -ImPy
	1536 β) 5'-W G C T T A C W-3'	ImPy- β -HpPyPy- γ -ImHpPy- β -ImPy
	1537 β) 5'-W G C T T G T W-3'	ImPy- β -HpImHp- γ -PyPyPy- β -ImPy
	1538 β) 5'-W G C T T G A W-3'	ImPy- β -HpImPy- γ -HpPyPy- β -ImPy
	1539 β) 5'-W G C T T G G W-3'	ImPy- β -HpImIm- γ -PyPyPy- β -ImPy
15	1540 β) 5'-W G C T T G C W-3'	ImPy- β -HpImPy- γ -ImPyPy- β -ImPy
	1541 β) 5'-W G C T T C T W-3'	ImPy- β -HpPyHp- γ -PyImPy- β -ImPy
	1542 β) 5'-W G C T T C A W-3'	ImPy- β -HpPyPy- γ -HpImPy- β -ImPy
	1543 β) 5'-W G C T T C G W-3'	ImPy- β -HpPyIm- γ -PyImPy- β -ImPy
	1544 β) 5'-W G C T T C C W-3'	ImPy- β -HpPyPy- γ -ImImPy- β -ImPy
20	1545 β) 5'-W G C T A T T W-3'	ImPy- β -PyHpHp- γ -PyPyHp- β -ImPy
	1546 β) 5'-W G C T A T A W-3'	ImPy- β -PyHpPy- γ -HpPyHp- β -ImPy
	1547 β) 5'-W G C T A T G W-3'	ImPy- β -PyHpIm- γ -PyPyHp- β -ImPy
	1548 β) 5'-W G C T A T C W-3'	ImPy- β -PyHpPy- γ -ImPyHp- β -ImPy
	1549 β) 5'-W G C T A A T W-3'	ImPy- β -PyPyHp- γ -PyHpHp- β -ImPy
25	1550 β) 5'-W G C T A A A W-3'	ImPy- β -PyPyPy- γ -HpHpHp- β -ImPy
	1551 β) 5'-W G C T A A G W-3'	ImPy- β -PyPyIm- γ -PyHpHp- β -ImPy
	1552 β) 5'-W G C T A A C W-3'	ImPy- β -PyPyPy- γ -ImHpHp- β -ImPy
	1553 β) 5'-W G C T A G T W-3'	ImPy- β -PyImHp- γ -PyPyHp- β -ImPy
	1554 β) 5'-W G C T A G A W-3'	ImPy- β -PyImPy- γ -HpPyHp- β -ImPy
30	1555 β) 5'-W G C T A G G W-3'	ImPy- β -PyImIm- γ -PyPyHp- β -ImPy
	1556 β) 5'-W G C T A G C W-3'	ImPy- β -PyImPy- γ -ImPyHp- β -ImPy
	1557 β) 5'-W G C T A C T W-3'	ImPy- β -PyPyHp- γ -PyImHp- β -ImPy
	1558 β) 5'-W G C T A C A W-3'	ImPy- β -PyPyPy- γ -HpImHp- β -ImPy
	1559 β) 5'-W G C T A C G W-3'	ImPy- β -PyPyIm- γ -PyImHp- β -ImPy
35	1560 β) 5'-W G C T A C C W-3'	ImPy- β -PyPyPy- γ -ImImHp- β -ImPy

TABLE 159: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCTSNW-3'

	DNA sequence	aromatic amino acid sequence
	1561 β) 5'-W G C T G T T W-3'	ImPy- β -ImHpHp- γ -PyPyPy- β -ImPy
5	1562 β) 5'-W G C T G T A W-3'	ImPy- β -ImHpPy- γ -HpPyPy- β -ImPy
	1563 β) 5'-W G C T G T G W-3'	ImPy- β -ImHpIm- γ -PyPyPy- β -ImPy
	1564 β) 5'-W G C T G T C W-3'	ImPy- β -ImHpPy- γ -ImPyPy- β -ImPy
	1565 β) 5'-W G C T G A T W-3'	ImPy- β -ImPyHp- γ -PyHpPy- β -ImPy
	1566 β) 5'-W G C T G A A W-3'	ImPy- β -ImPyPy- γ -HpHpPy- β -ImPy
10	1567 β) 5'-W G C T G A G W-3'	ImPy- β -ImPyIm- γ -PyHpPy- β -ImPy
	1568 β) 5'-W G C T G A C W-3'	ImPy- β -ImPyPy- γ -ImHpPy- β -ImPy
	1569 β) 5'-W G C T G G T W-3'	ImPy- β -ImImHp- γ -PyPyPy- β -ImPy
	1570 β) 5'-W G C T G G A W-3'	ImPy- β -ImImPy- γ -HpPyPy- β -ImPy
	1571 β) 5'-W G C T G C T W-3'	ImPy- β -ImPyHp- γ -PyImPy- β -ImPy
	1572 β) 5'-W G C T G C A W-3'	ImPy- β -ImPyPy- γ -HpImPy- β -ImPy
	1573 β) 5'-W G C T G G G W-3'	ImPy- β -ImImIm- γ -PyPyPy- β -ImPy
	1574 β) 5'-W G C T G G C W-3'	ImPy- β -ImImPy- γ -ImPyPy- β -ImPy
	1575 β) 5'-W G C T G C G W-3'	ImPy- β -ImPyIm- γ -PyImPy- β -ImPy
	1576 β) 5'-W G C T G C C W-3'	ImPy- β -ImPyPy- γ -ImImPy- β -ImPy
	1577 β) 5'-W G C T C T T W-3'	ImPy- β -PyHpHp- γ -PyPyIm- β -ImPy
	1578 β) 5'-W G C T C T A W-3'	ImPy- β -PyHpPy- γ -HpPyIm- β -ImPy
	1579 β) 5'-W G C T C T G W-3'	ImPy- β -PyHpIm- γ -PyPyIm- β -ImPy
	1580 β) 5'-W G C T C T C W-3'	ImPy- β -PyHpPy- γ -ImPyIm- β -ImPy
	1581 β) 5'-W G C T C A T W-3'	ImPy- β -PyPyHp- γ -PyHpIm- β -ImPy
25	1582 β) 5'-W G C T C A A W-3'	ImPy- β -PyPyPy- γ -HpHpIm- β -ImPy
	1583 β) 5'-W G C T C A G W-3'	ImPy- β -PyPyIm- γ -PyHpIm- β -ImPy
	1584 β) 5'-W G C T C A C W-3'	ImPy- β -PyPyPy- γ -ImHpIm- β -ImPy
	1585 β) 5'-W G C T C G T W-3'	ImPy- β -PyImHp- γ -PyPyIm- β -ImPy
	1586 β) 5'-W G C T C G A W-3'	ImPy- β -PyImPy- γ -HpPyIm- β -ImPy
30	1587 β) 5'-W G C T C C T W-3'	ImPy- β -PyPyHp- γ -PyImIm- β -ImPy
	1588 β) 5'-W G C T C C A W-3'	ImPy- β -PyPyPy- γ -HpImIm- β -ImPy
	1589 β) 5'-W G C T C G G W-3'	ImPy- β -PyImIm- γ -PyPyIm- β -ImPy
	1590 β) 5'-W G C T C G C W-3'	ImPy- β -PyImPy- γ -ImPyIm- β -ImPy
	1591 β) 5'-W G C T C C G W-3'	ImPy- β -PyPyIm- γ -PyImIm- β -ImPy
35	1592 β) 5'-W G C T C C C W-3'	ImPy- β -PyPyPy- γ -ImImIm- β -ImPy

TABLE 160: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1593 β) 5'-W G C A T T T W-3'	ImPy- β -HpHpHp- γ -PyPyPy- β -ImPy
5	1594 β) 5'-W G C A T T A W-3'	ImPy- β -HpHpPy- γ -HpPyPy- β -ImPy
	1595 β) 5'-W G C A T T G W-3'	ImPy- β -HpHpIm- γ -PyPyPy- β -ImPy
	1596 β) 5'-W G C A T T C W-3'	ImPy- β -HpHpPy- γ -ImPyPy- β -ImPy
	1597 β) 5'-W G C A T A T W-3'	ImPy- β -HpPyHp- γ -PyHpPy- β -ImPy
	1598 β) 5'-W G C A T A A W-3'	ImPy- β -HpPyPy- γ -HpHpPy- β -ImPy
10	1599 β) 5'-W G C A T A G W-3'	ImPy- β -HpPyIm- γ -PyHpPy- β -ImPy
	1600 β) 5'-W G C A T A C W-3'	ImPy- β -HpPyPy- γ -ImHpPy- β -ImPy
	1601 β) 5'-W G C A T G T W-3'	ImPy- β -HpImHp- γ -PyPyPy- β -ImPy
	1602 β) 5'-W G C A T G A W-3'	ImPy- β -HpImPy- γ -HpPyPy- β -ImPy
	1603 β) 5'-W G C A T G G W-3'	ImPy- β -HpImIm- γ -PyPyPy- β -ImPy
	1604 β) 5'-W G C A T G C W-3'	ImPy- β -HpImPy- γ -ImPyPy- β -ImPy
	1605 β) 5'-W G C A T C T W-3'	ImPy- β -HpPyHp- γ -PyImPy- β -ImPy
	1606 β) 5'-W G C A T C A W-3'	ImPy- β -HpPyPy- γ -HpImPy- β -ImPy
	1607 β) 5'-W G C A T C G W-3'	ImPy- β -HpPyIm- γ -PyImPy- β -ImPy
	1608 β) 5'-W G C A T C C W-3'	ImPy- β -HpPyPy- γ -ImImPy- β -ImPy
20	1609 β) 5'-W G C A A T T W-3'	ImPy- β -PyHpHp- γ -PyPyHp- β -ImPy
	1610 β) 5'-W G C A A T A W-3'	ImPy- β -PyHpPy- γ -HpPyHp- β -ImPy
	1611 β) 5'-W G C A A T G W-3'	ImPy- β -PyHpIm- γ -PyPyHp- β -ImPy
	1612 β) 5'-W G C A A T C W-3'	ImPy- β -PyHpPy- γ -ImPyHp- β -ImPy
	1613 β) 5'-W G C A A A T W-3'	ImPy- β -PyPyHp- γ -PyHpHp- β -ImPy
25	1614 β) 5'-W G C A A A A W-3'	ImPy- β -PyPyPy- γ -HpHpHp- β -ImPy
	1615 β) 5'-W G C A A A G W-3'	ImPy- β -PyPyIm- γ -PyHpHp- β -ImPy
	1616 β) 5'-W G C A A A C W-3'	ImPy- β -PyPyPy- γ -ImHpHp- β -ImPy
	1617 β) 5'-W G C A A A G T W-3'	ImPy- β -PyImHp- γ -PyPyHp- β -ImPy
	1618 β) 5'-W G C A A A G A W-3'	ImPy- β -PyImPy- γ -HpPyHp- β -ImPy
30	1619 β) 5'-W G C A A A G G W-3'	ImPy- β -PyImIm- γ -PyPyHp- β -ImPy
	1620 β) 5'-W G C A A A G C W-3'	ImPy- β -PyImPy- γ -ImPyHp- β -ImPy
	1621 β) 5'-W G C A A A C T W-3'	ImPy- β -PyPyHp- γ -PyImHp- β -ImPy
	1622 β) 5'-W G C A A A C A W-3'	ImPy- β -PyPyPy- γ -HpImHp- β -ImPy
	1623 β) 5'-W G C A A A C G W-3'	ImPy- β -PyPyIm- γ -PyImHp- β -ImPy
35	1624 β) 5'-W G C A A A C C W-3'	ImPy- β -PyPyPy- γ -ImImHp- β -ImPy

TABLE 161: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1625 β) 5'-W G C A G T T W-3'	ImPy- β -ImHpHp- γ -PyPyPy- β -ImPy
5	1626 β) 5'-W G C A G T A W-3'	ImPy- β -ImHpPy- γ -HpPyPy- β -ImPy
	1627 β) 5'-W G C A G T G W-3'	ImPy- β -ImHpIm- γ -PyPyPy- β -ImPy
	1628 β) 5'-W G C A G T C W-3'	ImPy- β -ImHpPy- γ -ImPyPy- β -ImPy
	1629 β) 5'-W G C A G A T W-3'	ImPy- β -ImPyHp- γ -PyHpPy- β -ImPy
	1630 β) 5'-W G C A G A A W-3'	ImPy- β -ImPyPy- γ -HpHpPy- β -ImPy
10	1631 β) 5'-W G C A G A G W-3'	ImPy- β -ImPyIm- γ -PyHpPy- β -ImPy
	1632 β) 5'-W G C A G A C W-3'	ImPy- β -ImPyPy- γ -ImHpPy- β -ImPy
	1633 β) 5'-W G C A G G T W-3'	ImPy- β -ImImHp- γ -PyPyPy- β -ImPy
	1634 β) 5'-W G C A G G A W-3'	ImPy- β -ImImPy- γ -HpPyPy- β -ImPy
	1635 β) 5'-W G C A G C T W-3'	ImPy- β -ImPyHp- γ -PyImPy- β -ImPy
	1636 β) 5'-W G C A G C A W-3'	ImPy- β -ImPyPy- γ -HpImPy- β -ImPy
	1637 β) 5'-W G C A G G G W-3'	ImPy- β -ImImIm- γ -PyPyPy- β -ImPy
	1638 β) 5'-W G C A G G C W-3'	ImPy- β -ImImPy- γ -ImPyPy- β -ImPy
	1639 β) 5'-W G C A G C G W-3'	ImPy- β -ImPyIm- γ -PyImPy- β -ImPy
	1640 β) 5'-W G C A G C C W-3'	ImPy- β -ImPyPy- γ -ImImPy- β -ImPy
20	1641 β) 5'-W G C A C T T W-3'	ImPy- β -PyHpHp- γ -PyPyIm- β -ImPy
	1642 β) 5'-W G C A C T A W-3'	ImPy- β -PyHpPy- γ -HpPyIm- β -ImPy
	1643 β) 5'-W G C A C T G W-3'	ImPy- β -PyHpIm- γ -PyPyIm- β -ImPy
	1644 β) 5'-W G C A C T C W-3'	ImPy- β -PyHpPy- γ -ImPyIm- β -ImPy
	1645 β) 5'-W G C A C A T W-3'	ImPy- β -PyPyHp- γ -PyHpIm- β -ImPy
25	1646 β) 5'-W G C A C A A W-3'	ImPy- β -PyPyPy- γ -HpHpIm- β -ImPy
	1647 β) 5'-W G C A C A G W-3'	ImPy- β -PyPyIm- γ -PyHpIm- β -ImPy
	1648 β) 5'-W G C A C A C W-3'	ImPy- β -PyPyPy- γ -ImHpIm- β -ImPy
	1649 β) 5'-W G C A C G T W-3'	ImPy- β -PyImHp- γ -PyPyIm- β -ImPy
	1650 β) 5'-W G C A C G A W-3'	ImPy- β -PyImPy- γ -HpPyIm- β -ImPy
30	1651 β) 5'-W G C A C C T W-3'	ImPy- β -PyPyHp- γ -PyImIm- β -ImPy
	1652 β) 5'-W G C A C C A W-3'	ImPy- β -PyPyPy- γ -HpImIm- β -ImPy
	1653 β) 5'-W G C A C G G W-3'	ImPy- β -PyImIm- γ -PyPyIm- β -ImPy
	1654 β) 5'-W G C A C G C W-3'	ImPy- β -PyImPy- γ -ImPyIm- β -ImPy
	1655 β) 5'-W G C A C C G W-3'	ImPy- β -PyPyIm- γ -PyImIm- β -ImPy
35	1656 β) 5'-W G C A C C C W-3'	ImPy- β -PyPyPy- γ -ImImIm- β -ImPy

TABLE 162: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCCWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1657 β) 5'-W G C C T T T W-3'	ImPyPy- β -HpHp- γ -PyPy- β -ImImPy
5	1658 β) 5'-W G C C T T A W-3'	ImPyPy- β -HpPy- γ -HpPy- β -ImImPy
	1659 β) 5'-W G C C T T G W-3'	ImPyPy- β -HpIm- γ -PyPy- β -ImImPy
	1660 β) 5'-W G C C T T C W-3'	ImPyPy- β -HpPy- γ -ImPy- β -ImImPy
	1661 β) 5'-W G C C T A T W-3'	ImPyPy- β -PyHp- γ -PyHp- β -ImImPy
	1662 β) 5'-W G C C T A A W-3'	ImPyPy- β -PyPy- γ -HpHp- β -ImImPy
10	1663 β) 5'-W G C C T A G W-3'	ImPyPy- β -PyIm- γ -PyHp- β -ImImPy
	1664 β) 5'-W G C C T A C W-3'	ImPyPy- β -PyPy- γ -ImHp- β -ImImPy
	1665 β) 5'-W G C C T G T W-3'	ImPyPy- β -ImHp- γ -PyPy- β -ImImPy
	1666 β) 5'-W G C C T G A W-3'	ImPyPy- β -ImPy- γ -HpPy- β -ImImPy
	1667 β) 5'-W G C C T G G W-3'	ImPyPy- β -ImIm- γ -PyPy- β -ImImPy
	1668 β) 5'-W G C C T G C W-3'	ImPyPy- β -ImPy- γ -ImPy- β -ImImPy
	1669 β) 5'-W G C C T C T W-3'	ImPyPy- β -PyHp- γ -PyIm- β -ImImPy
	1670 β) 5'-W G C C T C A W-3'	ImPyPy- β -PyPy- γ -HpIm- β -ImImPy
	1671 β) 5'-W G C C T C G W-3'	ImPyPy- β -PyIm- γ -PyIm- β -ImImPy
	1672 β) 5'-W G C C T C C W-3'	ImPyPy- β -PyPy- γ -ImIm- β -ImImPy
20	1673 β) 5'-W G C C A T T W-3'	ImPyPy- β -HpHp- γ -PyPy- β -ImImPy
	1674 β) 5'-W G C C A T A W-3'	ImPyPy- β -HpPy- γ -HpPy- β -ImImPy
	1675 β) 5'-W G C C A T G W-3'	ImPyPy- β -HpIm- γ -PyPy- β -ImImPy
	1676 β) 5'-W G C C A T C W-3'	ImPyPy- β -HpPy- γ -ImPy- β -ImImPy
	1677 β) 5'-W G C C A A T W-3'	ImPyPy- β -PyHp- γ -PyHp- β -ImImPy
25	1678 β) 5'-W G C C A A A W-3'	ImPyPy- β -PyPy- γ -HpHp- β -ImImPy
	1679 β) 5'-W G C C A A G W-3'	ImPyPy- β -PyIm- γ -PyHp- β -ImImPy
	1680 β) 5'-W G C C A A C W-3'	ImPyPy- β -PyPy- γ -ImHp- β -ImImPy
	1681 β) 5'-W G C C A G T W-3'	ImPyPy- β -ImHp- γ -PyPy- β -ImImPy
	1682 β) 5'-W G C C A G A W-3'	ImPyPy- β -ImPy- γ -HpPy- β -ImImPy
30	1683 β) 5'-W G C C A G G W-3'	ImPyPy- β -ImIm- γ -PyPy- β -ImImPy
	1684 β) 5'-W G C C A G C W-3'	ImPyPy- β -ImPy- γ -ImPy- β -ImImPy
	1685 β) 5'-W G C C A C T W-3'	ImPyPy- β -PyHp- γ -PyIm- β -ImImPy
	1686 β) 5'-W G C C A C A W-3'	ImPyPy- β -PyPy- γ -HpIm- β -ImImPy
	1687 β) 5'-W G C C A C G W-3'	ImPyPy- β -PyIm- γ -PyIm- β -ImImPy
35	1688 β) 5'-W G C C A C C W-3'	ImPyPy- β -PyPy- γ -ImIm- β -ImImPy

TABLE 163: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGCCSNW-3'

	DNA sequence	aromatic amino acid sequence
	1689 β) 5'-W G C C G T T W-3'	ImPy- β -ImHpHp- γ -PyPy- β -ImImPy
5	1690 β) 5'-W G C C G T A W-3'	ImPy- β -ImHpPy- γ -HpPy- β -ImImPy
	1691 β) 5'-W G C C G T G W-3'	ImPy- β -ImHpIm- γ -PyPy- β -ImImPy
	1692 β) 5'-W G C C G T C W-3'	ImPy- β -ImHpPy- γ -ImPy- β -ImImPy
	1693 β) 5'-W G C C G A T W-3'	ImPy- β -ImPyHp- γ -PyHp- β -ImImPy
	1694 β) 5'-W G C C G A A W-3'	ImPy- β -ImPyPy- γ -HpHp- β -ImImPy
10	1695 β) 5'-W G C C G A G W-3'	ImPy- β -ImPyIm- γ -PyHp- β -ImImPy
	1696 β) 5'-W G C C G A C W-3'	ImPy- β -ImPyPy- γ -ImHp- β -ImImPy
	1697 β) 5'-W G C C G G T W-3'	ImPy- β -ImImHp- γ -PyPy- β -ImImPy
	1698 β) 5'-W G C C G G A W-3'	ImPy- β -ImImPy- γ -HpPy- β -ImImPy
	1699 β) 5'-W G C C G C T W-3'	ImPy- β -ImPyHp- γ -PyIm- β -ImImPy
15	1700 β) 5'-W G C C G C A W-3'	ImPy- β -ImPyPy- γ -HpIm- β -ImImPy
	1701 β) 5'-W G C C C T T W-3'	ImPy- β -PyHpHp- γ -Py- β -ImImImPy
	1702 β) 5'-W G C C C T A W-3'	ImPy- β -PyHpPy- γ -Hp- β -ImImImPy
	1703 β) 5'-W G C C C T G W-3'	ImPy- β -PyHpIm- γ -Py- β -ImImImPy
	1704 β) 5'-W G C C C T C W-3'	ImPy- β -PyHpPy- γ -Im- β -ImImImPy
20	1705 β) 5'-W G C C C A T W-3'	ImPy- β -PyPyHp- γ -Py- β -ImImImPy
	1706 β) 5'-W G C C C A A W-3'	ImPy- β -PyPyPy- γ -Hp- β -ImImImPy
	1707 β) 5'-W G C C C A G W-3'	ImPy- β -PyPyIm- γ -Py- β -ImImImPy
	1708 β) 5'-W G C C C A C W-3'	ImPy- β -PyPyPy- γ -Im- β -ImImImPy
	1709 β) 5'-W G C C C G T W-3'	ImPy- β -PyImHp- γ -Py- β -ImImImPy
25	1710 β) 5'-W G C C C G A W-3'	ImPy- β -PyImPy- γ -Hp- β -ImImImPy
	G73 β) 5'-W G C C G G G W-3'	ImPy- β -ImImIm- γ -PyPy- β -ImImPy
	G74 β) 5'-W G C C G G C W-3'	ImPy- β -ImImPy- γ -ImPy- β -ImImPy
	G75 β) 5'-W G C C G C G W-3'	ImPy- β -ImPyIm- γ -PyIm- β -ImImPy
	G76 β) 5'-W G C C G C C W-3'	ImPy- β -ImPyPy- γ -ImIm- β -ImImPy
30	G77 β) 5'-W G C C C G G W-3'	ImPy- β -PyImIm- γ -Py- β -ImImImPy
	G78 β) 5'-W G C C C G C W-3'	ImPy- β -PyImPy- γ -Im- β -ImImImPy

TABLE 164: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGAGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1713 β) 5'-W G A G T T T W-3'	Im- β -ImHpHpHp- γ -PyPyPyPy- β -Py
	1714 β) 5'-W G A G T T A W-3'	Im- β -ImHpHpPy- γ -HpPyPyPy- β -Py
5	1715 β) 5'-W G A G T T G W-3'	Im- β -ImHpHpIm- γ -PyPyPyPy- β -Py
	1716 β) 5'-W G A G T T C W-3'	Im- β -ImHpHpPy- γ -ImPyPyPy- β -Py
	1717 β) 5'-W G A G T A T W-3'	Im- β -ImHpPyHp- γ -PyHpPyPy- β -Py
	1718 β) 5'-W G A G T A A W-3'	Im- β -ImHpPyPy- γ -HpHpPyPy- β -Py
	1719 β) 5'-W G A G T A G W-3'	Im- β -ImHpPyIm- γ -PyHpPyPy- β -Py
10	1720 β) 5'-W G A G T A C W-3'	Im- β -ImHpPyPy- γ -ImHpPyPy- β -Py
	1721 β) 5'-W G A G T G T W-3'	Im- β -ImHpImHp- γ -PyPyPyPy- β -Py
	1722 β) 5'-W G A G T G A W-3'	Im- β -ImHpImPy- γ -HpPyPyPy- β -Py
	1723 β) 5'-W G A G T G G W-3'	Im- β -ImHpImIm- γ -PyPyPyPy- β -Py
	1724 β) 5'-W G A G T G C W-3'	Im- β -ImHpImPy- γ -ImPyPyPy- β -Py
	1725 β) 5'-W G A G T C T W-3'	Im- β -ImHpPyHp- γ -PyImPyPy- β -Py
	1726 β) 5'-W G A G T C A W-3'	Im- β -ImHpPyPy- γ -HpImPyPy- β -Py
	1727 β) 5'-W G A G T C G W-3'	Im- β -ImHpPyIm- γ -PyImPyPy- β -Py
	1728 β) 5'-W G A G T C C W-3'	Im- β -ImHpPyPy- γ -ImImPyPy- β -Py
	1729 β) 5'-W G A G A T T W-3'	Im- β -ImPyHpHp- γ -PyPyHpPy- β -Py
20	1730 β) 5'-W G A G A T A W-3'	Im- β -ImPyHpPy- γ -HpPyHpPy- β -Py
	1731 β) 5'-W G A G A T G W-3'	Im- β -ImPyHpIm- γ -PyPyHpPy- β -Py
	1732 β) 5'-W G A G A T C W-3'	Im- β -ImPyHpPy- γ -ImPyHpPy- β -Py
	1733 β) 5'-W G A G A A T W-3'	Im- β -ImPyPyHp- γ -PyHpHpPy- β -Py
	1734 β) 5'-W G A G A A A W-3'	Im- β -ImPyPyPy- γ -HpHpHpPy- β -Py
25	1735 β) 5'-W G A G A A G W-3'	Im- β -ImPyPyIm- γ -PyHpHpPy- β -Py
	1736 β) 5'-W G A G A A C W-3'	Im- β -ImPyPyPy- γ -ImHpHpPy- β -Py
	1737 β) 5'-W G A G A G T W-3'	Im- β -ImPyImHp- γ -PyPyHpPy- β -Py
	1738 β) 5'-W G A G A G A W-3'	Im- β -ImPyImPy- γ -HpPyHpPy- β -Py
	1739 β) 5'-W G A G A G G W-3'	Im- β -ImPyImIm- γ -PyPyHpPy- β -Py
30	1740 β) 5'-W G A G A G C W-3'	Im- β -ImPyImPy- γ -ImPyHpPy- β -Py
	1741 β) 5'-W G A G A C T W-3'	Im- β -ImPyPyHp- γ -PyImHpPy- β -Py
	1742 β) 5'-W G A G A C A W-3'	Im- β -ImPyPyPy- γ -HpImHpPy- β -Py
	1743 β) 5'-W G A G A C G W-3'	Im- β -ImPyPyIm- γ -PyImHpPy- β -Py
	1744 β) 5'-W G A G A C C W-3'	Im- β -ImPyPyPy- γ -ImImHpPy- β -Py

TABLE 165: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGAGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	1745 β) 5'-W G A G G T T W-3'	Im- β -ImImHpHp- γ -PyPyPyPy- β -Py
5	1746 β) 5'-W G A G G T A W-3'	Im- β -ImImHpPy- γ -HpPyPyPy- β -Py
	1747 β) 5'-W G A G G T G W-3'	Im- β -ImImHpIm- γ -PyPyPyPy- β -Py
	1748 β) 5'-W G A G G T C W-3'	Im- β -ImImHpPy- γ -ImPyPyPy- β -Py
	1749 β) 5'-W G A G G A T W-3'	Im- β -ImImPyHp- γ -PyHpPyPy- β -Py
	1750 β) 5'-W G A G G A A W-3'	Im- β -ImImPyPy- γ -HpHpPyPy- β -Py
10	1751 β) 5'-W G A G G A G W-3'	Im- β -ImImPyIm- γ -PyHpPyPy- β -Py
	1752 β) 5'-W G A G G A C W-3'	Im- β -ImImPyPy- γ -ImHpPyPy- β -Py
	1753 β) 5'-W G A G G G T W-3'	Im- β -ImImImHp- γ -PyPyPyPy- β -Py
	1754 β) 5'-W G A G G G A W-3'	Im- β -ImImImPy- γ -HpPyPyPy- β -Py
	1755 β) 5'-W G A G G C T W-3'	Im- β -ImImPyHp- γ -PyImPyPy- β -Py
	1756 β) 5'-W G A G G C A W-3'	Im- β -ImImPyPy- γ -HpImPyPy- β -Py
	1757 β) 5'-W G A G C T T W-3'	Im- β -ImPyHpHp- γ -PyPyImPy- β -Py
	1758 β) 5'-W G A G C T A W-3'	Im- β -ImPyHpPy- γ -HpPyImPy- β -Py
	1759 β) 5'-W G A G C T G W-3'	Im- β -ImPyHpIm- γ -PyPyImPy- β -Py
	1760 β) 5'-W G A G C T C W-3'	Im- β -ImPyHpPy- γ -ImPyImPy- β -Py
20	1761 β) 5'-W G A G C A T W-3'	Im- β -ImPyPyHp- γ -PyHpImPy- β -Py
	1762 β) 5'-W G A G C A A W-3'	Im- β -ImPyPyPy- γ -HpHpImPy- β -Py
	1763 β) 5'-W G A G C A G W-3'	Im- β -ImPyPyIm- γ -PyHpImPy- β -Py
	1764 β) 5'-W G A G C A C W-3'	Im- β -ImPyPyPy- γ -ImHpImPy- β -Py
	1765 β) 5'-W G A G C G T W-3'	Im- β -ImPyImHp- γ -PyPyImPy- β -Py
25	1766 β) 5'-W G A G C G A W-3'	Im- β -ImPyImPy- γ -HpPyImPy- β -Py
	1767 β) 5'-W G A G C C T W-3'	Im- β -ImPyPyHp- γ -PyImImPy- β -Py
	1768 β) 5'-W G A G C C A W-3'	Im- β -ImPyPyPy- γ -HpImImPy- β -Py
	1769 β) 5'-W G A G G G G W-3'	Im- β -ImImImIm- γ -PyPyPyPy- β -Py
	1770 β) 5'-W G A G G G C W-3'	Im- β -ImImImPy- γ -ImPyPyPy- β -Py
30	1771 β) 5'-W G A G G C G W-3'	Im- β -ImImPyIm- γ -PyImPyPy- β -Py
	1772 β) 5'-W G A G G C C W-3'	Im- β -ImImPyPy- γ -ImImPyPy- β -Py
	1773 β) 5'-W G A G C G G W-3'	Im- β -ImPyImIm- γ -PyPyImPy- β -Py
	1774 β) 5'-W G A G C G C W-3'	Im- β -ImPyImPy- γ -ImPyImPy- β -Py
	1775 β) 5'-W G A G C C G W-3'	Im- β -ImPyPyIm- γ -PyImImPy- β -Py
35	1776 β) 5'-W G A G C C C W-3'	Im- β -ImPyPyPy- γ -ImImImPy- β -Py

TABLE 166: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGATWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1777 β) 5'-W G A T T T W-3'	ImPy- β -HpHpHp- γ -PyPyPy- β -HpPy
5	1778 β) 5'-W G A T T T A W-3'	ImPy- β -HpHpPy- γ -HpPyPy- β -HpPy
	1779 β) 5'-W G A T T T G W-3'	ImPy- β -HpHpIm- γ -PyPyPy- β -HpPy
	1780 β) 5'-W G A T T T C W-3'	ImPy- β -HpHpPy- γ -ImPyPy- β -HpPy
	1781 β) 5'-W G A T T A T W-3'	ImPy- β -HpPyHp- γ -PyHpPy- β -HpPy
	1782 β) 5'-W G A T T A A W-3'	ImPy- β -HpPyPy- γ -HpHpPy- β -HpPy
10	1783 β) 5'-W G A T T A G W-3'	ImPy- β -HpPyIm- γ -PyHpPy- β -HpPy
	1784 β) 5'-W G A T T A C W-3'	ImPy- β -HpPyPy- γ -ImHpPy- β -HpPy
	1785 β) 5'-W G A T T G T W-3'	ImPy- β -HpImHp- γ -PyPyPy- β -HpPy
	1786 β) 5'-W G A T T G A W-3'	ImPy- β -HpImPy- γ -HpPyPy- β -HpPy
	1787 β) 5'-W G A T T G G W-3'	ImPy- β -HpImIm- γ -PyPyPy- β -HpPy
	1788 β) 5'-W G A T T G C W-3'	ImPy- β -HpImPy- γ -ImPyPy- β -HpPy
	1789 β) 5'-W G A T T C T W-3'	ImPy- β -HpPyHp- γ -PyImPy- β -HpPy
	1790 β) 5'-W G A T T C A W-3'	ImPy- β -HpPyPy- γ -HpImPy- β -HpPy
	1791 β) 5'-W G A T T C G W-3'	ImPy- β -HpPyIm- γ -PyImPy- β -HpPy
	1792 β) 5'-W G A T T C C W-3'	ImPy- β -HpPyPy- γ -ImImPy- β -HpPy
20	1793 β) 5'-W G A T A T T W-3'	ImPy- β -PyHpHp- γ -PyPyHp- β -HpPy
	1794 β) 5'-W G A T A T A W-3'	ImPy- β -PyHpPy- γ -HpPyHp- β -HpPy
	1795 β) 5'-W G A T A T G W-3'	ImPy- β -PyHpIm- γ -PyPyHp- β -HpPy
	1796 β) 5'-W G A T A T C W-3'	ImPy- β -PyHpPy- γ -ImPyHp- β -HpPy
	1797 β) 5'-W G A T A A T W-3'	ImPy- β -PyPyHp- γ -PyHpHp- β -HpPy
25	1798 β) 5'-W G A T A A A W-3'	ImPy- β -PyPyPy- γ -HpHpHp- β -HpPy
	1799 β) 5'-W G A T A A G W-3'	ImPy- β -PyPyIm- γ -PyHpHp- β -HpPy
	1800 β) 5'-W G A T A A C W-3'	ImPy- β -PyPyPy- γ -ImHpHp- β -HpPy
	1801 β) 5'-W G A T A G T W-3'	ImPy- β -PyImHp- γ -PyPyHp- β -HpPy
	1802 β) 5'-W G A T A G A W-3'	ImPy- β -PyImPy- γ -HpPyHp- β -HpPy
30	1803 β) 5'-W G A T A G G W-3'	ImPy- β -PyImIm- γ -PyPyHp- β -HpPy
	1804 β) 5'-W G A T A G C W-3'	ImPy- β -PyImPy- γ -ImPyHp- β -HpPy
	1805 β) 5'-W G A T A C T W-3'	ImPy- β -PyPyHp- γ -PyImHp- β -HpPy
	1806 β) 5'-W G A T A C A W-3'	ImPy- β -PyPyPy- γ -HpImHp- β -HpPy
	1807 β) 5'-W G A T A C G W-3'	ImPy- β -PyPyIm- γ -PyImHp- β -HpPy
35	1808 β) 5'-W G A T A C C W-3'	ImPy- β -PyPyPy- γ -ImImHp- β -HpPy

TABLE 167: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGATSNW-3'

	DNA sequence	aromatic amino acid sequence
	1809 β) 5'-W G A T G T T W-3'	ImPy- β -ImHpHp- γ -PyPyPy- β -HpPy
5	1810 β) 5'-W G A T G T A W-3'	ImPy- β -ImHpPy- γ -HpPyPy- β -HpPy
	1811 β) 5'-W G A T G T G W-3'	ImPy- β -ImHpIm- γ -PyPyPy- β -HpPy
	1812 β) 5'-W G A T G T C W-3'	ImPy- β -ImHpPy- γ -ImPyPy- β -HpPy
	1813 β) 5'-W G A T G A T W-3'	ImPy- β -ImPyHp- γ -PyHpPy- β -HpPy
	1814 β) 5'-W G A T G A A W-3'	ImPy- β -ImPyPy- γ -HpHpPy- β -HpPy
10	1815 β) 5'-W G A T G A G W-3'	ImPy- β -ImPyIm- γ -PyHpPy- β -HpPy
	1816 β) 5'-W G A T G A C W-3'	ImPy- β -ImPyPy- γ -ImHpPy- β -HpPy
	1817 β) 5'-W G A T G G T W-3'	ImPy- β -ImImHp- γ -PyPyPy- β -HpPy
	1818 β) 5'-W G A T G G A W-3'	ImPy- β -ImImPy- γ -HpPyPy- β -HpPy
	1819 β) 5'-W G A T G C T W-3'	ImPy- β -ImPyHp- γ -PyImPy- β -HpPy
	1820 β) 5'-W G A T G C A W-3'	ImPy- β -ImPyPy- γ -HpImPy- β -HpPy
	1821 β) 5'-W G A T G G G W-3'	ImPy- β -ImImIm- γ -PyPyPy- β -HpPy
	1822 β) 5'-W G A T G G C W-3'	ImPy- β -ImImPy- γ -ImPyPy- β -HpPy
	1823 β) 5'-W G A T G C G W-3'	ImPy- β -ImPyIm- γ -PyImPy- β -HpPy
	1824 β) 5'-W G A T G C C W-3'	ImPy- β -ImPyPy- γ -ImImPy- β -HpPy
20	1825 β) 5'-W G A T C T T W-3'	ImPy- β -PyHpHp- γ -PyPyIm- β -HpPy
	1826 β) 5'-W G A T C T A W-3'	ImPy- β -PyHpPy- γ -HpPyIm- β -HpPy
	1827 β) 5'-W G A T C T G W-3'	ImPy- β -PyHpIm- γ -PyPyIm- β -HpPy
	1828 β) 5'-W G A T C T C W-3'	ImPy- β -PyHpPy- γ -ImPyIm- β -HpPy
	1829 β) 5'-W G A T C A T W-3'	ImPy- β -PyPyHp- γ -PyHpIm- β -HpPy
25	1830 β) 5'-W G A T C A A W-3'	ImPy- β -PyPyPy- γ -HpHpIm- β -HpPy
	1831 β) 5'-W G A T C A G W-3'	ImPy- β -PyPyIm- γ -PyHpIm- β -HpPy
	1832 β) 5'-W G A T C A C W-3'	ImPy- β -PyPyPy- γ -ImHpIm- β -HpPy
	1833 β) 5'-W G A T C G T W-3'	ImPy- β -PyImHp- γ -PyPyIm- β -HpPy
	1834 β) 5'-W G A T C G A W-3'	ImPy- β -PyImPy- γ -HpPyIm- β -HpPy
30	1835 β) 5'-W G A T C C T W-3'	ImPy- β -PyPyHp- γ -PyImIm- β -HpPy
	1836 β) 5'-W G A T C C A W-3'	ImPy- β -PyPyPy- γ -HpImIm- β -HpPy
	1837 β) 5'-W G A T C G G W-3'	ImPy- β -PyImIm- γ -PyPyIm- β -HpPy
	1838 β) 5'-W G A T C G C W-3'	ImPy- β -PyImPy- γ -ImPyIm- β -HpPy
	1839 β) 5'-W G A T C C G W-3'	ImPy- β -PyPyIm- γ -PyImIm- β -HpPy
35	1840 β) 5'-W G A T C C C W-3'	ImPy- β -PyPyPy- γ -ImImIm- β -HpPy

TABLE 168: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGAAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1841 β) 5'-W G A A T T T W-3'	ImPy- β -HpHpHp- γ -PyPyPy- β -HpPy
5	1842 β) 5'-W G A A T T A W-3'	ImPy- β -HpHpPy- γ -HpPyPy- β -HpPy
	1843 β) 5'-W G A A T T G W-3'	ImPy- β -HpHpIm- γ -PyPyPy- β -HpPy
	1844 β) 5'-W G A A T T C W-3'	ImPy- β -HpHpPy- γ -ImPyPy- β -HpPy
	1845 β) 5'-W G A A T A T W-3'	ImPy- β -HpPyHp- γ -PyHpPy- β -HpPy
	1846 β) 5'-W G A A T A A W-3'	ImPy- β -HpPyPy- γ -HpHpPy- β -HpPy
10	1847 β) 5'-W G A A T A G W-3'	ImPy- β -HpPyIm- γ -PyHpPy- β -HpPy
	1848 β) 5'-W G A A T A C W-3'	ImPy- β -HpPyPy- γ -ImHpPy- β -HpPy
	1849 β) 5'-W G A A T G T W-3'	ImPy- β -HpImHp- γ -PyPyPy- β -HpPy
	1850 β) 5'-W G A A T G A W-3'	ImPy- β -HpImPy- γ -HpPyPy- β -HpPy
	1851 β) 5'-W G A A T G G W-3'	ImPy- β -HpImIm- γ -PyPyPy- β -HpPy
	1852 β) 5'-W G A A T G C W-3'	ImPy- β -HpImPy- γ -ImPyPy- β -HpPy
	1853 β) 5'-W G A A T C T W-3'	ImPy- β -HpPyHp- γ -PyImPy- β -HpPy
	1854 β) 5'-W G A A T C A W-3'	ImPy- β -HpPyPy- γ -HpImPy- β -HpPy
	1855 β) 5'-W G A A T C G W-3'	ImPy- β -HpPyIm- γ -PyImPy- β -HpPy
	1856 β) 5'-W G A A T C C W-3'	ImPy- β -HpPyPy- γ -ImImPy- β -HpPy
20	1857 β) 5'-W G A A A T T W-3'	ImPy- β -PyHpHp- γ -PyPyHp- β -HpPy
	1858 β) 5'-W G A A A T A W-3'	ImPy- β -PyHpPy- γ -HpPyHp- β -HpPy
	1869 β) 5'-W G A A A T G W-3'	ImPy- β -PyHpIm- γ -PyPyHp- β -HpPy
	1860 β) 5'-W G A A A T C W-3'	ImPy- β -PyHpPy- γ -ImPyHp- β -HpPy
	1861 β) 5'-W G A A A A T W-3'	ImPy- β -PyPyHp- γ -PyHpHp- β -HpPy
25	1862 β) 5'-W G A A A A A W-3'	ImPy- β -PyPyPy- γ -HpHpHp- β -HpPy
	1863 β) 5'-W G A A A A G W-3'	ImPy- β -PyPyIm- γ -PyHpHp- β -HpPy
	1864 β) 5'-W G A A A A C W-3'	ImPy- β -PyPyPy- γ -ImHpHp- β -HpPy
	1865 β) 5'-W G A A A G T W-3'	ImPy- β -PyImHp- γ -PyPyHp- β -HpPy
	1866 β) 5'-W G A A A G A W-3'	ImPy- β -PyImPy- γ -HpPyHp- β -HpPy
30	1867 β) 5'-W G A A A G G W-3'	ImPy- β -PyImIm- γ -PyPyHp- β -HpPy
	1868 β) 5'-W G A A A G C W-3'	ImPy- β -PyImPy- γ -ImPyHp- β -HpPy
	1869 β) 5'-W G A A A C T W-3'	ImPy- β -PyPyHp- γ -PyImHp- β -HpPy
	1870 β) 5'-W G A A A C A W-3'	ImPy- β -PyPyPy- γ -HpImHp- β -HpPy
	1871 β) 5'-W G A A A C G W-3'	ImPy- β -PyPyIm- γ -PyImHp- β -HpPy
35	1872 β) 5'-W G A A A C C W-3'	ImPy- β -PyPyPy- γ -ImImHp- β -HpPy

TABLE 169: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGAASNNW-3'

	DNA sequence	aromatic amino acid sequence
	1873 β) 5'-W G A A G T T W-3'	ImPy- β -ImHpHp- γ -PyPyPy- β -HpPy
5	1874 β) 5'-W G A A G T A W-3'	ImPy- β -ImHpPy- γ -HpPyPy- β -HpPy
	1875 β) 5'-W G A A G T G W-3'	ImPy- β -ImHpIm- γ -PyPyPy- β -HpPy
	1876 β) 5'-W G A A G T C W-3'	ImPy- β -ImHpPy- γ -ImPyPy- β -HpPy
	1877 β) 5'-W G A A G A T W-3'	ImPy- β -ImPyHp- γ -PyHpPy- β -HpPy
	1878 β) 5'-W G A A G A A W-3'	ImPy- β -ImPyPy- γ -HpHpPy- β -HpPy
10	1879 β) 5'-W G A A G A G W-3'	ImPy- β -ImPyIm- γ -PyHpPy- β -HpPy
	1880 β) 5'-W G A A G A C W-3'	ImPy- β -ImPyPy- γ -ImHpPy- β -HpPy
	1881 β) 5'-W G A A G G T W-3'	ImPy- β -ImImHp- γ -PyPyPy- β -HpPy
	1882 β) 5'-W G A A G G A W-3'	ImPy- β -ImImPy- γ -HpPyPy- β -HpPy
	1883 β) 5'-W G A A G C T W-3'	ImPy- β -ImPyHp- γ -PyImPy- β -HpPy
	1884 β) 5'-W G A A G C A W-3'	ImPy- β -ImPyPy- γ -HpImPy- β -HpPy
	1885 β) 5'-W G A A G G G W-3'	ImPy- β -ImImIm- γ -PyPyPy- β -HpPy
	1886 β) 5'-W G A A G G C W-3'	ImPy- β -ImImPy- γ -ImPyPy- β -HpPy
	1887 β) 5'-W G A A G C G W-3'	ImPy- β -ImPyIm- γ -PyImPy- β -HpPy
	1888 β) 5'-W G A A G C C W-3'	ImPy- β -ImPyPy- γ -ImImPy- β -HpPy
20	1889 β) 5'-W G A A C T T W-3'	ImPy- β -PyHpHp- γ -PyPyIm- β -HpPy
	1890 β) 5'-W G A A C T A W-3'	ImPy- β -PyHpPy- γ -HpPyIm- β -HpPy
	1891 β) 5'-W G A A C T G W-3'	ImPy- β -PyHpIm- γ -PyPyIm- β -HpPy
	1892 β) 5'-W G A A C T C W-3'	ImPy- β -PyHpPy- γ -ImPyIm- β -HpPy
	1893 β) 5'-W G A A C A T W-3'	ImPy- β -PyPyHp- γ -PyHpIm- β -HpPy
25	1894 β) 5'-W G A A C A A W-3'	ImPy- β -PyPyPy- γ -HpHpIm- β -HpPy
	1895 β) 5'-W G A A C A G W-3'	ImPy- β -PyPyIm- γ -PyHpIm- β -HpPy
	1896 β) 5'-W G A A C A C W-3'	ImPy- β -PyPyPy- γ -ImHpIm- β -HpPy
	1897 β) 5'-W G A A C G T W-3'	ImPy- β -PyImHp- γ -PyPyIm- β -HpPy
	1898 β) 5'-W G A A C G A W-3'	ImPy- β -PyImPy- γ -HpPyIm- β -HpPy
30	1899 β) 5'-W G A A C C T W-3'	ImPy- β -PyPyHp- γ -PyImIm- β -HpPy
	1900 β) 5'-W G A A C C A W-3'	ImPy- β -PyPyPy- γ -HpImIm- β -HpPy
	1901 β) 5'-W G A A C G G W-3'	ImPy- β -PyImIm- γ -PyPyIm- β -HpPy
	1902 β) 5'-W G A A C G C W-3'	ImPy- β -PyImPy- γ -ImPyIm- β -HpPy
	1903 β) 5'-W G A A C C G W-3'	ImPy- β -PyPyIm- γ -PyImIm- β -HpPy
35	1904 β) 5'-W G A A C C C W-3'	ImPy- β -PyPyPy- γ -ImImIm- β -HpPy

TABLE 170: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGACWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1905 β) 5'-W G A C T T T W-3'	ImPyPy- β -HpHp- γ -PyPy- β -ImHpPy
5	1906 β) 5'-W G A C T T A W-3'	ImPyPy- β -HpPy- γ -HpPy- β -ImHpPy
	1907 β) 5'-W G A C T T G W-3'	ImPyPy- β -HpIm- γ -PyPy- β -ImHpPy
	1908 β) 5'-W G A C T T C W-3'	ImPyPy- β -HpPy- γ -ImPy- β -ImHpPy
	1909 β) 5'-W G A C T A T W-3'	ImPyPy- β -PyHp- γ -PyHp- β -ImHpPy
	1910 β) 5'-W G A C T A A W-3'	ImPyPy- β -PyPy- γ -HpHp- β -ImHpPy
10	1911 β) 5'-W G A C T A G W-3'	ImPyPy- β -PyIm- γ -PyHp- β -ImHpPy
	1912 β) 5'-W G A C T A C W-3'	ImPyPy- β -PyPy- γ -ImHp- β -ImHpPy
	1913 β) 5'-W G A C T G T W-3'	ImPyPy- β -ImHp- γ -PyPy- β -ImHpPy
	1914 β) 5'-W G A C T G A W-3'	ImPyPy- β -ImPy- γ -HpPy- β -ImHpPy
	1915 β) 5'-W G A C T G G W-3'	ImPyPy- β -ImIm- γ -PyPy- β -ImHpPy
15	1916 β) 5'-W G A C T G C W-3'	ImPyPy- β -ImPy- γ -ImPy- β -ImHpPy
	1917 β) 5'-W G A C T C T W-3'	ImPyPy- β -PyHp- γ -PyIm- β -ImHpPy
	1918 β) 5'-W G A C T C A W-3'	ImPyPy- β -PyPy- γ -HpIm- β -ImHpPy
	1919 β) 5'-W G A C T C G W-3'	ImPyPy- β -PyIm- γ -PyIm- β -ImHpPy
	1920 β) 5'-W G A C T C C W-3'	ImPyPy- β -PyPy- γ -ImIm- β -ImHpPy
20	1921 β) 5'-W G A C A T T W-3'	ImPyPy- β -HpHp- γ -PyPy- β -ImHpPy
	1922 β) 5'-W G A C A T A W-3'	ImPyPy- β -HpPy- γ -HpPy- β -ImHpPy
	1923 β) 5'-W G A C A T G W-3'	ImPyPy- β -HpIm- γ -PyPy- β -ImHpPy
	1924 β) 5'-W G A C A T C W-3'	ImPyPy- β -HpPy- γ -ImPy- β -ImHpPy
	1925 β) 5'-W G A C A A T W-3'	ImPyPy- β -PyHp- γ -PyHp- β -ImHpPy
25	1926 β) 5'-W G A C A A A W-3'	ImPyPy- β -PyPy- γ -HpHp- β -ImHpPy
	1927 β) 5'-W G A C A A G W-3'	ImPyPy- β -PyIm- γ -PyHp- β -ImHpPy
	1928 β) 5'-W G A C A A C W-3'	ImPyPy- β -PyPy- γ -ImHp- β -ImHpPy
	1929 β) 5'-W G A C A G T W-3'	ImPyPy- β -ImHp- γ -PyPy- β -ImHpPy
	1930 β) 5'-W G A C A G A W-3'	ImPyPy- β -ImPy- γ -HpPy- β -ImHpPy
30	1931 β) 5'-W G A C A G G W-3'	ImPyPy- β -ImIm- γ -PyPy- β -ImHpPy
	1932 β) 5'-W G A C A G C W-3'	ImPyPy- β -ImPy- γ -ImPy- β -ImHpPy
	1933 β) 5'-W G A C A C T W-3'	ImPyPy- β -PyHp- γ -PyIm- β -ImHpPy
	1934 β) 5'-W G A C A C A W-3'	ImPyPy- β -PyPy- γ -HpIm- β -ImHpPy
	1935 β) 5'-W G A C A C G W-3'	ImPyPy- β -PyIm- γ -PyIm- β -ImHpPy
35	1936 β) 5'-W G A C A C C W-3'	ImPyPy- β -PyPy- γ -ImIm- β -ImHpPy

TABLE 171: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGACSNW-3'

	DNA sequence	aromatic amino acid sequence
	1937 β) 5'-W G A C G T T W-3'	ImPy- β -ImHpHp- γ -PyPy- β -ImHpPy
5	1938 β) 5'-W G A C G T A W-3'	ImPy- β -ImHpPy- γ -HpPy- β -ImHpPy
	1939 β) 5'-W G A C G T G W-3'	ImPy- β -ImHpIm- γ -PyPy- β -ImHpPy
	1940 β) 5'-W G A C G T C W-3'	ImPy- β -ImHpPy- γ -ImPy- β -ImHpPy
	1941 β) 5'-W G A C G A T W-3'	ImPy- β -ImPyHp- γ -PyHp- β -ImHpPy
	1942 β) 5'-W G A C G A A W-3'	ImPy- β -ImPyPy- γ -HpHp- β -ImHpPy
10	1943 β) 5'-W G A C G A G W-3'	ImPy- β -ImPyIm- γ -PyHp- β -ImHpPy
	1944 β) 5'-W G A C G A C W-3'	ImPy- β -ImPyPy- γ -ImHp- β -ImHpPy
	1945 β) 5'-W G A C G G T W-3'	ImPy- β -ImImHp- γ -PyPy- β -ImHpPy
	1946 β) 5'-W G A C G G A W-3'	ImPy- β -ImImPy- γ -HpPy- β -ImHpPy
	1947 β) 5'-W G A C G C T W-3'	ImPy- β -ImPyHp- γ -PyIm- β -ImHpPy
	1948 β) 5'-W G A C G C A W-3'	ImPy- β -ImPyPy- γ -HpIm- β -ImHpPy
	1949 β) 5'-W G A C C T T W-3'	ImPy- β -PyHpHp- γ -Py- β -ImImHpPy
	1950 β) 5'-W G A C C T A W-3'	ImPy- β -PyHpPy- γ -Hp- β -ImImHpPy
	1951 β) 5'-W G A C C T G W-3'	ImPy- β -PyHpIm- γ -Py- β -ImImHpPy
	1952 β) 5'-W G A C C T C W-3'	ImPy- β -PyHpPy- γ -Im- β -ImImHpPy
20	1953 β) 5'-W G A C C A T W-3'	ImPy- β -PyPyHp- γ -Py- β -ImImHpPy
	1954 β) 5'-W G A C C A A W-3'	ImPy- β -PyPyPy- γ -Hp- β -ImImHpPy
	1955 β) 5'-W G A C C A G W-3'	ImPy- β -PyPyIm- γ -Py- β -ImImHpPy
	1956 β) 5'-W G A C C A C W-3'	ImPy- β -PyPyPy- γ -Im- β -ImImHpPy
	1957 β) 5'-W G A C C G T W-3'	ImPy- β -PyImHp- γ -Py- β -ImImHpPy
25	1958 β) 5'-W G A C C G A W-3'	ImPy- β -PyImPy- γ -Hp- β -ImImHpPy
	1959 β) 5'-W G A C C C T W-3'	ImPy- β -PyPyHp- γ -PyImImIm- β -Py
	1960 β) 5'-W G A C C C A W-3'	ImPy- β -PyPyPy- γ -HpImImIm- β -Py
	1961 β) 5'-W G A C G G G W-3'	ImPy- β -ImImIm- γ -PyPy- β -ImHpPy
	1962 β) 5'-W G A C G G C W-3'	ImPy- β -ImImPy- γ -ImPy- β -ImHpPy
30	1963 β) 5'-W G A C G C G W-3'	ImPy- β -ImPyIm- γ -PyIm- β -ImHpPy
	1964 β) 5'-W G A C G C C W-3'	ImPy- β -ImPyPy- γ -ImIm- β -ImHpPy
	1965 β) 5'-W G A C C G G W-3'	ImPy- β -PyImIm- γ -Py- β -ImImHpPy
	1966 β) 5'-W G A C C G C W-3'	ImPy- β -PyImPy- γ -Im- β -ImImHpPy
	1967 β) 5'-W G A C C C G W-3'	ImPy- β -PyPyIm- γ -PyImImIm- β -Py
35	1968 β) 5'-W G A C C C C W-3'	ImPy- β -PyPyPy- γ -ImImImIm- β -Py

TABLE 172: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTGWNNW-3'

	DNA sequence	aromatic amino acid sequence
	1969 β) 5'-W G T G T T T W-3'	Im- β -ImHpHpHp- γ -PyPyPyPy- β -Py
5	1970 β) 5'-W G T G T T A W-3'	Im- β -ImHpHpPy- γ -HpPyPyPy- β -Py
	1971 β) 5'-W G T G T T G W-3'	Im- β -ImHpHpIm- γ -PyPyPyPy- β -Py
	1972 β) 5'-W G T G T T C W-3'	Im- β -ImHpHpPy- γ -ImPyPyPy- β -Py
	1973 β) 5'-W G T G T A T W-3'	Im- β -ImHpPyHp- γ -PyHpPyPy- β -Py
	1974 β) 5'-W G T G T A A W-3'	Im- β -ImHpPyPy- γ -HpHpPyPy- β -Py
10	1975 β) 5'-W G T G T A G W-3'	Im- β -ImHpPyIm- γ -PyHpPyPy- β -Py
	1976 β) 5'-W G T G T A C W-3'	Im- β -ImHpPyPy- γ -ImHpPyPy- β -Py
	1977 β) 5'-W G T G T G T W-3'	Im- β -ImHpImHp- γ -PyPyPyPy- β -Py
	1978 β) 5'-W G T G T G A W-3'	Im- β -ImHpImPy- γ -HpPyPyPy- β -Py
	1979 β) 5'-W G T G T G G W-3'	Im- β -ImHpImIm- γ -PyPyPyPy- β -Py
	1980 β) 5'-W G T G T G C W-3'	Im- β -ImHpImPy- γ -ImPyPyPy- β -Py
	1981 β) 5'-W G T G T C T W-3'	Im- β -ImHpPyHp- γ -PyImPyPy- β -Py
	1982 β) 5'-W G T G T C A W-3'	Im- β -ImHpPyPy- γ -HpImPyPy- β -Py
	1983 β) 5'-W G T G T C G W-3'	Im- β -ImHpPyIm- γ -PyImPyPy- β -Py
	1984 β) 5'-W G T G T C C W-3'	Im- β -ImHpPyPy- γ -ImImPyPy- β -Py
20	1985 β) 5'-W G T G A T T W-3'	Im- β -ImPyHpHp- γ -PyPyHpPy- β -Py
	1986 β) 5'-W G T G A T A W-3'	Im- β -ImPyHpPy- γ -HpPyHpPy- β -Py
	1987 β) 5'-W G T G A T G W-3'	Im- β -ImPyHpIm- γ -PyPyHpPy- β -Py
	1988 β) 5'-W G T G A T C W-3'	Im- β -ImPyHpPy- γ -ImPyHpPy- β -Py
	1989 β) 5'-W G T G A A T W-3'	Im- β -ImPyPyHp- γ -PyHpHpPy- β -Py
25	1990 β) 5'-W G T G A A A W-3'	Im- β -ImPyPyPy- γ -HpHpHpPy- β -Py
	1991 β) 5'-W G T G A A G W-3'	Im- β -ImPyPyIm- γ -PyHpHpPy- β -Py
	1992 β) 5'-W G T G A A C W-3'	Im- β -ImPyPyPy- γ -ImHpHpPy- β -Py
	1993 β) 5'-W G T G A G T W-3'	Im- β -ImPyImHp- γ -PyPyHpPy- β -Py
	1994 β) 5'-W G T G A G A W-3'	Im- β -ImPyImPy- γ -HpPyHpPy- β -Py
30	1995 β) 5'-W G T G A G G W-3'	Im- β -ImPyImIm- γ -PyPyHpPy- β -Py
	1996 β) 5'-W G T G A G C W-3'	Im- β -ImPyImPy- γ -ImPyHpPy- β -Py
	1997 β) 5'-W G T G A C T W-3'	Im- β -ImPyPyHp- γ -PyImHpPy- β -Py
	1998 β) 5'-W G T G A C A W-3'	Im- β -ImPyPyPy- γ -HpImHpPy- β -Py
	1999 β) 5'-W G T G A C G W-3'	Im- β -ImPyPyIm- γ -PyImHpPy- β -Py
35	2000 β) 5'-W G T G A C C W-3'	Im- β -ImPyPyPy- γ -ImImHpPy- β -Py

TABLE 173: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTGSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2001 β) 5'-W G T G G T T W-3'	Im- β -ImImHpHp- γ -PyPyPyPy- β -Py
5	2002 β) 5'-W G T G G T A W-3'	Im- β -ImImHpPy- γ -HpPyPyPy- β -Py
	2003 β) 5'-W G T G G T G W-3'	Im- β -ImImHpIm- γ -PyPyPyPy- β -Py
	2004 β) 5'-W G T G G T C W-3'	Im- β -ImImHpPy- γ -ImPyPyPy- β -Py
	2005 β) 5'-W G T G G A T W-3'	Im- β -ImImPyHp- γ -PyHpPyPy- β -Py
	2006 β) 5'-W G T G G A A W-3'	Im- β -ImImPyPy- γ -HpHpPyPy- β -Py
10	2007 β) 5'-W G T G G A G W-3'	Im- β -ImImPyIm- γ -PyHpPyPy- β -Py
	2008 β) 5'-W G T G G A C W-3'	Im- β -ImImPyPy- γ -ImHpPyPy- β -Py
	2009 β) 5'-W G T G G G T W-3'	Im- β -ImImImHp- γ -PyPyPyPy- β -Py
	2010 β) 5'-W G T G G G A W-3'	Im- β -ImImImPy- γ -HpPyPyPy- β -Py
	2011 β) 5'-W G T G G C T W-3'	Im- β -ImImPyHp- γ -PyImPyPy- β -Py
	2012 β) 5'-W G T G G C A W-3'	Im- β -ImImPyPy- γ -HpImPyPy- β -Py
	2013 β) 5'-W G T G C T T W-3'	Im- β -ImPyHpHp- γ -PyPyImPy- β -Py
	2014 β) 5'-W G T G C T A W-3'	Im- β -ImPyHpPy- γ -HpPyImPy- β -Py
	2015 β) 5'-W G T G C T G W-3'	Im- β -ImPyHpIm- γ -PyPyImPy- β -Py
	2016 β) 5'-W G T G C T C W-3'	Im- β -ImPyHpPy- γ -ImPyImPy- β -Py
20	2017 β) 5'-W G T G C A T W-3'	Im- β -ImPyPyHp- γ -PyHpImPy- β -Py
	2018 β) 5'-W G T G C A A W-3'	Im- β -ImPyPyPy- γ -HpHpImPy- β -Py
	2019 β) 5'-W G T G C A G W-3'	Im- β -ImPyPyIm- γ -PyHpImPy- β -Py
	2020 β) 5'-W G T G C A C W-3'	Im- β -ImPyPyPy- γ -ImHpImPy- β -Py
	2021 β) 5'-W G T G C G T W-3'	Im- β -ImPyImHp- γ -PyPyImPy- β -Py
25	2022 β) 5'-W G T G C G A W-3'	Im- β -ImPyImPy- γ -HpPyImPy- β -Py
	2023 β) 5'-W G T G C C T W-3'	Im- β -ImPyPyHp- γ -PyImImPy- β -Py
	2024 β) 5'-W G T G C C A W-3'	Im- β -ImPyPyPy- γ -HpImImPy- β -Py
	2025 β) 5'-W G T G G G G W-3'	Im- β -ImImImIm- γ -PyPyPyPy- β -Py
	2026 β) 5'-W G T G G G C W-3'	Im- β -ImImImPy- γ -ImPyPyPy- β -Py
30	2027 β) 5'-W G T G G C G W-3'	Im- β -ImImPyIm- γ -PyImPyPy- β -Py
	2028 β) 5'-W G T G G C C W-3'	Im- β -ImImPyPy- γ -ImImPyPy- β -Py
	2029 β) 5'-W G T G C G G W-3'	Im- β -mPyImIm- γ -PyPyImPy- β -Py
	2030 β) 5'-W G T G C G C W-3'	Im- β -ImPyImPy- γ -ImPyImPy- β -Py
	2031 β) 5'-W G T G C C G W-3'	Im- β -ImPyPyIm- γ -PyImImPy- β -Py
35	2032 β) 5'-W G T G C C C W-3'	Im- β -ImPyPyPy- γ -ImImImPy- β -Py

TABLE 174: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2033 β) 5'-W G T T T T W-3'	ImHp- β -HpHpHp- γ -PyPyPy- β -PyPy
5	2034 β) 5'-W G T T T T A W-3'	ImHp- β -HpHpPy- γ -HpPyPy- β -PyPy
	2035 β) 5'-W G T T T T G W-3'	ImHp- β -HpHpIm- γ -PyPyPy- β -PyPy
	2036 β) 5'-W G T T T T C W-3'	ImHp- β -HpHpPy- γ -ImPyPy- β -PyPy
	2037 β) 5'-W G T T T A T W-3'	ImHp- β -HpPyHp- γ -PyHpPy- β -PyPy
	2038 β) 5'-W G T T T A A W-3'	ImHp- β -HpPyPy- γ -HpHpPy- β -PyPy
10	2039 β) 5'-W G T T T A G W-3'	ImHp- β -HpPyIm- γ -PyHpPy- β -PyPy
	2040 β) 5'-W G T T T A C W-3'	ImHp- β -HpPyPy- γ -ImHpPy- β -PyPy
	2041 β) 5'-W G T T T G T W-3'	ImHp- β -HpImHp- γ -PyPyPy- β -PyPy
	2042 β) 5'-W G T T T G A W-3'	ImHp- β -HpImPy- γ -HpPyPy- β -PyPy
	2043 β) 5'-W G T T T G G W-3'	ImHp- β -HpImIm- γ -PyPyPy- β -PyPy
	2044 β) 5'-W G T T T G C W-3'	ImHp- β -HpImPy- γ -ImPyPy- β -PyPy
	2045 β) 5'-W G T T T C T W-3'	ImHp- β -HpPyHp- γ -PyImPy- β -PyPy
	2046 β) 5'-W G T T T C A W-3'	ImHp- β -HpPyPy- γ -HpImPy- β -PyPy
	2047 β) 5'-W G T T T C G W-3'	ImHp- β -HpPyIm- γ -PyImPy- β -PyPy
	2048 β) 5'-W G T T T C C W-3'	ImHp- β -HpPyPy- γ -ImImPy- β -PyPy
20	2049 β) 5'-W G T T A T T W-3'	ImHp- β -PyHpHp- γ -PyPyHp- β -PyPy
	2050 β) 5'-W G T T A T A W-3'	ImHp- β -PyHpPy- γ -HpPyHp- β -PyPy
	2051 β) 5'-W G T T A T G W-3'	ImHp- β -PyHpIm- γ -PyPyHp- β -PyPy
	2052 β) 5'-W G T T A T C W-3'	ImHp- β -PyHpPy- γ -ImPyHp- β -PyPy
	2053 β) 5'-W G T T A A T W-3'	ImHp- β -PyPyHp- γ -PyHpHp- β -PyPy
25	2054 β) 5'-W G T T A A A W-3'	ImHp- β -PyPyPy- γ -HpHpHp- β -PyPy
	2055 β) 5'-W G T T A A G W-3'	ImHp- β -PyPyIm- γ -PyHpHp- β -PyPy
	2056 β) 5'-W G T T A A C W-3'	ImHp- β -PyPyPy- γ -ImHpHp- β -PyPy
	2057 β) 5'-W G T T A G T W-3'	ImHp- β -PyImHp- γ -PyPyHp- β -PyPy
	2058 β) 5'-W G T T A G A W-3'	ImHp- β -PyImPy- γ -HpPyHp- β -PyPy
30	2059 β) 5'-W G T T A G G W-3'	ImHp- β -PyImIm- γ -PyPyHp- β -PyPy
	2060 β) 5'-W G T T A G C W-3'	ImHp- β -PyImPy- γ -ImPyHp- β -PyPy
	2061 β) 5'-W G T T A C T W-3'	ImHp- β -PyPyHp- γ -PyImHp- β -PyPy
	2062 β) 5'-W G T T A C A W-3'	ImHp- β -PyPyPy- γ -HpImHp- β -PyPy
	2063 β) 5'-W G T T A C G W-3'	ImHp- β -PyPyIm- γ -PyImHp- β -PyPy
35	2064 β) 5'-W G T T A C C W-3'	ImHp- β -PyPyPy- γ -ImImHp- β -PyPy

TABLE 175: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTTSNNW-3'

	DNA sequence	aromatic amino acid sequence
	2065 β) 5'-W G T T G T T W-3'	ImHp- β -ImHpHp- γ -PyPyPy- β -PyPy
5	2066 β) 5'-W G T T G T A W-3'	ImHp- β -ImHpPy- γ -HpPyPy- β -PyPy
	2067 β) 5'-W G T T G T G W-3'	ImHp- β -ImHpIm- γ -PyPyPy- β -PyPy
	2068 β) 5'-W G T T G T C W-3'	ImHp- β -ImHpPy- γ -ImPyPy- β -PyPy
	2069 β) 5'-W G T T G A T W-3'	ImHp- β -ImPyHp- γ -PyHpPy- β -PyPy
	2070 β) 5'-W G T T G A A W-3'	ImHp- β -ImPyPy- γ -HpHpPy- β -PyPy
10	2071 β) 5'-W G T T G A G W-3'	ImHp- β -ImPyIm- γ -PyHpPy- β -PyPy
	2072 β) 5'-W G T T G A C W-3'	ImHp- β -ImPyPy- γ -ImHpPy- β -PyPy
	2073 β) 5'-W G T T G G T W-3'	ImHp- β -ImImHp- γ -PyPyPy- β -PyPy
	2074 β) 5'-W G T T G G A W-3'	ImHp- β -ImImPy- γ -HpPyPy- β -PyPy
	2075 β) 5'-W G T T G C T W-3'	ImHp- β -ImPyHp- γ -PyImPy- β -PyPy
	2076 β) 5'-W G T T G C A W-3'	ImHp- β -ImPyPy- γ -HpImPy- β -PyPy
	2077 β) 5'-W G T T G G G W-3'	ImHp- β -ImImIm- γ -PyPyPy- β -PyPy
	2078 β) 5'-W G T T G G C W-3'	ImHp- β -ImImPy- γ -ImPyPy- β -PyPy
	2079 β) 5'-W G T T G C G W-3'	ImHp- β -ImPyIm- γ -PyImPy- β -PyPy
	2080 β) 5'-W G T T G C C W-3'	ImHp- β -ImPyPy- γ -ImImPy- β -PyPy
	2081 β) 5'-W G T T C T T W-3'	ImHp- β -PyHpHp- γ -PyPyIm- β -PyPy
	2082 β) 5'-W G T T C T A W-3'	ImHp- β -PyHpPy- γ -HpPyIm- β -PyPy
	2083 β) 5'-W G T T C T G W-3'	ImHp- β -PyHpIm- γ -PyPyIm- β -PyPy
	2084 β) 5'-W G T T C T C W-3'	ImHp- β -PyHpPy- γ -ImPyIm- β -PyPy
	2085 β) 5'-W G T T C A T W-3'	ImHp- β -PyPyHp- γ -PyHpIm- β -PyPy
25	2086 β) 5'-W G T T C A A W-3'	ImHp- β -PyPyPy- γ -HpHpIm- β -PyPy
	2087 β) 5'-W G T T C A G W-3'	ImHp- β -PyPyIm- γ -PyHpIm- β -PyPy
	2088 β) 5'-W G T T C A C W-3'	ImHp- β -PyPyPy- γ -ImHpIm- β -PyPy
	2089 β) 5'-W G T T C G T W-3'	ImHp- β -PyImHp- γ -PyPyIm- β -PyPy
	2090 β) 5'-W G T T C G A W-3'	ImHp- β -PyImPy- γ -HpPyIm- β -PyPy
30	2091 β) 5'-W G T T C C T W-3'	ImHp- β -PyPyHp- γ -PyImIm- β -PyPy
	2092 β) 5'-W G T T C C A W-3'	ImHp- β -PyPyPy- γ -HpImIm- β -PyPy
	2093 β) 5'-W G T T C G G W-3'	ImHp- β -PyImIm- γ -PyPyIm- β -PyPy
	2094 β) 5'-W G T T C G C W-3'	ImHp- β -PyImPy- γ -ImPyIm- β -PyPy
	2095 β) 5'-W G T T C C G W-3'	ImHp- β -PyPyIm- γ -PyImIm- β -PyPy
35	2096 β) 5'-W G T T C C C W-3'	ImHp- β -PyPyPy- γ -ImImIm- β -PyPy

TABLE 176: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTAWNNW-3'

	DNA sequence	aromatic amino acid sequence
	2097 β) 5'-W G T A T T T W-3'	ImHp- β -HpHpHp- γ -PyPyPy- β -PyPy
5	2098 β) 5'-W G T A T T A W-3'	ImHp- β -HpHpPy- γ -HpPyPy- β -PyPy
	2099 β) 5'-W G T A T T G W-3'	ImHp- β -HpHpIm- γ -PyPyPy- β -PyPy
	2100 β) 5'-W G T A T T C W-3'	ImHp- β -HpHpPy- γ -ImPyPy- β -PyPy
	2101 β) 5'-W G T A T A T W-3'	ImHp- β -HpPyHp- γ -PyHpPy- β -PyPy
	2102 β) 5'-W G T A T A A W-3'	ImHp- β -HpPyPy- γ -HpHpPy- β -PyPy
10	2103 β) 5'-W G T A T A G W-3'	ImHp- β -HpPyIm- γ -PyHpPy- β -PyPy
	2104 β) 5'-W G T A T A C W-3'	ImHp- β -HpPyPy- γ -ImHpPy- β -PyPy
	2105 β) 5'-W G T A T G T W-3'	ImHp- β -HpImHp- γ -PyPyPy- β -PyPy
	2106 β) 5'-W G T A T G A W-3'	ImHp- β -HpImPy- γ -HpPyPy- β -PyPy
	2107 β) 5'-W G T A T G G W-3'	ImHp- β -HpImIm- γ -PyPyPy- β -PyPy
	2108 β) 5'-W G T A T G C W-3'	ImHp- β -HpImPy- γ -ImPyPy- β -PyPy
	2109 β) 5'-W G T A T C T W-3'	ImHp- β -HpPyHp- γ -PyImPy- β -PyPy
	2110 β) 5'-W G T A T C A W-3'	ImHp- β -HpPyPy- γ -HpImPy- β -PyPy
	2111 β) 5'-W G T A T C G W-3'	ImHp- β -HpPyIm- γ -PyImPy- β -PyPy
	2112 β) 5'-W G T A T C C W-3'	ImHp- β -HpPyPy- γ -ImImPy- β -PyPy
	2113 β) 5'-W G T A A T T W-3'	ImHp- β -PyHpHp- γ -PyPyHp- β -PyPy
	2114 β) 5'-W G T A A T A W-3'	ImHp- β -PyHpPy- γ -HpPyHp- β -PyPy
	2115 β) 5'-W G T A A T G W-3'	ImHp- β -PyHpIm- γ -PyPyHp- β -PyPy
	2116 β) 5'-W G T A A T C W-3'	ImHp- β -PyHpPy- γ -ImPyHp- β -PyPy
	2117 β) 5'-W G T A A A T W-3'	ImHp- β -PyPyHp- γ -PyHpHp- β -PyPy
25	2118 β) 5'-W G T A A A A W-3'	ImHp- β -PyPyPy- γ -HpHpHp- β -PyPy
	2119 β) 5'-W G T A A A G W-3'	ImHp- β -PyPyIm- γ -PyHpHp- β -PyPy
	2120 β) 5'-W G T A A A C W-3'	ImHp- β -PyPyPy- γ -ImHpHp- β -PyPy
	2121 β) 5'-W G T A A G T W-3'	ImHp- β -PyImHp- γ -PyPyHp- β -PyPy
	2122 β) 5'-W G T A A G A W-3'	ImHp- β -PyImPy- γ -HpPyHp- β -PyPy
30	2123 β) 5'-W G T A A G G W-3'	ImHp- β -PyImIm- γ -PyPyHp- β -PyPy
	2124 β) 5'-W G T A A G C W-3'	ImHp- β -PyImPy- γ -ImPyHp- β -PyPy
	2125 β) 5'-W G T A A C T W-3'	ImHpPyPyPyHp- γ -PyImHp- β -PyPy
	2126 β) 5'-W G T A A C A W-3'	ImHpPyPyPyPy- γ -HpImHp- β -PyPy
	2127 β) 5'-W G T A A C G W-3'	ImHpPyPyPyIm- γ -PyImHp- β -PyPy
35	2128 β) 5'-W G T A A C C W-3'	ImHpPyPyPyPy- γ -ImImHp- β -PyPy

TABLE 177: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTASNNW-3'

	DNA sequence	aromatic amino acid sequence
	2129 β) 5'-W G T A G T T W-3'	ImHp- β -ImHpHp- γ -PyPyPy- β -PyPy
5	2130 β) 5'-W G T A G T A W-3'	ImHp- β -ImHpPy- γ -HpPyPy- β -PyPy
	2131 β) 5'-W G T A G T G W-3'	ImHp- β -ImHpIm- γ -PyPyPy- β -PyPy
	2132 β) 5'-W G T A G T C W-3'	ImHp- β -ImHpPy- γ -ImPyPy- β -PyPy
	2133 β) 5'-W G T A G A T W-3'	ImHp- β -ImPyHp- γ -PyHpPy- β -PyPy
	2134 β) 5'-W G T A G A A W-3'	ImHp- β -ImPyPy- γ -HpHpPy- β -PyPy
10	2135 β) 5'-W G T A G A G W-3'	ImHp- β -ImPyIm- γ -PyHpPy- β -PyPy
	2136 β) 5'-W G T A G A C W-3'	ImHp- β -ImPyPy- γ -ImHpPy- β -PyPy
	2137 β) 5'-W G T A G G T W-3'	ImHp- β -ImImHp- γ -PyPyPy- β -PyPy
	2138 β) 5'-W G T A G G A W-3'	ImHp- β -ImImPy- γ -HpPyPy- β -PyPy
	2139 β) 5'-W G T A G C T W-3'	ImHp- β -ImPyHp- γ -PyImPy- β -PyPy
	2140 β) 5'-W G T A G C A W-3'	ImHp- β -ImPyPy- γ -HpImPy- β -PyPy
	2141 β) 5'-W G T A G G G W-3'	ImHp- β -ImImIm- γ -PyPyPy- β -PyPy
	2142 β) 5'-W G T A G G C W-3'	ImHp- β -ImImPy- γ -ImPyPy- β -PyPy
	2143 β) 5'-W G T A G C G W-3'	ImHp- β -ImPyIm- γ -PyImPy- β -PyPy
	2144 β) 5'-W G T A G C C W-3'	ImHp- β -ImPyPy- γ -ImImPy- β -PyPy
20	2145 β) 5'-W G T A C T T W-3'	ImHp- β -PyHpHp- γ -PyPyIm- β -PyPy
	2146 β) 5'-W G T A C T A W-3'	ImHp- β -PyHpPy- γ -HpPyIm- β -PyPy
	2147 β) 5'-W G T A C T G W-3'	ImHp- β -PyHpIm- γ -PyPyIm- β -PyPy
	2148 β) 5'-W G T A C T C W-3'	ImHp- β -PyHpPy- γ -ImPyIm- β -PyPy
	2149 β) 5'-W G T A C A T W-3'	ImHp- β -PyPyHp- γ -PyHpIm- β -PyPy
25	2150 β) 5'-W G T A C A A W-3'	ImHp- β -PyPyPy- γ -HpHpIm- β -PyPy
	2151 β) 5'-W G T A C A G W-3'	ImHp- β -PyPyIm- γ -PyHpIm- β -PyPy
	2152 β) 5'-W G T A C A C W-3'	ImHp- β -PyPyPy- γ -ImHpIm- β -PyPy
	2153 β) 5'-W G T A C G T W-3'	ImHp- β -PyImHp- γ -PyPyIm- β -PyPy
	2154 β) 5'-W G T A C G A W-3'	ImHp- β -PyImPy- γ -HpPyIm- β -PyPy
30	2155 β) 5'-W G T A C C T W-3'	ImHp- β -PyPyHp- γ -PyImIm- β -PyPy
	2156 β) 5'-W G T A C C A W-3'	ImHp- β -PyPyPy- γ -HpImIm- β -PyPy
	2157 β) 5'-W G T A C G G W-3'	ImHp- β -PyImIm- γ -PyPyIm- β -PyPy
	2158 β) 5'-W G T A C G C W-3'	ImHp- β -PyImPy- γ -ImPyIm- β -PyPy
	2159 β) 5'-W G T A C C G W-3'	ImHp- β -PyPyIm- γ -PyImIm- β -PyPy
35	2160 β) 5'-W G T A C C C W-3'	ImHp- β -PyPyPy- γ -ImImIm- β -PyPy

TABLE 178: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTCWNNW-3'


	DNA sequence	aromatic amino acid sequence
	2161 β) 5'-W G T C T T T W-3'	ImHpPy- β -HpHp- γ -PyPy- β -ImPyPy
5	2162 β) 5'-W G T C T T A W-3'	ImHpPy- β -HpPy- γ -HpPy- β -ImPyPy
	2163 β) 5'-W G T C T T G W-3'	ImHpPy- β -HpIm- γ -PyPy- β -ImPyPy
	2164 β) 5'-W G T C T T C W-3'	ImHpPy- β -HpPy- γ -ImPy- β -ImPyPy
	2165 β) 5'-W G T C T A T W-3'	ImHpPy- β -PyHp- γ -PyHp- β -ImPyPy
	2166 β) 5'-W G T C T A A W-3'	ImHpPy- β -PyPy- γ -HpHp- β -ImPyPy
10	2167 β) 5'-W G T C T A G W-3'	ImHpPy- β -PyIm- γ -PyHp- β -ImPyPy
	2168 β) 5'-W G T C T A C W-3'	ImHpPy- β -PyPy- γ -ImHp- β -ImPyPy
	2169 β) 5'-W G T C T G T W-3'	ImHpPy- β -ImHp- γ -PyPy- β -ImPyPy
	2170 β) 5'-W G T C T G A W-3'	ImHpPy- β -ImPy- γ -HpPy- β -ImPyPy
	2171 β) 5'-W G T C T G G W-3'	ImHpPy- β -ImIm- γ -PyPy- β -ImPyPy
	2172 β) 5'-W G T C T G C W-3'	ImHpPy- β -ImPy- γ -ImPy- β -ImPyPy
	2173 β) 5'-W G T C T C T W-3'	ImHpPy- β -PyHp- γ -PyIm- β -ImPyPy
	2174 β) 5'-W G T C T C A W-3'	ImHpPy- β -PyPy- γ -HpIm- β -ImPyPy
	2175 β) 5'-W G T C T C G W-3'	ImHpPy- β -PyIm- γ -PyIm- β -ImPyPy
	2176 β) 5'-W G T C T C C W-3'	ImHpPy- β -PyPy- γ -ImIm- β -ImPyPy
	2177 β) 5'-W G T C A T T W-3'	ImHpPy- β -HpHp- γ -PyPy- β -ImPyPy
	2178 β) 5'-W G T C A T A W-3'	ImHpPy- β -HpPy- γ -HpPy- β -ImPyPy
	2179 β) 5'-W G T C A T G W-3'	ImHpPy- β -HpIm- γ -PyPy- β -ImPyPy
	2180 β) 5'-W G T C A T C W-3'	ImHpPy- β -HpPy- γ -ImPy- β -ImPyPy
	2181 β) 5'-W G T C A A T W-3'	ImHpPy- β -PyHp- γ -PyHp- β -ImPyPy
25	2182 β) 5'-W G T C A A A W-3'	ImHpPy- β -PyPy- γ -HpHp- β -ImPyPy
	2183 β) 5'-W G T C A A G W-3'	ImHpPy- β -PyIm- γ -PyHp- β -ImPyPy
	2184 β) 5'-W G T C A A C W-3'	ImHpPy- β -PyPy- γ -ImHp- β -ImPyPy
	2185 β) 5'-W G T C A G T W-3'	ImHpPy- β -ImHp- γ -PyPy- β -ImPyPy
	2186 β) 5'-W G T C A G A W-3'	ImHpPy- β -ImPy- γ -HpPy- β -ImPyPy
30	2187 β) 5'-W G T C A G G W-3'	ImHpPy- β -ImIm- γ -PyPy- β -ImPyPy
	2188 β) 5'-W G T C A G C W-3'	ImHpPy- β -ImPy- γ -ImPy- β -ImPyPy
	2189 β) 5'-W G T C A C T W-3'	ImHpPy- β -PyHp- γ -PyIm- β -ImPyPy
	2190 β) 5'-W G T C A C A W-3'	ImHpPy- β -PyPy- γ -HpIm- β -ImPyPy
	2191 β) 5'-W G T C A C G W-3'	ImHpPy- β -PyIm- γ -PyIm- β -ImPyPy
35	2192 β) 5'-W G T C A C C W-3'	ImHpPy- β -PyPy- γ -ImIm- β -ImPyPy

TABLE 179: 12-ring β -Hairpin Polyamides for recognition of 8-bp 5'-WGTCNNW-3'

	DNA sequence	aromatic amino acid sequence
	2193 β) 5'-W G T C G T T W-3'	ImHp- β -ImHpHp- γ -PyPy- β -ImPyPy
5	2194 β) 5'-W G T C G T A W-3'	ImHp- β -ImHpPy- γ -HpPy- β -ImPyPy
	2195 β) 5'-W G T C G T G W-3'	ImHp- β -ImHpIm- γ -PyPy- β -ImPyPy
	2196 β) 5'-W G T C G T C W-3'	ImHp- β -ImHpPy- γ -ImPy- β -ImPyPy
	2197 β) 5'-W G T C G A T W-3'	ImHp- β -ImPyHp- γ -PyHp- β -ImPyPy
	2198 β) 5'-W G T C G A A W-3'	ImHp- β -ImPyPy- γ -HpHp- β -ImPyPy
10	2199 β) 5'-W G T C G A G W-3'	ImHp- β -ImPyIm- γ -PyHp- β -ImPyPy
	2200 β) 5'-W G T C G A C W-3'	ImHp- β -ImPyPy- γ -ImHp- β -ImPyPy
	2201 β) 5'-W G T C G G T W-3'	ImHp- β -ImImHp- γ -PyPy- β -ImPyPy
	2202 β) 5'-W G T C G G A W-3'	ImHp- β -ImImPy- γ -HpPy- β -ImPyPy
	2203 β) 5'-W G T C G C T W-3'	ImHp- β -ImPyHp- γ -PyIm- β -ImPyPy
	2204 β) 5'-W G T C G C A W-3'	ImHp- β -ImPyPy- γ -HpIm- β -ImPyPy
	2205 β) 5'-W G T C C T T W-3'	ImHp- β -PyHpHp- γ -Py- β -ImImPyPy
	2206 β) 5'-W G T C C T A W-3'	ImHp- β -PyHpPy- γ -Hp- β -ImImPyPy
	2207 β) 5'-W G T C C T G W-3'	ImHp- β -PyHpIm- γ -Py- β -ImImPyPy
	2208 β) 5'-W G T C C T C W-3'	ImHp- β -PyHpPy- γ -Im- β -ImImPyPy
	2209 β) 5'-W G T C C A T W-3'	ImHp- β -PyPyHp- γ -Py- β -ImImPyPy
	2210 β) 5'-W G T C C A A W-3'	ImHp- β -PyPyPy- γ -Hp- β -ImImPyPy
	2211 β) 5'-W G T C C A G W-3'	ImHp- β -PyPyIm- γ -Py- β -ImImPyPy
	2212 β) 5'-W G T C C A C W-3'	ImHp- β -PyPyPy- γ -Im- β -ImImPyPy
	2213 β) 5'-W G T C C G T W-3'	ImHp- β -PyImHp- γ -Py- β -ImImPyPy
25	2214 β) 5'-W G T C C G A W-3'	ImHp- β -PyImPy- γ -Hp- β -ImImPyPy
	2215 β) 5'-W G T C C C T W-3'	ImHp- β -PyPyHp- γ -PyImImIm- β -Py
	2216 β) 5'-W G T C C C A W-3'	ImHp- β -PyPyPy- γ -HpImImIm- β -Py
	2217 β) 5'-W G T C G G G W-3'	ImHp- β -ImImIm- γ -PyPy- β -ImPyPy
	2218 β) 5'-W G T C G G C W-3'	ImHp- β -ImImPy- γ -ImPy- β -ImPyPy
30	2219 β) 5'-W G T C G C G W-3'	ImHp- β -ImPyIm- γ -PyIm- β -ImPyPy
	2220 β) 5'-W G T C G C C W-3'	ImHp- β -ImPyPy- γ -ImIm- β -ImPyPy
	2221 β) 5'-W G T C C G G W-3'	ImHp- β -PyImIm- γ -Py- β -ImImPyPy
	2222 β) 5'-W G T C C G C W-3'	ImHp- β -PyImPy- γ -Im- β -ImImPyPy
	2223 β) 5'-W G T C C C G W-3'	ImHp- β -PyPyIm- γ -PyImImIm- β -Py
35	2224 β) 5'-W G T C C C C W-3'	ImHp- β -PyPyPy- γ -ImImImIm- β -Py

What is claimed is:


1. A method for designing a specific polyamide



wherein X_1 , X_2 , X_m , $X_{(m+1)}$, $X_{(2m-1)}$, and X_{2m} are carboxamide residues forming carboxamide binding pairs X_1/X_{2m} , $X_2/X_{(2m-1)}$, $X_m/X_{(m+1)}$, and γ is γ -aminobutyric acid or 2,4 diaminobutyric acid and Dp is dimethylaminopropylamide, suitable for use as a DNA-binding ligand that is selective for identified target DNA sequences $5'-WN_1N_2 \dots N_mW-3'$ where m is an integer having a value from 3 to 6, comprising the steps of:

- a. identifying a target sequence of double stranded DNA having the form $5'-WN_1N_2 \dots N_mW-3'$, $N_1N_2 \dots N_m$ being the sequence to be bound by carboxamide residues, wherein each N is independently chosen from the group A, G, C, and T, each W is independently chosen from the group A and T, and m is an integer having a value from 3 to 6;
- b. representing the identified sequence as $5'-Wab \dots xW-3'$, wherein a is a first nucleotide to be bound by the X_1 carboxamide residue, b is a second nucleotide to be bound by the X_2 carboxamide residue, and x is the corresponding nucleotide to be bound by the X_m carboxamide residue;
- c. defining a as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified sequence;
- d. selecting Im as the X_1 carboxamide residue and Py as the X_{2m} carboxamide residue if $a = G$;
- e. selecting Py as the X_1 carboxamide residue and Im as the X_{2m} carboxamide residue if $a = C$;
- f. selecting Hp as the X_1 carboxamide residue and Py as the X_{2m} carboxamide residue if $a = T$;
- g. selecting Py as the X_1 carboxamide residue and Hp as the X_{2m} carboxamide residue if $a = A$; and
- h. repeating steps c – g for b through x until all carboxamide residues are selected.

2. The method of claim 1 further comprising the step of synthesizing the polyamide



3. The method of claim 2 further comprising the step of determining if the binding affinity of the polyamide to the identified sequence is subnanomolar.

4. The method of claim 2 further comprising the step of determining if the sequence specificity of the polyamide is greater or equal to ten.
5. The method of claim 2 further comprising the step of replacing at least one pyrrole residue with a β -alanine residue.

5 6. A method for designing a selective polyamide molecule $X_1X_2X_3X_4\text{-}\gamma\text{-}X_5X_6X_7X_8$, wherein $X_1, X_2, X_3, X_4, X_5, X_6, X_7$, and X_8 , are carboxamide residues forming binding pairs $X_1/X_8, X_2/X_7, X_3/X_6$ and X_4/X_5 , and γ is γ -aminobutyric acid or 2,4 diaminobutyric acid suitable for binding to a six base pair sequence of the form 5'-WNNNNW-3' in the minor groove of double stranded DNA, comprising the steps of:

- 10 a. identifying a six base pair sequence of double stranded DNA having the form 5'-WNNNNW-3', wherein W is either A or T, NNNN is the sequence to be bound by carboxamide residues, and each N is independently A, G, C, or T;
- b. representing the identified sequence as 5'-WabcdW-3', wherein *a* is a first nucleotide to be bound by a carboxamide residue, *b* is a second nucleotide to be bound by a carboxamide residue, *c* is a third nucleotide to be bound by a carboxamide residue, and *d* is a fourth nucleotide to be bound by a carboxamide residue;
- c. defining *a* as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
- d. selecting Im as the X_1 carboxamide residue and Py as the X_8 carboxamide residue if *a* = G;
- e. selecting Py as the X_1 carboxamide residue and Im as the X_8 carboxamide residue if *a* = C;
- f. selecting Hp as the X_1 carboxamide residue and Py as the X_8 carboxamide residue if *a* = T;
- 25 g. selecting Py as the X_1 carboxamide residue and Hp as the X_8 carboxamide residue if *a* = A;
- h. defining *b* as A, G, C, or T to correspond to the second nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
- i. selecting Im as the X_2 carboxamide residue and Py as the X_7 carboxamide residue if *b* = G;
- 30 j. selecting Py as the X_2 carboxamide residue and Im as the X_7 carboxamide residue if *b* = C;

- k. selecting Hp as the X₂ carboxamide residue and Py as the X₇ carboxamide residue if $b = T$;
- l. selecting Py as the X₂ carboxamide residue and Hp as the X₇ carboxamide residue if $b = A$;
- 5 m. defining c as A, G, C, or T to correspond to the third nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
- n. selecting Im as the X₃ carboxamide residue and Py as the X₆ carboxamide residue if $c = G$;
- 10 o. selecting Py as the X₃ carboxamide residue and Im as the X₆ carboxamide residue if $c = C$;
- p. selecting Hp as the X₃ carboxamide residue and Py as the X₆ carboxamide residue if $c = T$;
- q. selecting Py as the X₃ carboxamide residue and Hp as the X₆ carboxamide residue if $c = A$;
- r. defining d as A, G, C, or T to correspond to the fourth nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
- s. selecting Im as the X₄ carboxamide residue and Py as the X₅ carboxamide residue if $d = G$;
- 20 t. selecting Py as the X₄ carboxamide residue and Im as the X₅ carboxamide residue if $d = C$;
- u. selecting Hp as the X₄ carboxamide residue and Py as the X₅ carboxamide residue if $d = T$; and
- v. selecting Py as the X₄ carboxamide residue and Hp as the X₅ carboxamide residue if $d = A$.

- 25 7. The method of claim 6 further comprising the step of synthesizing the polyamide X₁X₂X₃X₄- γ -X₅X₆X₇X₈.
- 8. The method of claim 7 further comprising the step of determining if the binding affinity of the polyamide to the identified sequence is subnanomolar.
- 9. The method of claim 7 further comprising the step of determining if the sequence specificity of the polyamide is greater or equal to ten.
- 30 10. The method of claim 7 further comprising the step of replacing at least one pyrrole residue with a β -alanine residue at a position chosen from the group consisting of X₂, X₃, X₆, and X₇.

11. The method of claim 7 further comprising the step of replacing at least one 3-hydroxypyrrole residue with a β -alanine residue at a position chosen from the group consisting of X₂, X₃, X₆, and X₇.
12. A polyamide composition produced by the process comprising the steps of:
 - a. identifying a six base pair sequence of double stranded DNA having the form 5'-WNNNNW-3', wherein W is either A or T, NNNN is the sequence to be bound by carboxamide residues, and each N is independently A, G, C, or T;
 - b. representing the identified sequence as 5'-*Wabcd*W-3', wherein *a* is a first nucleotide to be bound by a carboxamide residue, *b* is a second nucleotide to be bound by a carboxamide residue, *c* is a third nucleotide to be bound by a carboxamide residue, and *d* is a fourth nucleotide to be bound by a carboxamide residue;
 - c. defining *a* as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
 - d. selecting Im as the X₁ carboxamide residue and Py as the X₈ carboxamide residue if *a* = G;
 - e. selecting Py as the X₁ carboxamide residue and Im as the X₈ carboxamide residue if *a* = C;
 - f. selecting Hp as the X₁ carboxamide residue and Py as the X₈ carboxamide residue if *a* = T;
 - g. selecting Py as the X₁ carboxamide residue and Hp as the X₈ carboxamide residue if *a* = A;
 - h. defining *b* as A, G, C, or T to correspond to the second nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;
 - i. selecting Im as the X₂ carboxamide residue and Py as the X₇ carboxamide residue if *b* = G;
 - j. selecting Py as the X₂ carboxamide residue and Im as the X₇ carboxamide residue if *b* = C;
 - k. selecting Hp as the X₂ carboxamide residue and Py as the X₇ carboxamide residue if *b* = T;
 - l. selecting Py as the X₂ carboxamide residue and Hp as the X₇ carboxamide residue if *b* = A;
 - m. defining *c* as A, G, C, or T to correspond to the third nucleotide to be bound by a carboxamide residue in the identified six base pair sequence;

- n. selecting Im as the X₃ carboxamide residue and Py as the X₆ carboxamide residue if $c = G$;
- o. selecting Py as the X₃ carboxamide residue and Im as the X₆ carboxamide residue if $c = C$;
- 5 p. selecting Hp as the X₃ carboxamide residue and Py as the X₆ carboxamide residue if $c = T$;
- q. selecting Py as the X₃ carboxamide residue and Hp as the X₆ carboxamide residue if $c = A$;
- r. defining d as A, G, C, or T to correspond to the fourth nucleotide to be bound by a
10 carboxamide residue in the identified six base pair sequence;
- s. selecting Im as the X₄ carboxamide residue and Py as the X₅ carboxamide residue if $d = G$;
- t. selecting Py as the X₄ carboxamide residue and Im as the X₅ carboxamide residue if $d = C$;
- u. selecting Hp as the X₄ carboxamide residue and Py as the X₅ carboxamide residue if $d = T$;
- v. selecting Py as the X₄ carboxamide residue and Hp as the X₅ carboxamide residue if $d = A$; and
- w. synthesizing the polyamide X₁X₂X₃X₄- γ -X₅X₆X₇X₈.

13. The polyamides described by the formulas listed in Tables 4 – 19.

14. The polyamides described by the formulas listed in Tables 20 – 83.

15. The polyamides described by the formulas listed in Tables 84 – 179.

16. A method for designing a selective polyamide molecule X₁X₂X₃X₄X₅- γ -X₆X₇X₈X₉X₁₀, wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, and X₁₀ are
25 carboxamide residues forming binding pairs X₁/X₁₀, X₂/X₉, X₃/X₈, X₄/X₇, and X₅/X₆, and γ is γ -aminobutyric acid or 2,4 diaminobutyric acid suitable for binding to a six base pair sequence of the form 5'-WNNNNNW-3' in the minor groove of double stranded DNA, comprising the steps of:

- a. identifying a seven base pair sequence of double stranded DNA having the form 5'-
30 WNNNNNW-3', wherein W is either A or T, NNNNN is the sequence to be bound by carboxamide residues, and each N is independently A, G, C, or T;
- b. representing the identified sequence as 5'-W**a***bcde*W-3', wherein **a** is a first nucleotide to be bound by a carboxamide residue, **b** is a second nucleotide to be


bound by a carboxamide residue, *c* is a third nucleotide to be bound by a carboxamide residue, *d* is a fourth nucleotide to be bound by a carboxamide residue, and *e* is a fifth nucleotide to be bound by a carboxamide residue;

- c. defining *a* as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified seven base pair sequence;
- d. selecting Im as the X₁ carboxamide residue and Py as the X₁₀ carboxamide residue if *a* = G;
- e. selecting Py as the X₁ carboxamide residue and Im as the X₁₀ carboxamide residue if *a* = C;
- f. selecting Hp as the X₁ carboxamide residue and Py as the X₁₀ carboxamide residue if *a* = T;
- g. selecting Py as the X₁ carboxamide residue and Hp as the X₁₀ carboxamide residue if *a* = A;
- h. defining *b* as A, G, C, or T to correspond to the second nucleotide to be bound by a carboxamide residue in the identified seven base pair sequence;
- i. selecting Im as the X₂ carboxamide residue and Py as the X₉ carboxamide residue if *b* = G;
- j. selecting Py as the X₂ carboxamide residue and Im as the X₉ carboxamide residue if *b* = C;
- k. selecting Hp as the X₂ carboxamide residue and Py as the X₉ carboxamide residue if *b* = T;
- l. selecting Py as the X₂ carboxamide residue and Hp as the X₉ carboxamide residue if *b* = A;
- m. defining *c* as A, G, C, or T to correspond to the third nucleotide to be bound by a carboxamide residue in the identified seven base pair sequence;
- n. selecting Im as the X₃ carboxamide residue and Py as the X₈ carboxamide residue if *c* = G;
- o. selecting Py as the X₃ carboxamide residue and Im as the X₈ carboxamide residue if *c* = C;
- p. selecting Hp as the X₃ carboxamide residue and Py as the X₈ carboxamide residue if *c* = T;
- q. selecting Py as the X₃ carboxamide residue and Hp as the X₈ carboxamide residue if *c* = A;

- r. defining d as A, G, C, or T to correspond to the fourth nucleotide to be bound by a carboxamide residue in the seven base pair sequence identified sequence;
- s. selecting Im as the X4 carboxamide residue and Py as the X7 carboxamide residue if $d = G$;
- 5 t. selecting Py as the X4 carboxamide residue and Im as the X7 carboxamide residue if $d = C$;
- u. selecting Hp as the X4 carboxamide residue and Py as the X7 carboxamide residue if $d = T$;
- v. selecting Py as the X4 carboxamide residue and Hp as the X7 carboxamide residue if $d = A$;
- 10 w. defining e as A, G, C, or T to correspond to the fifth nucleotide to be bound by a carboxamide residue in the seven base pair sequence identified sequence;
- x. selecting Im as the X5 carboxamide residue and Py as the X6 carboxamide residue if $e = G$;
- y. selecting Py as the X5 carboxamide residue and Im as the X6 carboxamide residue if $e = C$;
- z. selecting Hp as the X5 carboxamide residue and Py as the X6 carboxamide residue if $e = T$; and
- aa. selecting Py as the X5 carboxamide residue and Hp as the X6 carboxamide residue if $e = A$.

- 17. The method of claim 16 further comprising the step of synthesizing the polyamide $X_1X_2X_3X_4X_5\text{-}\gamma\text{-}X_6X_7X_8X_9X_{10}$.
- 18. The method of claim 17 further comprising the step of determining if the binding affinity of the polyamide to the identified sequence is subnanomolar.
- 19. The method of claim 17 further comprising the step of determining if the sequence specificity of the polyamide is greater or equal to ten.
- 20. The method of claim 17 further comprising the step of replacing at least one pyrrole residue with a β -alanine residue at a position chosen from the group consisting of X2, X3, X4, X7, X8, and X9.
- 21. The method of claim 17 further comprising the step of replacing at least one 3-hydroxypyrrole residue with a β -alanine residue at a position chosen from the group consisting of X2, X3, X4, X7, X8, and X9.

22. A polyamide composition produced by the method of claim 17.
23. A polyamide composition produced by the method of claim 18.
24. A polyamide composition produced by the method of claim 19.
25. A polyamide composition produced by the method of claim 20.
5 26. A polyamide composition produced by the method of claim 21.
27. A method for designing a selective polyamide molecule



wherein X_1 , X_2 , X_3 , X_4 , X_5 , X_6 , X_7 , X_8 , X_9 , X_{10} , X_{11} , and X_{12} , are
carboxamide residues forming binding pairs X_1/X_{12} , X_2/X_{11} , X_3/X_{10} , X_4/X_9 , X_5/X_8 ,
10 and X_6/X_7 , and γ is γ -aminobutyric acid or 2,4 diaminobutyric acid

suitable for binding to a eight base pair sequence of the form 5'-WNNNNNNW-
3' in the minor groove of double stranded DNA, comprising the steps of:


- a. identifying a eight base pair sequence of double stranded DNA having the form 5'-
WNNNNNNW-3', wherein W is either A or T, NNNNNN is the sequence to be
bound by carboxamide residues, and each N is independently A, G, C, or T;
b. representing the identified sequence as 5'-W**a***bcdef*W-3', wherein **a** is a first
nucleotide to be bound by a carboxamide residue, **b** is a second nucleotide to be
bound by a carboxamide residue, **c** is a third nucleotide to be bound by a
carboxamide residue, **d** is a fourth nucleotide to be bound by a carboxamide residue,
e is a fifth nucleotide to be bound by a carboxamide residue and **f** is a sixth
nucleotide to be bound by a carboxamide residue;
c. defining **a** as A, G, C, or T to correspond to the first nucleotide to be bound by a
carboxamide residue in the identified eight base pair sequence;
d. selecting Im as the X_1 carboxamide residue and Py as the X_{12} carboxamide residue
25 if **a** = G;
e. selecting Py as the X_1 carboxamide residue and Im as the X_{10} carboxamide residue
if **a** = C;
f. selecting Hp as the X_1 carboxamide residue and Py as the X_{12} carboxamide residue
if **a** = T;
30 g. selecting Py as the X_1 carboxamide residue and Hp as the X_{12} carboxamide residue
if **a** = A;
h. defining **b** as A, G, C, or T to correspond to the second nucleotide to be bound by a
carboxamide residue in the identified eight base pair sequence;

- i. selecting Im as the X₂ carboxamide residue and Py as the X₁₁ carboxamide residue if $b = G$;
- j. selecting Py as the X₂ carboxamide residue and Im as the X₁₁ carboxamide residue if $b = C$;
- 5 k. selecting Hp as the X₂ carboxamide residue and Py as the X₁₁ carboxamide residue if $b = T$;
- l. selecting Py as the X₂ carboxamide residue and Hp as the X₁₁ carboxamide residue if $b = A$;
- 10 m. defining c as A, G, C, or T to correspond to the third nucleotide to be bound by a carboxamide residue in the identified eight base pair sequence;
- n. selecting Im as the X₃ carboxamide residue and Py as the X₁₀ carboxamide residue if $c = G$;
- o. selecting Py as the X₃ carboxamide residue and Im as the X₁₀ carboxamide residue if $c = C$;
- p. selecting Hp as the X₃ carboxamide residue and Py as the X₁₀ carboxamide residue if $c = T$;
- q. selecting Py as the X₃ carboxamide residue and Hp as the X₁₀ carboxamide residue if $c = A$;
- 20 r. defining d as A, G, C, or T to correspond to the fourth nucleotide to be bound by a carboxamide residue in the eight base pair sequence identified sequence;
- s. selecting Im as the X₄ carboxamide residue and Py as the X₉ carboxamide residue if $d = G$;
- t. selecting Py as the X₄ carboxamide residue and Im as the X₉ carboxamide residue if $d = C$;
- 25 u. selecting Hp as the X₄ carboxamide residue and Py as the X₉ carboxamide residue if $d = T$;
- v. selecting Py as the X₄ carboxamide residue and Hp as the X₉ carboxamide residue if $d = A$;
- 30 w. defining e as A, G, C, or T to correspond to the fifth nucleotide to be bound by a carboxamide residue in the eight base pair sequence identified sequence;
- x. selecting Im as the X₅ carboxamide residue and Py as the X₈ carboxamide residue if $e = G$;

- y. selecting Py as the X5 carboxamide residue and Im as the X8 carboxamide residue if $e = C$;
- z. selecting Hp as the X5 carboxamide residue and Py as the X8 carboxamide residue if $e = T$;
- 5 aa. selecting Py as the X5 carboxamide residue and Hp as the X8 carboxamide residue if $e = A$;
- bb. defining f as A, G, C, or T to correspond to the sixth nucleotide to be bound by a carboxamide residue in the eight base pair sequence identified sequence;
- cc. selecting Im as the X6 carboxamide residue and Py as the X7 carboxamide residue if $f = G$;
- 10 dd. selecting Py as the X6 carboxamide residue and Im as the X7 carboxamide residue if $f = C$;
- ee. selecting Hp as the X6 carboxamide residue and Py as the X7 carboxamide residue if $f = T$; and
- ff. selecting Py as the X6 carboxamide residue and Hp as the X7 carboxamide residue if $f = A$.

- 28. The method of claim 17 further comprising the step of synthesizing the polyamide $X_1X_2X_3X_4X_5X_6\text{-}\gamma\text{-}X_7X_8X_9X_{10}X_{11}X_{12}$.
- 29. The method of claim 28 further comprising the step of determining if the binding affinity of the polyamide to the identified sequence is subnanomolar.
- 30. The method of claim 28 further comprising the step of determining if the sequence specificity of the polyamide is greater or equal to ten.
- 31. The method of claim 28 further comprising the step of replacing at least one pyrrole residue with a β -alanine residue at a position chosen from the group consisting of X_2 , X_3 , X_4 , X_5 , X_8 , X_9 , X_{10} , and X_{11} .
- 25 32. The method of claim 28 further comprising the step of replacing at least one 3-hydroxypyrrole residue with a β -alanine residue at a position chosen from the group consisting of X_2 , X_3 , X_4 , X_5 , X_8 , X_9 , X_{10} , and X_{11} .
- 33. A polyamide composition produced by the method of claim 28.
- 30 34. A polyamide composition produced by the method of claim 29.
- 35. A polyamide composition produced by the method of claim 30.
- 36. A polyamide composition produced by the method of claim 31.
- 37. A polyamide composition produced by the method of claim 32.

38. A polyamide composition produced by the method of claim 2 wherein one carboxamide binding pair is β/β .
39. A polyamide composition produced by the method of claim 7 wherein one carboxamide binding pair is β/β .
- 5 40. A polyamide composition produced by the method of claim 17 wherein one carboxamide binding pair is β/β .
41. A selective polyamide according to claim 1 whereby the polyamide is of the formula:



or a pharmaceutically acceptable salt wherein:

R^1 is chosen from H, NH_2 , SH, Cl, Br, F, N-acetyl, or N-formyl;

R^2 is chosen from H, $(CH_2)_mCH_3$, $(CH_2)_mNH_2$, $(CH_2)_mSH$, $(CH_2)_mOH$, $(CH_2)_mNR^5_2$, $(CH_2)_mOR^5$, $(CH_2)_mSR^5$, where $R^5 = (CH_2)_mCH_3$, $(CH_2)_mNH_2$, $(CH_2)_mSH$, $(CH_2)_mOH$ and m is an integer from 0 to 6;

R^3 is chosen from H, NH_2 , OH, SH, Br, Cl, F, OMe, CH_2OH , CH_2SH , CH_2NH_2 ;

R^4 is chosen from $-NH(CH_2)_{0-100}NR^6R^7$ or $NH(CH_2)_pCO NH(CH_2)_{0-100}NR^6R^7$ or NHR^6 or $NH(CH_2)_pCONHR^6$, where R^6 and R^7 are independently chosen from H, Cl, NO, N-acetyl, benzyl, C_{1-100} alkyl, C_{1-100} alkylamine, C_{1-100} alkyldiamine, C_{1-100} alkylcarboxylate, C_{1-100} alkenyl, a C_{1-100} alkynyl, or a $C_{1-100}L$, where L groups can be independently chosen from but is not limited to arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, oligodeoxynucleotide, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, an oligodeoxynucleotide, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- α -tocopheral;

where X and Y are chosen from the group consisting of N, CH, COH, CCH₃, CNH₂, CCl, CF;

a is an integer having values of 0 or 1;

b is an integer ranging from 1 to 5 inclusive; and

c is an integer value ranging from 2 to 10 inclusive.

42. The polyamide of claim 1 wherein the duplex DNA sequence is a regulatory sequence.

43. The polyamide of claim 1 wherein the duplex DNA sequence is a promoter sequence.

44. The polyamide of claim 1 wherein the duplex DNA sequence is a coding sequence.

45. The polyamide of claim 1 wherein the duplex DNA sequence is a non-coding sequence.

46. The polyamide of claim 1 wherein the binding of the carboxamide binding pairs to the identified target DNA sequence modulates the expression of a gene.

47. A composition comprising an effective amount of the polyamide of claim 1 and a pharmologically suitable excipient.

48. A diagnostic kit comprising the polyamide of claim 1.

ABSTRACT

The invention encompasses improved selective polyamides for binding to specific nucleotide sequences of double stranded DNA as well as methods for designing and synthesizing polyamide DNA binding ligands that are selective for an identified specific nucleotide sequence. The 3-hydroxy-N-methylpyrrole/N-methylpyrrole carboxamide pair specifically recognizes the T•A base pair, while the N-methylpyrrole/3-hydroxy-N-methylpyrrole pair recognizes A•T nucleotide pairs. Similarly, an N-methylimidazole/N-methylpyrrole carboxamide pair specifically recognizes the G•C nucleotide pair, and the N-methylpyrrole/N-methylimidazole carboxamide pair recognizes the C•G nucleotide pair.







FIG. 1

2/14




FIG. 2











FIG. 3

6-Ring Hairpin Hp-Py-Im-Polyamides



8-Ring Hairpin Hp-Py-Im-Polyamides



10-Ring Hairpin Hp-Py-Im-Polyamides






FIG. 4

5/14



Identify
5'-WNNNNW-3'
Sequence

define a

a = G

X₁ = Im
X₈ = Py

a = C

X₁ = Py
X₈ = Im

a = T

X₁ = Hp
X₈ = Py

a = A

X₁ = Py
X₈ = Hp

define b

b = G

X₂ = Im
X₇ = Py

b = C

X₂ = Py
X₇ = Im

b = T

X₂ = Hp
X₇ = Py

b = A

X₂ = Py
X₇ = Hp

define c

c = G

X₃ = Im
X₆ = Py

c = C

X₃ = Py
X₆ = Im

c = T

X₃ = Hp
X₆ = Py

c = A

X₃ = Py
X₆ = Hp

define d

d = G

X₄ = Im
X₅ = Py

d = C

X₄ = Py
X₅ = Im

d = T

X₄ = Hp
X₅ = Py

d = A

X₄ = Py
X₅ = Hp

Synthesize all-rng polyamide
X₁X₂X₃X₄-γ-X₅X₆X₇X₈

Determine Polyamide
Binding Affinity

subnanomolar
affinity

yes

no

> 10 fold
specificity

yes

no

End

A

FIG. 5

6/14




FIG. 6

7/14




FIG. 7a

8/14




FIG. 7b




FIG. 8




FIG. 9

11/14




FIG 10A

12/14




FIG 10B




FIG 11A




FIG 11B

**DECLARATION
AND POWER OF ATTORNEY
Utility Application**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **DESIGN, SYNTHESIS AND USE OF SPECIFIC POLYAMIDE DNA-BINDING LIGANDS**, the specification of which

(Check One) ☐ is attached hereto OR
☒ was filed on herewith, as United States Application Serial No.
to be assigned or PCT International Application No. PCT/US98/01714 and
was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Date of Filing	Priority Claimed	
			Yes	No
97/03332	PCT	February 20, 1997	X	
97/12722	PCT	July 21, 1997	X	

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date
60/038,384	February 14, 1997
60/023,309	July 31, 1996
60/024,374	August 1, 1996
60/026,713	September 25, 1996

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date	Status-Patented, Pending or Abandoned
08/853,522		May 8, 1997	Pending
08/837,524		April 21, 1997	Pending
08/607,078		February 26, 1996	Pending

POWER OF ATTORNEY: As a named inventor, I hereby appoint as my attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Roland N. Smoot, Reg. No. 18,718; Conrad R. Solum, Jr. Reg. No. 20,467; James W. Geriak, Reg. No. 20,233; Robert M. Taylor, Jr., Reg. No. 19,848; Samuel B. Stone, Reg. No. 19,297; Douglas E. Olson, Reg. No. 22,798; Robert E. Lyon, Reg. No. 24,171; Robert C. Weiss, Reg. No. 24,939; Richard E. Lyon, Jr., Reg. No. 26,300; John D. McConaghy, Reg. No. 26,773; William C. Steffin, Reg. No. 26,811; Coe A. Bloomberg, Reg. No. 26,605; J. Donald McCarthy, Reg. No. 25,119; John M. Benassi, Reg. No. 27,483; James H. Shalek, Reg. No. 29,749; Allan W. Jansen, Reg. No. 29,035; Robert W. Dickerson, Reg. No. 29,914; Roy L. Anderson, Reg. No. 30,240; David B. Murphy, Reg. No. 31,125; James C. Brooks, Reg. No. 29,898; Jeffrey M. Olson, Reg. No. 30,790; Steven D. Hemminger, Reg. No. 30,755; Jerrold B. Reilly, Reg. No. 32,293; Paul H. Meier, Reg. No. 32,274; John A. Rafter, Jr., Reg. No. 31,653; Kenneth H. Ohriner, Reg. No. 31,646; Mary S. Consalvi, Reg. No. 32,212; Lois M. Kwasigroch, Reg. No. 35,579; Lawrence R. LaPorte, Reg. No. 38,948; Robert C. Laurenson, Reg. No. 34,206; Bradford J. Duft, Reg. No. 32,219; Suzanne L. Biggs, Reg. No. 30,158; Richard J. Warburg, Reg. No. 32,327; John M. Johnson, Reg. No. 33,334; Daniel M. Chambers, Reg. No. 34,561; Troy M. Schmelzer, Reg. No. 36,667; Jessica R. Wolff, Reg. No. 37,261; Douglas C. Murdock, Reg. No. 37,549; Jeffrey W. Guise, Reg. No. 34,613; F.T. Alexandra Mahaney, Reg. No. 37,668; Charles S. Berkman, Reg. No. 38,077; Edward M. Jordan, Reg. No. 40,666; John C. Kappos, Reg. No. 37,861; Howard N. Wisnia, Reg. No. 37,502; Keith Kind, Reg. No. 42,735; Stephen S. Korniczky, Reg. No. 34,853; Jonathan T. Losk, Reg. No. 39,755; Theodore S. Maceiko, Reg. No. 35,593; Hope E. Melville, Reg. No. 34,874; Jeffrey A. Miller, Reg. No. 35,874; Kurt T. Mulville, Reg. No. 37,194; Clarke W. Neumann, Reg. No. 39,789; Vicki Norton, Reg. No. 40,745; Kenneth S. Roberts, Reg. No. 38,283; Thomas R. Rouse, Reg. No. 40,793; James K. Sakaguchi, Reg. No. 41,285; James K. Sakaguchi, Reg. No. 41,285; Carol A. Schneider, Reg. No. 34,923; Sheryl R. Silverstein, Reg. No. 40,812; Brent D. Sokol, Reg. No. 38,621; Jeffrey D. Tekanic, Reg. No. 36,031; Christopher A. Vanderlaan, Reg. No. 37,747; David E. Wang, Reg. No. 38,358; Lisa Ward Karmelich, Reg. No. 41,421; Michael J. Wise, Reg. No. 34,047; Wesley B. Ames, Reg. No. 40,893; Charles Balgenorth, Reg. No. 37,586; Thomas J. Brindisi, Reg. No. 40,348; James P. Brogan, Reg. No. 35,833; David T. Burse, Reg. No. 37,104; Bruce G. Chapman, Reg. No. 33,846; Farshad Farjami, Reg. No. 41,014; Charles C. Fowler, Reg. No. 39,675; and Corrine M. Freeman, Reg. No. 37,625.; of LYON & LYON, 633 West Fifth Street, Suite 4700, Los Angeles, California 90071-2066 telephone (619) 552-8400.

Send Correspondence to: Richard J. Warburg	LYON & LYON LLP 633 W Fifth St., Suite 4700 Los Angeles, CA 90071	Direct Telephone calls to: (619) 552-8400
---	---	--

	FULL NAME OF INVENTOR	FIRST Name Eldon	MIDDLE Initial E.	LAST Name Baird	
201	RESIDENCE & CITIZENSHIP	City Foster City	State or Foreign Country California	Country of Citizenship USA	
	POST OFFICE ADDRESS	311 Halyard Lane	City Foster City	State or Country California	Zip Code 94404
	FULL NAME OF INVENTOR	FIRST Name Peter	MIDDLE Initial B.	LAST Name Dervan	
202	RESIDENCE & CITIZENSHIP	City San Marino	State or Foreign Country California	Country of Citizenship USA	
	POST OFFICE ADDRESS	1235 St. Albans Road	City San Marino	State or Country California	Zip Code 91108
	FULL NAME OF INVENTOR	FIRST Name	MIDDLE Initial	LAST Name	
203	RESIDENCE & CITIZENSHIP	City	State or Foreign Country	Country of Citizenship	
	POST OFFICE ADDRESS		City	State or Country	Zip Code
	FULL NAME OF INVENTOR	FIRST Name	MIDDLE Initial	LAST Name	

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18, United States Code §1001, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor	201
Date	8-6-99
Signature of Inventor	202
Date	
Signature of Inventor	203
Date	

Signature of Inventor	204
Date	8-6-99
Signature of Inventor	205
Date	
Signature of Inventor	206
Date	

(Signatures should conform to names as presented at 201 et seq. above.)